arXiv:1210.0697v1 [math.LO] 2 Oct 2012

INSIDE THE MUCHNIK DEGREES:
DISCONTINUITY, LEARNABILITY, AND CONSTRUCTIVISM

KOJIRO HIGUCHI AND TAKAYUKI KIHARA

Abstract. It is known that infinitely many Medvedev degrees exist inside the Muchnik degree of any nontrivial Π^0_1 subset of Cantor space. We shed light on the fine structures inside these Muchnik degrees related to Learning Theory, Constructive Mathematics, and Computability Theory for discontinuous functions. First of all, we classify nonuniformly computable functions on Baire space from the viewpoint of Learning Theory, and then we investigate separation techniques to these classes in the strongest sense possible. By applying these techniques, among other results, we show that every nonzero Medvedev degree of Π^0_1 sets has the strong anti-cupping property, and that the Σ^0_2 law of excluded middle (Σ^0_2-LEM) is sufficient to transform each 3-valued diagonally noncomputable function (DNR_3) into a 2-valued diagonally noncomputable function (DNR_2), whereas neither the Σ^0_2 double negation elimination (Σ^0_2-DNE) nor the Σ^0_2 lessor limited principle of omniscience (Σ^0_2-LLPO) is sufficient. We also give characterizations of our classes of nonuniformly computable functions by piecewise computability, constructive principle, and Wadge games.

1. Summary

1.1. **Introduction.** To formulate a semantics for intuitionistic propositional calculus, Kolmogorov tried to interpret each proposition as a problem. Medvedev [49] formalized his idea by interpreting each proposition p as a mass problem $[\![p]\!] \subseteq \mathbb{N}^\mathbb{N}$. Under the interpretation, p is *provable* if $[\![p]\!] \neq \emptyset$ is witnessed algorithmically, i.e., $[\![p]\!]$ contains a computable element. Formally, $P \subseteq \mathbb{N}^\mathbb{N}$ *is Medvedev reducible to* $Q \subseteq \mathbb{N}^\mathbb{N}$ if there is a computable function $\Phi : Q \to P$. The degree structure induced by the Medvedev reduction forms a Brouwer algebra (the dual of a Heyting algebra). Therefore, for instance, the (intuitionistic) disjunction is interpreted as some binary operation on the subsets of Baire space which realizes the infimum in the Medvedev lattice.

The Medvedev lattice of nonempty Π^0_1 subsets (also called *effectively closed* or *co-c.e. closed* subsets) of Cantor space $2^\mathbb{N}$ has been widely studied from another perspective unrelated to non-classical logic (see Simpson [61]). In this context, Cenzer-Kihara-Weber-Wu [19] found the notion of tree-immunity for closed subsets of Cantor space $2^\mathbb{N}$. Their method to prove some theorems on tree-immunity gives rise to a kind of "disjunction" under the limit-BHK interpretation of *Limit Computable Mathematics* [36] (abbreviated as LCM), a kind of constructive mathematics based on Learning Theory, whose positive arithmetical fragment is captured by Heyting arithmetic with the recursive ω-rule and the Σ^0_1 law of excluded middle [6, 71]. Then the LCM-disjunctions are defined as binary operations on the subsets of Baire space. When disjunctive notions are represented as operations on the subsets of Baire space, this enable us to compare *degrees of difficulty* of disjunctive notions. This allows us to formalize the intuition that the intuitionistic disjunction is somehow *more difficult* than the classical one. We then introduce the notion of *relative learnability* for subsets of Baire space by defining the notion of learnability for functions on Baire space. They are useful for understanding how does degrees of difficulty of disjunctive notions behave.

Our another motivation is the development and study of computability of discontinuous functions. Imagine the floor function $\lfloor \cdot \rfloor : \mathbb{R} \to \mathbb{R}$ which is defined by $\lfloor x \rfloor = \max\{n \in \mathbb{Z} : n \leq x\}$. Although it

---

2010 *Mathematics Subject Classification.* 03D30, 03D78, 03F60, 03E15, 03B55, 68Q32.

Key words and phrases. Π^0_1 classes, computable analysis, limit computable mathematics, identification in the limit, constructive mathematics.

seems to be easy to draw a rough graph of the floor function $\lfloor\cdot\rfloor$, it is not computable since any computable function must be continuous. Such a *seemingly-computable* function is usually *nonuniformly computable*, where we say that a function f is nonuniformly computable if the output $f(x)$ is computable in a given input x. The degree structure induced by nonuniform computability is introduced by Muchnik [52], and nonuniformly computability naturally arises in Computable Analysis [12, 81]. Of course, most of nonuniformly computable discontinuous functions are far from being computable. Then, what kind of discontinuous functions are *nearly computable* such as the floor function or the step function? There have been several challenges [76, 77] to develop computability theory of (nonuniformly computable) discontinuous functions by using the notion of *learnability*. There are also a number of approaches to developing computability theory of (nonuniformly computable) discontinuous functions (see [15, 80, 79, 78, 81]). In accordance with their attempt and insight, we study various classes of nonuniformly computable functions.

This paper mainly deals with *computably-refutable cases*: In the situation that a scientist tries to verify a statement P, someday he can algorithmically find his verification to be refuted, whenever his verification is incorrect. This *refutability principle* holds only when P is represented as a Π^0_1 subset of the space. The restriction to Π^0_1 subsets of Cantor space $2^{\mathbb{N}}$ is motivated by several arguments. First of all, many mathematical problems can be represented as Π^0_1 subsets of certain topological spaces (see Cenzer and Remmel [22]). Indeed, the Π^0_1 sets in such spaces have become important notions in many branches of Computability Theory, such as *Recursive Mathematics* [31], *Reverse Mathematics* [64], *Computable Analysis* [75], *Effective Randomness* [55, 29], and *Effective Descriptive Set Theory* [50]. Secondly, the restriction to Π^0_1 sets can be regarded as an analogy of *the Popperian learning* [17] because of the refutability principle. From these perspective, the universe of Π^0_1 sets is expected to be a good playground of Learning Theory [38]. Thirdly, degree structures on Π^0_1 subsets of Cantor space $2^{\mathbb{N}}$ are widely studied from the viewpoint of *Computability Theory* and *Reverse Mathematics*. Cenzer and Hinman [20] showed that the Medvedev degrees of Π^0_1 subsets of Cantor space are dense. Then Simpson [61] asked whether the Muchnik degrees of Π^0_1 subsets of Cantor space are also dense. However, this question remains open. The reason for this might be that, even if A is Muchnik reducible to B (i.e., there is a nonuniformly computable function from B to A), this observation does not give us any knowledge about an algorithm solving A from a solution of B. Actually, as shown by Simpson-Slaman [66] and Cole-Simpson [25], there are infinitely many Medvedev degrees in the Muchnik degree of any nontrivial Π^0_1 subsets of Cantor space. Our learnability consideration for subsets of Baire space also appears to be in accordance with this point of view. In this paper, we shed light on a hidden, but extremely deep structure inside the Muchnik degree of each Π^0_1 subset of Cantor space.

1.2. **Results.** In section 2, we formalize the disjunction operations. Medvedev interpreted the intuitionistic disjunction as the direct sum $\oplus : \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \times \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$. We will introduce the following disjunction operations:

- (The LCM disjunction with mind-changes $< n+1$) $\triangledown_{n+1} : \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \times \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$;
- (The LCM disjunction) $\triangledown_{\omega} : \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \times \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$;
- (The classical disjunction) $\triangledown_{\infty} : \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \times \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$.

Then, the direct sum $\oplus$ is characterized as the LCM disjunction without mind-changes $\triangledown_1$. In the section, we will also introduce the notion of $(\alpha,\beta|\gamma)$-computability for partial functions on $\mathbb{N}^{\mathbb{N}}$, for each ordinal $\alpha,\beta,\gamma \leq \omega$. Then, the notion of $(\alpha,\beta|\gamma)$-computability induces just seven classes closed under composition.

- $[\mathfrak{C}_T]^1_1$ denotes the set of all partial computable functions on $\mathbb{N}^{\mathbb{N}}$.
- $[\mathfrak{C}_T]^1_{<\omega}$ denotes the set of all partial functions on $\mathbb{N}^{\mathbb{N}}$ learnable with bounded mind changes.
- $[\mathfrak{C}_T]^1_{\omega|<\omega}$ denotes the set of all partial functions on $\mathbb{N}^{\mathbb{N}}$ learnable with bounded errors.
- $[\mathfrak{C}_T]^1_{\omega}$ denotes the set of all partial learnable functions on $\mathbb{N}^{\mathbb{N}}$.

- $[\mathfrak{C}_T]^{<\omega}_1$ denotes the set of all partial k-wise computable functions on $\mathbb{N}^\mathbb{N}$ for some $k \in \mathbb{N}$.
- $[\mathfrak{C}_T]^{<\omega}_\omega$ denotes the set of all partial functions on $\mathbb{N}^\mathbb{N}$ learnable by a team.
- $[\mathfrak{C}_T]^{\omega}_1$ denotes the set of all partial nonuniformly computable functions on $\mathbb{N}^\mathbb{N}$ (i.e., all functions f satisfying $f(x) \leq_T x$ for any $x \in \mathrm{dom}(f)$).

These notions are also characterized as piecewise computability. We will see that the following inclusions hold.

$$\begin{tikzcd}[column sep=small, row sep=tiny]
 & & & {[\mathfrak{C}_T]^{<\omega}_1} \arrow[dr, phantom, sloped, "\subset"] & & \\
{[\mathtt{COMP}] = [\mathfrak{C}_T]^1_1} \arrow[r, phantom, "\subset"] & {[\mathfrak{C}_T]^1_{<\omega}} \arrow[r, phantom, "\subset"] & {[\mathfrak{C}_T]^1_{\omega|<\omega}} \arrow[ur, phantom, sloped, "\subset"] \arrow[dr, phantom, sloped, "\subset"] & & {[\mathfrak{C}_T]^{<\omega}_\omega} \arrow[r, phantom, "\subset"] & {[\mathfrak{C}_T]^\omega_1 = [\mathtt{NONU\ COMP}]} \\
 & & & {[\mathfrak{C}_T]^1_\omega} \arrow[ur, phantom, sloped, "\subset"] & &
\end{tikzcd}$$

Here, by [`COMP`] we denote the set of all partial computable functions on $\mathbb{N}^\mathbb{N}$, and by [`NONU COMP`] we denote the set of all partial nonuniformly computable functions on $\mathbb{N}^\mathbb{N}$. Our disjunction operations are useful to show that all of the above inclusions are proper. Beyond the properness of these inclusions, there are four LEVELs signifying the differences between two classes $\mathfrak{F}$ and $\mathfrak{G}$ of partial functions on $\mathbb{N}^\mathbb{N}$ (lying between [`COMP`] and [`NONU COMP`]).

(1) There is a function $\Gamma \in \mathfrak{F} \setminus \mathfrak{G}$.
(2) There are sets $X, Y \subseteq \mathbb{N}^\mathbb{N}$ such that $\mathfrak{F}$ has a function $\Gamma_\mathfrak{F} : X \to Y$, but $\mathfrak{G}$ has *no* function $\Gamma_\mathfrak{G} : X \to Y$.
(3) There are Π^0_1 sets $X, Y \subseteq 2^\mathbb{N}$ such that $\Gamma_\mathfrak{F}$ has a function $\Gamma_\mathfrak{F} : X \to Y$, but $\Gamma_\mathfrak{G}$ has *no* function $\Gamma_\mathfrak{G} : X \to Y$.
(4) For every special Π^0_1 set $Y \subseteq 2^\mathbb{N}$, there is a Π^0_1 set $X \subseteq 2^\mathbb{N}$ such that $\Gamma_\mathfrak{F}$ has a function $\Gamma_\mathfrak{F} : X \to Y$, but $\mathfrak{G}$ has *no* function $\Gamma_\mathfrak{G} : X \to Y$.

The LEVEL 1 separation just represents $\mathfrak{F} \not\subseteq \mathfrak{G}$. Clearly, $4 \to 3 \to 2 \to 1$. Note that the LEVEL 2 separation holds for *no* Σ^0_1 sets $X, Y \subseteq \mathbb{N}^\mathbb{N}$, since Π^0_1 is the first level in the arithmetical hierarchy which can define a nonempty set $S \subseteq \mathbb{N}^\mathbb{N}$ without computable element. Such a Π^0_1 set is called *special*, i.e., a subset of Baire space is special if it is nonempty and contains no computable points. Under Medvedev's interpretation, the specialness of $[\![p]\!]$ indicates that neither p nor $\neg p$ is provable (i.e., p is undecidable).

In section 3, we study interactions between the disjunction operations and the learnable functions. In this way, the LEVEL 3 separations are successful for $[\mathfrak{C}_T]^1_1$, $[\mathfrak{C}_T]^1_{<\omega}$, $[\mathfrak{C}_T]^1_{\omega|<\omega}$, and $[\mathfrak{C}_T]^{<\omega}_1$. We show that there exist Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that all of the following conditions are satisfied.

(1) (a) There is *no* computable function $\Gamma^1_1 : P \triangledown_2 Q \to P \triangledown_1 Q$;
    (b) There is a function $\Gamma^1_{<\omega} : P \triangledown_2 Q \to P \triangledown_1 Q$ learnable with bounded mind-changes.
(2) (a) There is *no* function $\Gamma^1_{<\omega} : P \triangledown_\omega Q \to P \triangledown_1 Q$ learnable with bounded mind-changes;
    (b) There is a function $\Gamma^1_{\omega|<\omega} : P \triangledown_\omega Q \to P \triangledown_1 Q$ learnable with bounded errors.
(3) (a) There is *no* function $\Gamma^1_{\omega|<\omega} : P \triangledown_\infty Q \to P \triangledown_1 Q$ learnable with bounded errors;
    (b) There is a k-wise computable function $\Gamma^{<\omega}_1 : P \triangledown_\infty Q \to P \triangledown_1 Q$ for some $k \in \mathbb{N}$.

The above conditions also suggest how does degrees of difficulty of our disjunction operations behave. For instance, (under Medvedev's interpretation) we cannot extract the proof of the intuitionistic disjunction of p and q from the proof of the bounded-mind-changes-LCM disjunction of p and q, even if these propositions p and q are interpreted as Π^0_1 sets.

In contrast to the above results, in section 4, we will see that the hierarchy between $[\mathfrak{C}_T]^1_{<\omega}$ and $[\mathfrak{C}_T]^{<\omega}_1$ collapses for *homogeneous* Π^0_1 subsets of Cantor space $2^\mathbb{N}$. In other words, the LEVEL 4 separations *fail* for $[\mathfrak{C}_T]^1_{<\omega}$, $[\mathfrak{C}_T]^1_{\omega|<\omega}$, and $[\mathfrak{C}_T]^{<\omega}_1$. For other classes, is the LEVEL 4 separation successful? In most parts of section 4, we develop techniques to archive the LEVEL 4 separations. In the section, we introduce many well-behaved operators $O : \mathcal{P}(\mathbb{N}^\mathbb{N}) \to \mathcal{P}(\mathbb{N}^\mathbb{N})$. The first operator is defined by $P \mapsto P^\frown P$, the concatenation of two closed sets $P \subseteq \mathbb{N}^\mathbb{N}$. Intuitively, the mass problem $P^\frown P$ means that "solve P by a learning proof process with mind-change-bound 2". By iterating the concatenation operation, we may introduce the mass problem which means that “solve P by a learning proof process (with a given mind-change-bound)”. Actually, we will introduce any transfinite iteration of the concatenation operation, where a learning proof process with an ordinal bounded mind changes is usually represented as *a learning proof process along a well-founded tree*. We will also introduce *a learning proof process along an ill-founded tree*, which will be called the *hyperconcatenation* $\blacktriangledown$ of closed sets. The hyperconcatenation $\blacktriangledown$ can be viewed as a special case of a team-learning proof process.

Our operators turn out to be extremely useful to establish the LEVEL 4 separation results. Some of these results will be proved by applying priority argument *inside* some learning proof model of P.

(1) The LEVEL 4 separation succeeds for $[\mathfrak{C}_T]^1_1$ and $[\mathfrak{C}_T]^1_{<\omega}$, via the map $P \mapsto P^\frown P$.
(2) The LEVEL 4 separation succeeds for $[\mathfrak{C}_T]^{<\omega}_1$ and $[\mathfrak{C}_T]^1_\omega$, via the map

$$P \mapsto \bigcup_{m\in\mathbb{N}} (P^\frown P^\frown \dots (m \text{ times}) \dots {}^\frown P^\frown P).$$

(3) The LEVEL 4 separation succeeds for $[\mathfrak{C}_T]^1_\omega$ and $[\mathfrak{C}_T]^{<\omega}_\omega$, via the map $P \mapsto P \blacktriangledown P$.
(4) The LEVEL 4 separation succeeds for $[\mathfrak{C}_T]^{<\omega}_\omega$ and $[\mathfrak{C}_T]^\omega_1$, via the map $P \mapsto \widehat{\mathrm{Deg}}(P)$, where $\widehat{\mathrm{Deg}}(P)$ denotes the Turing upward closure of P.

The method we use to show the first item also implies that every nonzero Medvedev degree of Π^0_1 subsets of Cantor space has *the strong anticupping property*, i.e., for every nonzero $\mathbf{a} \in \mathcal{P}$, there is a nonzero $\mathbf{b} \in \mathcal{P}$ below $\mathbf{a}$ such that $\mathbf{a} \not\leq \mathbf{b} \vee \mathbf{c}$ for any $\mathbf{a} \not\leq \mathbf{c}$, where the set of all Medvedev degrees of Π^0_1 subsets of Cantor space is denoted by $\mathcal{P}$. Under Medvedev’s interpretation of Intuitionism, our results seem to suggest, informally, that any Π^0_1-interpreted proposition p is not IPC-derivable from $p^\frown p \wedge q$ whenever p is not IPC-derivable from the Π^0_1-interpreted proposition q. Similarly, under Medvedev-style mass problem interpretation of LCM, any Π^0_1-interpreted proposition p is not LCM-derivable from $p \blacktriangledown p \wedge q$ whenever p is not LCM-derivable from the Π^0_1-interpreted proposition q. Here, IPC and LCM represent intuitionistic propositional caluculas and limit computable mathematics, respectively.

In section 5, we apply our results to sharpen Jockusch’s theorem [43] and Simpson’s Embedding Lemma [62]. Jockusch showed the following nonuniform computability result for DNR_k, the set of all k-valued *diagonally noncomputable functions*.

(1) There is *no* (uniformly) computable function $\Gamma^1_1 : \mathsf{DNR}_3 \to \mathsf{DNR}_2$.
(2) There is a nonuniformly computable function $\Gamma^\omega_1 : \mathsf{DNR}_3 \to \mathsf{DNR}_2$.

This result will be sharpened by using our learnability notions as follows.

(1) There is *no* learnable function $\Gamma^1_\omega : \mathsf{DNR}_3 \to \mathsf{DNR}_2$.
(2) There is *no* k-wise computable function $\Gamma^{<\omega}_1 : \mathsf{DNR}_3 \to \mathsf{DNR}_2$ for $k \in \mathbb{N}$.
(3) There is a (uniformly) computable function $\Gamma^1_1 : \mathsf{DNR}_3 \to \mathsf{DNR}_2 \blacktriangledown \mathsf{DNR}_2$. Hence, there is a function $\Gamma^{<\omega}_\omega : \mathsf{DNR}_3 \to \mathsf{DNR}_2$ learnable by a team of two learners.

In this section, we characterize our classes $[\mathfrak{C}_T]^\alpha_{\beta|\gamma}$ and operations in the context of constructive principle, Borel measurability, and topological games. We will see that how our classes of nonuniformly computable functions relate to the arithmetical hierarchy of non-intuitionistic principles such as *the law of excluded middle* (LEM), *the lessor limited principle of omniscience* or *de Morgan’s law* (LLPO), and *the double negation elimination* (DNE). The arithmetical hierarchy of non-intuitionistic principles is illustrated as follows:

$$\begin{tikzcd}
 & & \Sigma^0_2\text{-}\mathsf{LLPO} \arrow[dr, dash] & & \\
\Sigma^0_1\text{-}\mathsf{LEM} \arrow[r, dash] & \Delta^0_2\text{-}\mathsf{LEM} \arrow[ur, dash] \arrow[dr, dash] & & \Sigma^0_2\text{-}\mathsf{LEM} \arrow[r, dash] & \Sigma^0_3\text{-}\mathsf{DNE} \\
 & & \Sigma^0_2\text{-}\mathsf{DNE} \arrow[ur, dash] & &
\end{tikzcd}$$

Here, Γ-LEM represents the sentence $\varphi \vee \neg\varphi$ for Γ-sentences φ; Γ-LLPO represents the sentence $\neg(\varphi \wedge \psi) \to \neg\varphi \vee \neg\psi$ for Γ-sentences φ, ψ; and Γ-DNE represents the sentence $\neg\neg\varphi \to \varphi$ for Γ-sentences φ. Of course, in our context (i.e., in classical logic), these principles are provable. However, in some sense, these principles are not computable. For example, the concatenation and $[\mathfrak{C}_T]^1_{<\omega}$ corresponds to Σ^0_1-LEM; the bounded-error learnability $[\mathfrak{C}_T]^1_{\omega|<\omega}$ is related to Δ^0_2-LEM; the learnability $[\mathfrak{C}_T]^1_{\omega}$ corresponds to Σ^0_2-DNE; $[\mathfrak{C}_T]^{<\omega}_1$ is related to Σ^0_2-LLPO; the hyperconcatenation $\blacktriangledown$ is related to Σ^0_2-LEM; the nonuniform computability $[\mathfrak{C}_T]^{\omega}_1$ is related to Σ^0_3-DNE, and so on. Hence, every DNR_3-function can be transformed into a DNR_2-function whenever we are allowed to use Σ^0_2-LEM, whereas *neither* Σ^0_2-DNE *nor* Σ^0_2-LLPO is sufficient to transform each DNR_3-function into a DNR_2-function.

1.3. Notations and Conventions.

For any sets X and Y, we say that *f is a function from X to Y* (written $f : X \to Y$) if the domain $\mathrm{dom}(f)$ of f includes X, and the range $\mathrm{range}(f)$ of f is included in Y. We also use the notation $f :\subseteq X \to Y$ to denote that f is a partial function from X to Y, i.e., the domain $\mathrm{dom}(f)$ of f is included in X, and the range $\mathrm{range}(f)$ of f is also included in Y.

For basic terminology in Computability Theory, see Soare [67]. For $\sigma \in \mathbb{N}^{<\mathbb{N}}$, we let $|\sigma|$ denote the length of σ. For $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $f \in \mathbb{N}^{<\mathbb{N}} \cup \mathbb{N}^{\mathbb{N}}$, we say that σ is *an initial segment* of f (denoted by $\sigma \subset f$) if $\sigma(n) = f(n)$ for each $n < |\sigma|$. Moreover, $f \upharpoonright n$ denotes the unique initial segment of f of length n. let σ^- denote an immediate predecessor node of σ, i.e. $\sigma^- = \sigma \upharpoonright (|\sigma| - 1)$. We also define $[\sigma] = \{f \in \mathbb{N}^{\mathbb{N}} : f \supset \sigma\}$. A *tree* is a subset of $\mathbb{N}^{<\mathbb{N}}$ closed under taking initial segments. For any tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we also let $[T]$ be the set of all infinite paths of T, i.e., f belongs to $[T]$ if $f \upharpoonright n$ belongs to T for each $n \in \mathbb{N}$. A node $\sigma \in T$ is *extendible* if $[T] \cap [\sigma] \neq \emptyset$. Let T^{ext} denote the set of all extendible nodes of T. We say that $\sigma \in T$ is *a leaf* or *a dead end* if there is no $\tau \in T$ with $\tau \supsetneq \sigma$.

For any set X, the tree $X^{<\mathbb{N}}$ of finite words on X forms a monoid under concatenation $^\frown$. Here *the concatenation of σ and τ* is defined by $(\sigma^\frown\tau)(n) = \sigma(n)$ for $n < |\sigma|$ and $(\sigma^\frown\tau)(|\sigma| + n) = \tau(n)$ for $n < |\tau|$. We use symbols $^\frown$ and $\bigsqcap$ for the operation on this monoid, where $\bigsqcap_{i \leq n} \sigma_i$ denotes $\sigma_0{}^\frown\sigma_1{}^\frown \dots {}^\frown\sigma_n$. To avoid confusion, the symbols $\times$ and $\prod$ are only used for a product of sets. We often consider the following three left monoid actions of $X^{<\mathbb{N}}$: The first one is the set $X^{\mathbb{N}}$ of infinite words on X with an operation $^\frown : X^{<\mathbb{N}} \times X^{\mathbb{N}} \to X^{\mathbb{N}}$; $(\sigma^\frown f)(n) = \sigma(n)$ for $n < |\sigma|$ and $(\sigma^\frown f)(|\sigma| + n) = f(n)$ for $n \in \mathbb{N}$. The second one is the set $\mathcal{T}(X)$ of subtrees $T \subseteq X^{<\mathbb{N}}$ with an operation $^\frown : X^{<\mathbb{N}} \times \mathcal{T}(X) \to \mathcal{T}(X)$; $\sigma^\frown T = \{\sigma^\frown\tau : \tau \in T\}$. The third one is the power set $\mathcal{P}(X^{\mathbb{N}})$ of $X^{\mathbb{N}}$ with an operation $^\frown : X^{<\mathbb{N}} \times \mathcal{P}(X^{\mathbb{N}}) \to \mathcal{P}(X^{\mathbb{N}})$; $\sigma^\frown P = \{\sigma^\frown f : f \in P\}$.

We say that a set $P \subseteq \mathbb{N}^{\mathbb{N}}$ is Π^0_1 if there is a computable relation R such that $P = \{f \in \mathbb{N}^{\mathbb{N}} : (\forall n) R(n, f)\}$ holds. Equivalently, $P = [T_P]$ for some computable tree $T_P \subseteq \mathbb{N}^{\mathbb{N}}$. Let $\{\Phi_e\}_{e \in \mathbb{N}}$ be an effective enumeration of all partial computable functionals. Then the e-th Π^0_1 subset of $2^{\mathbb{N}}$ is defined by $P_e = \{f \in 2^{\mathbb{N}} : \Phi_e(f; 0) \uparrow\}$. Note that $\{P_e\}_{e \in \mathbb{N}}$ is an effective enumeration of all Π^0_1 subsets of Cantor space $2^{\mathbb{N}}$. If (an index e of) a Π^0_1 set $P_e \subseteq 2^{\mathbb{N}}$ is given, then $T_e = \{\sigma \in 2^{<\mathbb{N}} : \Phi_e(\sigma; 0) \uparrow\}$ is called *the corresponding tree for P_e*. Here $\Phi(\sigma; n)$ for $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $n \in \mathbb{N}$ denotes the computation of Φ with an oracle σ, an input n, and step $|\sigma|$. If $P \subseteq 2^{\mathbb{N}}$ is Π^0_1, then the corresponding tree $T_P \subseteq 2^{<\mathbb{N}}$ of P is computable, and $[T] = P$. Moreover, the set L_P of all leaves of the computable tree T_P is also computable. We also say that a sequence of $\{P_i\}_{i \in I}$ of Π^0_1 subsets of a space X is *computable* or *uniform* if the set $\{(i, f) \in I \times X : f \in P_i\}$ is again a Π^0_1 subset of the product space $I \times X$. A set $P \subseteq \mathbb{N}^{\mathbb{N}}$ is *special* if P is nonempty and P has no computable member. For $f, g \in \mathbb{N}^{\mathbb{N}}$, $f \oplus g$ is defined by $(f \oplus g)(2n) = f(n)$ and $(f \oplus g)(2n + 1) = g(n)$ for each $n \in \mathbb{N}$. For $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, put $P \oplus Q = (\langle 0 \rangle^\frown P) \cup (\langle 1 \rangle^\frown Q)$ and $P \otimes Q = \{f \oplus g : f \in P \& g \in Q\}$.

Contents

1. Summary 1
1.1. Introduction 1
1.2. Results 2
1.3. Notations and Conventions 5
2. Motivation and Terminology 7
2.1. Medvedev's Semantics for Intuitionism 7
2.2. Disjunction Operations Based on Learning Theory 8
2.3. Concatenations versus Disjunction Operations 13
2.4. Nonuniformly Computable Discontinuous Functions 15
2.5. Nonuniform Computability on Falsifiable Problems 21
3. Degrees of Difficulty of Disjunctions 24
3.1. The Disjunction $\oplus$ versus the Disjunction $\cup$ 24
3.2. The Disjunction $\cup$ versus the Disjunction $\triangledown$ 24
3.3. The Disjunction $\triangledown$ versus the Disjunction $\triangledown_\omega$ 25
3.4. The Disjunction $\triangledown_\omega$ versus the Disjunction $\triangledown_\infty$ 26
3.5. Infinite-Tape Models 26
3.6. The Disjunction $\bigoplus$ versus the Disjunction $\bigtriangledown_\infty$ 28
4. Contiguous Degrees and Dynamic Infinitary Disjunctions 31
4.1. When the Hierarchy Collapses 31
4.2. Dynamic Proof Models 33
4.3. Concatenation, Dynamic Disjunctions, and Contiguous Degrees 36
4.4. Infinitary Disjunctions along well-Founded Trees 39
4.5. Infinitary Disjunctions along the Straight Line 42
4.6. Infinitary Disjunctions along ill-Founded Trees 44
4.7. Infinitary Disjunctions along Nested ill-Founded Trees 48
4.8. Infinitary Disjunctions along Infinite Complete Graphs 54
5. Characterizations, Applications, and Questions 56
5.1. Characterization by Piecewise Computability 56
5.2. Diagonally Noncomputable Functions 64
5.3. Weihrauch Degrees and Constructive Principles 66
5.4. Duality between Dynamic Operations and Nonconstructive Principles 72
5.5. Some Intermediate Lattices are Not Brouwerian 75
5.6. Borel Measurability, and Backtrack Games 78
5.7. Open Questions 81
Bibliography 82
References 82

2. Motivation and Terminology

2.1. **Medvedev's Semantics for Intuitionism.**

To formulate a semantics for IPC, Kolmogorov tried to interpret each proposition as a problem. Medvedev [49] formalized his idea by interpreting each proposition p as a mass problem $[\![p]\!] \subseteq \mathbb{N}^{\mathbb{N}}$. Under the interpretation:

1. A *proof* π is a dynamical process represented by an infinite sequence of natural numbers, i.e., $\pi \in \mathbb{N}^{\mathbb{N}}$.
2. $[\![p]\!]$ is the set of all proofs of a proposition p, i.e., $[\![p]\!] \subseteq \mathbb{N}^{\mathbb{N}}$.
3. A proposition p is *provable* if p has a computable proof, i.e., $[\![p]\!] \subseteq \mathbb{N}^{\mathbb{N}}$ contains a computable element.

To prove the disjunction $p_0 \vee p_1$, we need to algorithmically decide which part is valid, i.e., we first declare one part to be valid and then construct a witness for this part. Consequently, $p_0 \vee p_1$ is provable under that interpretation if and only if we can algorithmically construct an element of $[\![p_0 \vee p_1]\!] = [\![p_0]\!] \oplus [\![p_1]\!] = \{\langle i\rangle^\frown f : i < 2 \ \& \ f \in [\![p_i]\!]\}$. Generally, let Form denote the all propositional formulas. Medvedev's idea is defining a mass-problem-interpretation of IPC by a function $[\![\cdot]\!] : \mathsf{Form} \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$ as in Definition 1.

Definition 1. We say that a function $[\![\cdot]\!] : \mathsf{Form} \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$ is *a Medvedev interpretation* if it satisfies the following six conditions.

1. $[\![\top]\!]$ contains a computable element.
2. $[\![\bot]\!] = \emptyset$.
3. $[\![\varphi \wedge \psi]\!] = [\![\varphi]\!] \otimes [\![\psi]\!] = \{f \oplus g : f \in [\![\varphi]\!] \ \& \ g \in [\![\psi]\!]\}$.
4. $[\![\varphi \vee \psi]\!] = [\![\varphi]\!] \oplus [\![\psi]\!] = \{\langle 0\rangle^\frown f : f \in [\![\varphi]\!]\} \cup \{\langle 1\rangle^\frown g : g \in [\![\psi]\!]\}$.
5. $[\![\varphi \to \psi]\!] = [\![\varphi]\!] \mathbf{\to} [\![\psi]\!] = \{e^\frown g \mid \Phi_e(g \oplus *) : [\![\varphi]\!] \to [\![\psi]\!]\}$.
6. $[\![\neg\varphi]\!] = [\![\varphi \to \bot]\!]$.

Here, $\Phi(g \oplus *)$ denotes the partial function $\lambda f.\Phi(g \oplus f) :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$, and recall that Φ_e is the e-th partial computable function on $\mathbb{N}^{\mathbb{N}}$.

Definition 2 (Medvedev [49])**.** For sets $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, *$P$ is Medvedev reducible to Q* (written $P \leq^1_1 Q$) if a computable function $\Phi : Q \to P$ exists. In other words, $P \leq^1_1 Q$ means the existence of a partial computable function Φ on Baire space $\mathbb{N}^{\mathbb{N}}$ with $\mathrm{dom}(\Phi) \supseteq Q$ and $\mathrm{rng}(\Phi) \subseteq P$. If both $P \leq^1_1 Q$ and $Q \leq^1_1 P$, then we say that *P is Medvedev equivalent to Q* (written $P \equiv^1_1 Q$).

If a lattice $(L, \leq, \vee, \wedge)$ has the top element 1, the bottom element 0, and $\max\{c : c \wedge a \leq b\}$ (denoted by $a \to_L b$) exists for any $a, b \in L$, then $\mathcal{L} = (L, \leq, \vee, \wedge, \to_L, 0, 1)$ is called a *Heyting algebra*. An algebra $\mathcal{L} = (L, \leq, \vee, \wedge, \to, \bot, \top)$ is *a Brouwer algebra* if its dual $\mathcal{L}^{\mathrm{op}} = (L, \geq, \wedge, \vee, \leftarrow, \top, \bot)$ is a Heyting algebra.

Theorem 3 (Medvedev [49])**.** *The quotient algebra $\mathcal{D}^1_1$ called the Medvedev lattice is Brouwerian. Here, $\mathcal{D}^1_1 = (\mathcal{P}(\mathbb{N}^{\mathbb{N}}), \leq^1_1, \otimes, \oplus, \mathbf{\to}, \mathbb{N}^{\mathbb{N}}, \emptyset)/\equiv^1_1$.*

Following Medvedev [49], Muchnik [52] introduced another calculus of mass problems. His calculus is non-uniform, but preserves computability in the following sense. A function $f :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is *nonuniformly computable* if $f(x) \leq_T x$ holds for any $x \in \mathrm{dom}(f)$, where the symbol $\leq_T$ denotes the Turing reducibility. Let $[\mathfrak{C}_T]^1_1$ denote the class of all computable functions $f :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$, and let $[\mathfrak{C}_T]^\omega_1$ denote the class of all nonuniformly computable functions $f :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$. Clearly, $[\mathfrak{C}_T]^1_1 \subsetneq [\mathfrak{C}_T]^\omega_1$. Actually, we have $\#[\mathfrak{C}_T]^\omega_1 = 2^{2^{\aleph_0}}$ while $[\mathfrak{C}_T]^1_1$ is countable.

Definition 4 (Muchnik [52])**.** For sets $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, *$P$ is Muchnik reducible to Q* (written $P \leq^\omega_1 Q$) if a nonuniformly computable function $\Phi : Q \to P$ exists. If both $P \leq^\omega_1 Q$ and $Q \leq^\omega_1 P$, then we say that *P is Muchnik equivalent to Q* (written $P \equiv^\omega_1 Q$).

Theorem 5 (Muchnik [52])**.** *The quotient algebra $\mathcal{D}^\omega_1$ called the Muchnik lattice is Brouwerian. Here, $\mathcal{D}^\omega_1 = (\mathcal{P}(\mathbb{N}^\mathbb{N}), \leq^\omega_1, \otimes, \oplus, \rightarrow', \mathbb{N}^\mathbb{N}, \emptyset)/ \equiv^\omega_1$ with the (dual) implication $P \rightarrow' Q = \{g \in \mathbb{N}^\mathbb{N} \mid (\exists \Gamma \in [\mathfrak{C}_T]^\omega_1)\ \Gamma(g \oplus *) : P \to Q\}$.*

Remark.

(1) Usually, the Medvedev reducibility is written as $\leq_M$ or $\leq_s$, and the Muchnik reducibility is written by $\leq_w$. However, later we will introduce the notion of (α,β)-reducibility for $\alpha,\beta \in \{1, <\omega, \omega\}$, and then the Medvedev reducibility and the Muchnik reducibility are treated as *the* $(1,1)$-*reducibility* and *the* $(\omega,1)$-*reducibility*, respectively. This is the reason why we use the notations $\leq^1_1$ and $\leq^\omega_1$ for the Medvedev reducibility and the Muchnik reducibility, respectively.

(2) Both of the Medvedev lattice $\mathcal{D}^1_1$ and the Muchnik lattice $\mathcal{D}^\omega_1$ provide sound and complete semantics for *Jankov's Logic* $\mathsf{KC} = \mathsf{IPC} + \neg p \vee \neg\neg p$, the intuitionistic propositional logic with *the weak law of excluded middle*, which is also called *De Morgan logic*. The Medvedev lattice and the Muchnik lattice are extensively studied from the aspect of Intermediate Logic. See Sorbi-Terwijn [69] and Hinman [37].

(3) Forty years after the pioneering work by Muchnik, the Muchnik reducibility become useful in the context of Reverse Mathematics (see Simpson [64]). The reason is that the Muchnik reducibility $\leq^\omega_1$ is strongly associated with the provability relation in RCA, *the recursive comprehension axiom*. Then, the Muchnik degrees of Π^0_1 subsets of $2^\mathbb{N}$ might be seen as instances of WKL, *the weak König's lemma*. For example, by using a result of Binns and Simpson [8] for the Muchnik degrees of Π^0_1 subsets of $2^\mathbb{N}$, Mummert [53] obtains an embedding theorem about the Lindenbaum algebra between RCA_0 and WKL_0.

(4) For more basic results about the Medvedev and Muchnik degrees of Π^0_1 subsets of $2^\mathbb{N}$, see Simpson [61, 62, 63, 65]. There are lots of research on the algebraic structure of the Medvedev degrees of Π^0_1 subsets of $2^\mathbb{N}$, such as density [20], embeddability of distributive lattices [8], join-reducibility [7], meet-irreducibility [2], noncuppability [19], decidability [24], and undecidability [60]. As a computable-analysistic approach of (Constructive) Reverse Mathematics, the Weihrauch degrees which is a common extension of the Medvedev degrees and the Wadge degrees is also studied [11, 13, 12].

2.2. **Disjunction Operations Based on Learning Theory.** Hayashi [35, 36] introduced *Limit Computable Mathematics* (LCM), an extended constructive mathematics based on *Learning Theory*. Like the BHK-interpretation for intuitionistic logic, there is a *limit-BHK interpratation* for Limit Computable Mathematics. We introduce three mass-problem-interpretations $[\![\cdot]\!]^i_{\mathsf{LCM}} : \mathsf{Form} \to \mathcal{P}(\mathbb{N}^\mathbb{N})$ of LCM based on the limit-BHK interpretation. To formulate a mass-problem-style interpretation of LCM, imagine the following *dynamic* proof models.

The one-tape model is defined as follows: When a verifier Ψ tries to prove that "P_0 or P_1", a tape Λ is given. At each stage, Ψ declares 0 or 1, and writes one letter on the tape Λ.

- **Intuitionism**: Ψ does not change his declaration, say $i \in \{0,1\}$, and the infinite word written on the tape Λ witnesses the validity of P_i.
- **LCM**: the sequence of declarations of Ψ converges, say $i \in \{0,1\}$, and the infinite word written on the tape Λ witnesses the validity of P_i.
- **Classical**: any declaration of Ψ is nonsense, and the infinite word written on the tape Λ witnesses the validity of P_0 or P_1.

The two-tape model is follows: When a verifier Ψ tries to prove "P_0 or P_1", two tapes Λ_0 and Λ_1 are given. At each stage, Ψ declares 0 or 1, say i, and he writes one letter on the tape Λ_i.

- **Intuitionism**: For either $i < 2$, the word written on Λ_{1-i} is empty, and the infinite word written on Λ_i witnesses the validity of P_i.

- **LCM**: For either $i < 2$, the word written on Λ_{1-i} is finite, and the infinite word written on Λ_i witnesses the validity of P_i.
- **Classical**: For either $i < 2$, the infinite word written on Λ_i witnesses the validity of P_i.

The backtrack-tape model is follows: When a verifier Ψ tries to prove that "P_0 or P_1", a paper $\square$, and two infinite tapes Λ, Δ are given. The paper $\square$ is called *the declaration*, Λ is called *the working tape*, and Δ is called *the record tape*. At each stage, the verifier Ψ works as follows.

1. If no letter is written on the declaration $\square$, then Ψ declares 0 or 1 and this is written on the declaration $\square$ and the record tape Δ.
2. When some letter is written on the declaration $\square$, the verifier Ψ chooses one letter k from $\mathbb{N} \cup \{\sharp\}$, and his choice k is written on the record tape Δ.
   (a) In the case $k \neq \sharp$, it expresses that Ψ writes the letter k on the working tape Λ.
   (b) In the case $k = \sharp$, it expresses that Ψ erases all letters from the declaration $\square$ and the working tape Λ.

- **Intuitionism**: Ψ does not choose $\sharp$, hence he does not change his declaration, say i, and the infinite word written on the tape Λ witnesses the validity of P_i.
- **LCM**: Ψ chooses $\sharp$ at most finitely often, hence the sequence of declarations of Ψ converges, say i, and the infinite word written on the tape Λ witnesses the validity of P_i.
- **Classical**: No classical counterpart.

To give formal definitions of these dynamic proof models, we introduce some auxiliary definitions.

Definition 6.

1. For $x = (x_0, x_1) \in I \times \mathbb{N}$, the first (second, resp.) coordinate x_0 (x_1, resp.) is denoted by $(x)_0$ ($(x)_1$, resp.). Fix $\sigma \in (I \times \mathbb{N})^{<\mathbb{N}}$, and $i \in I$. Then *the i-th projection of σ* is inductively defined as follows.
$$\mathtt{pr}_i(\langle\rangle) = \langle\rangle, \qquad \mathtt{pr}_i(\sigma) = \begin{cases} \mathtt{pr}_i(\sigma^-)^\frown n, & \text{if } \sigma = \sigma^{-\frown}\langle(i,n)\rangle, \\ \mathtt{pr}_i(\sigma^-), & \text{otherwise.} \end{cases}$$
Moreover, *the mind-change of σ* is given by $\mathtt{mc}(\sigma) = \#\{n < |\sigma| - 1 : (\sigma(n))_0 \neq (\sigma(n+1))_0\}$. Furthermore, for $f \in (I \times \mathbb{N})^{\mathbb{N}}$, we define $\mathtt{pr}_i(f) = \bigcup_{n \in \mathbb{N}} \mathtt{pr}_i(f \upharpoonright n)$ for each $i \in I$, and $\mathtt{mc}(f) = \lim_n \mathtt{mc}(f \upharpoonright n)$.
2. For any set X and string $\sigma \in X^{<\mathbb{N}}$, *the n-th shift* $\sigma^{\leftharpoonup n}$ is defined as $\sigma^{\leftharpoonup n}(m) = \sigma(n+m)$ for each $m < |\sigma| - n$. The *tail of σ* is defined as $\mathtt{tail}(\sigma) = \sigma^{\leftharpoonup m}$, for the least $m \in \mathbb{N}$ such that $\sigma(k) \neq \sharp$ for all $k \geq m$. Furthermore, for $f \in X^{\mathbb{N}}$, we define $f^{\leftharpoonup n} = \bigcup_{m \geq n}(f \upharpoonright m)^{\leftharpoonup n}$, and $\mathtt{tail}(f) = \lim_m \mathtt{tail}(f \upharpoonright m)$ if the limit exists.

Example 7. We consider two functions $\sigma \in (2 \times \mathbb{N})^{<\mathbb{N}}$ and $\tau \in (\mathbb{N} \cup \{\sharp\})^{<\mathbb{N}}$.

1. If $\sigma = \langle(1,3),(1,1),(0,4),(0,15),(1,9),(0,26),(0,5)\rangle$, then the projections of σ are $\mathtt{pr}_0(\sigma) = \langle 4, 15, 26, 5\rangle$, and $\mathtt{pr}_1(\sigma) = \langle 3, 1, 9\rangle$. Moreover, $\mathtt{mc}(\sigma) = 3$.
2. If $\tau = \langle 0,2,7,18,28,\sharp,1,8,2,8,45,9,\sharp,0,4,52,35,3,6\rangle$, then the tail of τ is $\mathtt{tail}(\tau) = \tau^{\leftharpoonup 13} = \langle 0,4,52,35,3,6\rangle$.

Definition 8 (One-Tape Disjunctions)**.** Let P_0 and P_1 be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.

1. $[\![P_0 \vee P_1]\!]^1_{\mathsf{Int}} = \bigcup_{i<2}(\{i^{\mathbb{N}}\} \otimes P_i)$.
2. $[\![P_0 \vee P_1]\!]^1_{\mathsf{LCM}} = \bigcup_{i<2}(\{f \in 2^{\mathbb{N}} : (\forall^\infty n)\ f(n) = i\} \otimes P_i)$.
3. $[\![P_0 \vee P_1]\!]^1_{\mathsf{CL}} = \bigcup_{i<2}(2^{\mathbb{N}} \otimes P_i)$.

Here, $i^{\mathbb{N}}$ denotes the infinite sequence consisting of i's, i.e., $i^{\mathbb{N}} = \langle i, i, i, \dots, i, i, i, \dots\rangle$.

Definition 9 (Two-Tape Disjunctions)**.** Let P_0 and P_1 be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.

1. $[\![P_0 \vee P_1]\!]^2_{\mathsf{Int}} = \{f \in (2 \times \mathbb{N})^{\mathbb{N}} : ((\exists i < 2)\ \mathtt{pr}_i(f) \in P_i)\ \&\ \mathtt{mc}(f) = 0\}$.
2. $[\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}} = \{f \in (2 \times \mathbb{N})^{\mathbb{N}} : ((\exists i < 2)\ \mathtt{pr}_i(f) \in P_i)\ \&\ \mathtt{mc}(f) < \infty\}$.

(3) $[\![P_0 \vee P_1]\!]^2_{\mathsf{CL}} = \{f \in (2 \times \mathbb{N})^{\mathbb{N}} : (\exists i < 2)\ \mathtt{pr}_i(f) \in P_i\}$.

Definition 10 (Backtrack Disjunctions)**.** Let P_0 and P_1 be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.

(1) $[\![P_0 \vee P_1]\!]^3_{\mathsf{Int}} = \{f \in (\mathbb{N} \cup \{\sharp\})^{\mathbb{N}} : \mathtt{tail}(f)^{\leftharpoonup 1} \in P_{\mathtt{tail}(f;0)}\ \&\ (\forall n)\ f(n) \neq \sharp\}$.
(2) $[\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}} = \{f \in (\mathbb{N} \cup \{\sharp\})^{\mathbb{N}} : \mathtt{tail}(f)^{\leftharpoonup 1} \in P_{\mathtt{tail}(f;0)}\ \&\ (\forall^{\infty} n)\ f(n) \neq \sharp\}$.

Remark. Note that we always have to choose a new symbol $\sharp$ which has not been already used. We consider the set S_t of all symbols which has been already used, where $S_0 = \mathbb{N}$. Then, each symbol $s \in S_t$ always codes $\langle 0, s\rangle$, and the new symbol $\sharp$ codes $\langle 1, 0\rangle$. The symbol $\sharp$ is enumerated into S_{t+1}, when we define some set $X \subseteq (S \cup \{\sharp\})^{\mathbb{N}}$. Here the symbol $S \cup \{\sharp\}$ means that $(\{0\} \times S) \cup \{\langle 1, 0\rangle\}$. Usually, both of S_t and $S_{t+1} = S_t \cup \{\sharp\}$ are coded as subsets of $\mathbb{N}$, by using a fixed bijection between $\mathbb{N}$ and $2 \times \mathbb{N}$.

Proposition 11. *Let P and Q be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) $[\![P \vee P]\!]^1_X \equiv^1_1 P$ *for each* $X \in \{\mathsf{Int}, \mathsf{LCM}, \mathsf{CL}\}$.
(2) $[\![P \vee Q]\!]^i_{\mathsf{CL}} \leq^1_1 [\![P \vee Q]\!]^i_{\mathsf{LCM}} \leq^1_1 [\![P \vee Q]\!]^i_{\mathsf{Int}}$ *for each* $i \in \{1, 2, 3\}$.
(3) $[\![P \vee Q]\!]^i_X \leq^1_1 [\![P \vee Q]\!]^j_X$ *for each* $j \leq i$ *and* $X \in \{\mathsf{Int}, \mathsf{LCM}, \mathsf{CL}\}$.
(4) $P \oplus Q \equiv^1_1 [\![P \vee Q]\!]^i_{\mathsf{Int}}$ *for each* $i \in \{1, 2, 3\}$.
(5) $P \cup Q \equiv^1_1 [\![P \vee Q]\!]^1_{\mathsf{CL}}$.

Proof. (1) The reduction $f \oplus g \mapsto g$ witnesses $P \leq^1_1 [\![P \vee P]\!]^1_X$, and the reduction $f \mapsto \mathtt{write}(0, f)$ witnesses $[\![P \vee P]\!]^1_X \leq^1_1 P$, for each $X \in \{\mathsf{Int}, \mathsf{LCM}, \mathsf{CL}\}$. Here, $\mathtt{write}(0, f)$ represents $\mathtt{write}(0, f) = 0^{\mathbb{N}} \oplus f = \langle (0, f(0)), (0, f(1)), (0, f(2)), \dots, (0, f(n)), (0, f(n+1)), \dots\rangle$ which indicates the instruction, in the one-tape model, to declare “P_0 is correct” at each stage and to write the infinite word f on the tape Λ.

(2) Clearly, $[\![P \vee Q]\!]^i_{\mathsf{CL}} \supseteq [\![P \vee Q]\!]^i_{\mathsf{LCM}} \supseteq [\![P \vee Q]\!]^i_{\mathsf{Int}}$ for each $i \in \{1, 2, 3\}$.

(3) Fix $X \in \{\mathsf{Int}, \mathsf{LCM}, \mathsf{CL}\}$. We inductively construct a computable function Ξ witnessing $[\![P \vee Q]\!]^2_X \leq^1_1 [\![P \vee Q]\!]^1_X$. First set $\Xi(\langle\rangle) = \langle\rangle$, and assume that $\Xi(\alpha \oplus \beta)$ has been already defined for every strings α and β of length s. Then we now define $\Xi(\sigma \oplus \tau)$ for each strings σ and τ of length $s+1$. For $p = |\mathtt{pr}_{\sigma(s)}(\Xi(\sigma^- \oplus \tau^-))|$ (recall that σ^- denotes the immediate predecessor of σ), we put $\Xi(\sigma \oplus \tau) = \Xi(\sigma^- \oplus \tau^-)^\frown \mathtt{write}(\sigma(s), \tau^{\leftharpoonup p})$. Here, $\mathtt{write}(i, \alpha)$ for $i \in I$ and $\alpha \in \mathbb{N}^{<\mathbb{N}}$ is defined as $i^{|\alpha|} \oplus \alpha = \langle (i, \alpha(0)), (i, \alpha(1)), \dots, (i, \alpha(|\alpha| - 1))\rangle$, i.e., $\mathtt{write}(i, \alpha)$ indicates the instruction to write the finite word α on the i-th tape Λ_i in the two-tape model. Then, we can inductively ensure the following condition.

$$\mathtt{pr}_{\sigma(s)}(\Xi(\sigma \oplus \tau)) = \mathtt{pr}_{\sigma(s)}(\Xi(\sigma^- \oplus \tau^-))^\frown(\tau^{\leftharpoonup p}) = \tau.$$

Finally, we set $\Xi(f \oplus g) = \bigcup_{n \in \mathbb{N}} \Xi((f \upharpoonright n) \oplus (g \upharpoonright n))$, for any $f, g \in \mathbb{N}^{\mathbb{N}}$. Therefore, for any $f \oplus g \in [\![P \vee Q]\!]^1_X$ and each $i < 2$, if $f(n) = i$ for infinitely many $n \in \mathbb{N}$, then $\mathtt{pr}_i(\Xi(f \oplus g))$ is total, and $\mathtt{pr}_i(\Xi(f \oplus g)) = g$. By definition, $\mathtt{pr}_i(\Xi(f \oplus g)) = g \in P_i$ for some $i < 2$. Hence, $\Xi(f \oplus g) \in [\![P \vee Q]\!]^2_X$.

Fix $X \in \{\mathsf{Int}, \mathsf{LCM}\}$. We inductively construct a computable function Ξ witnessing $[\![P \vee Q]\!]^3_X \leq^1_1 [\![P \vee Q]\!]^2_X$. First set $\Xi(\langle (i, n)\rangle) = \langle i, n\rangle$ for each $(i, n) \in 2 \times \mathbb{N}$. Fix $\sigma = \sigma^{--\frown}\langle (i, m), (j, n)\rangle \in (2 \times \mathbb{N})^{<\mathbb{N}}$, and assume that $\Xi(\sigma^-)$ has been already defined. Then, let us define $\Xi(\sigma)$ as follows:

$$\Xi(\sigma^{--\frown}\langle (i, m), (j, n)\rangle) = \begin{cases} \Xi(\sigma^-)^\frown\langle n\rangle & \text{if } j = i; \\ \Xi(\sigma^-)^\frown\langle \sharp, j\rangle^\frown \mathtt{pr}_j(\sigma) & \text{otherwise.} \end{cases}$$

Finally set $\Xi(f) = \bigcup_n \Xi(f \upharpoonright n)$, for any $f \in (2 \times \mathbb{N})^{\mathbb{N}}$. It is easy to see that $\mathtt{tail}(f)$ is defined for any $f \in [\![P \vee Q]\!]^2_X$, since $\#\{k \in \mathbb{N} : \Xi(f; k) = \sharp\} = \mathtt{mc}(f)$. Therefore, $\mathtt{tail}^{\leftharpoonup 1}(\Xi(f)) \in P_{\mathtt{tail}(\Xi(f);0)}$. If $X = \mathsf{Int}$, then no $\sharp$ occurs in $\Xi(f)$.

(4) By definition, $[\![P \vee Q]\!]^3_{\mathsf{Int}} = P \oplus Q$. (5) The reduction $f \oplus g \mapsto g$ witnesses $P \cup Q \leq^1_1 [\![P \vee Q]\!]^1_{\mathsf{CL}}$, and the reduction $f \mapsto \mathtt{write}(0, f) = 0^{\mathbb{N}} \oplus f$ witnesses $[\![P \vee Q]\!]^1_{\mathsf{CL}} \leq^1_1 P \cup Q$. □

Definition 12. For each proof model, there are variations of LCM disjunctions, for any *bound of mind changes*. Let P_0, P_1 be any subset of Baire space $\mathbb{N}^\mathbb{N}$, and n be any natural number.

(1) *The one-tape* LCM *disjunction of P_0 and P_1 with mind-changes-bound n* is defined as follows.

$$[\![P_0 \vee P_1]\!]^1_{\mathsf{LCM}[n]} = [\![P_0 \vee P_1]\!]^1_{\mathsf{LCM}} \cap \{f \in 2^\mathbb{N} : \#\{n \in \mathbb{N} : f(n+1) \neq f(n)\} < n\} \otimes 2^\mathbb{N}.$$

(2) *The two-tape* LCM *disjunction of P_0 and P_1 with mind-changes-bound n* is defined as follows.

$$[\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}[n]} = [\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}} \cap \{f \in (2 \times \mathbb{N})^\mathbb{N} : \mathtt{mc}(f) < n\}.$$

(3) *The backtrack-tape* LCM *disjunction of P_0 and P_1 with mind-changes-bound n* is defined as follows.

$$[\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}[n]} = [\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}} \cap \{f \in (\mathbb{N} \cup \{\sharp\})^\mathbb{N} : \#\{k \in \mathbb{N} : f(k) = \sharp\} < n\}.$$

Proposition 13. *Let P, Q be subsets of Baire space $\mathbb{N}^\mathbb{N}$.*

(1) $P \oplus Q \equiv^1_1 [\![P \vee Q]\!]^i_{\mathsf{LCM}[1]}$ *for each $i \in \{2,3\}$.*

(2) $[\![P \vee P]\!]^2_{\mathsf{LCM}[2]} \equiv^1_1 [\![P \vee P]\!]^3_{\mathsf{LCM}[2]}$. *Indeed,* $[\![\bigvee_{i<n} P_i]\!]^2_{\mathsf{LCM}[n]} \equiv^1_1 [\![P \vee P]\!]^3_{\mathsf{LCM}[n]}$, *where $P_i = P$ for each $i < n$. Here, for each collection $\{P_i\}_{i<k}$ of subsets of Baire space,* $[\![\bigvee_{i<k} P_i]\!]^2_{\mathsf{LCM}[n]}$ *is defined as follows.*

$$\{f \in (k \times \mathbb{N})^\mathbb{N} : ((\exists i < k)\ \mathtt{pr}_i(f) \in P_i)\ \&\ \mathtt{mc}(f) < n\}.$$

Proof. (1) Clearly $[\![P \vee Q]\!]^i_{\mathsf{LCM}[1]} = [\![P \vee Q]\!]^i_{\mathsf{Int}}$ for each $i \in \{1,2,3\}$. By Proposition 11 (4), we have $P \oplus Q \equiv^1_1 [\![P \vee Q]\!]^i_{\mathsf{Int}}$.

(2) The reduction $\Xi : h \mapsto h^*$ in the proof of Proposition 11 (3) also witnesses $[\![P \vee P]\!]^3_{\mathsf{LCM}[n]} \leq^1_1 [\![\bigvee_{i<n} P_i]\!]^2_{\mathsf{LCM}[n]}$. We inductively define a computable function Ξ^* witnessing $[\![\bigvee_{i<n} P_i]\!]^2_{\mathsf{LCM}[n]} \leq^1_1 [\![P \vee P]\!]^3_{\mathsf{LCM}[n]}$. Put $\Xi^*(\langle\rangle) = \langle\rangle$, and fix $\sigma = \sigma^{-}{}^\frown\langle k\rangle \in (\mathbb{N} \cup \{\sharp\})^{<\mathbb{N}}$. Assume that $\Xi^*(\sigma^-)$ has been already defined. Then, $\Xi^*(\sigma)$ is defined as follows.

$$\mathtt{count}(\sigma) = \#\{m < |\sigma| : \sigma(m) = \sharp\},$$

$$\Xi^*(\sigma^{-}{}^\frown\langle k\rangle) = \begin{cases} \Xi^*(\sigma^-)^\frown\langle(\mathtt{count}(\sigma), k)\rangle & \text{if } k \neq \sharp, \\ \Xi^*(\sigma^-) & \text{otherwise.} \end{cases}$$

For any $g \in [\![P \vee P]\!]^3_{\mathsf{LCM}[n]}$, we have $\mathtt{count}(g \upharpoonright s) < n$ for any $s \in \mathbb{N}$, and hence $\mathtt{mc}(\Xi^*(g)) < n$, since g contains at most n many $\sharp$'s. Moreover, $\mathtt{pr}_{\lim_s \mathtt{count}(g \upharpoonright s)}(\Xi^*(g)) = \mathtt{tail}(g)^{\leftarrow 1} \in P$. □

Proposition 14. *Let P_0, P_1, Q_0, and Q_1 be subsets of Baire space $\mathbb{N}^\mathbb{N}$, and fix $i \in \{2,3\}$ and $X \in \{\mathsf{Int}, \mathsf{LCM}, \mathsf{CL}\} \cup \{\mathsf{LCM}[n] : n \in \mathbb{N}\}$. If $P_0 \leq^1_1 Q_0$ and $P_1 \leq^1_1 Q_1$, then $[\![P_0 \vee P_1]\!]^i_X \leq^1_1 [\![Q_0 \vee Q_1]\!]^i_X$. Hence, the operator $\mathbf{D}^i_X : \mathcal{D}^1_1 \times \mathcal{D}^1_1 \to \mathcal{D}^1_1$ introduced by $\mathbf{D}^i_X(\deg^1_1(P), \deg^1_1(Q)) = \deg^1_1([\![P \vee Q]\!]^i_X)$ is well-defined. Here, $\deg^1_1(P)$ denotes the equivalent class $\{R \subseteq \mathbb{N}^\mathbb{N} : R \equiv^1_1 P\}$.*

Proof. We first consider the two-tape model. Assume that $P_0 \leq^1_1 Q_0$ and $P_1 \leq^1_1 Q_1$ via computable functions Γ_0 and Γ_1, respectively. We construct a computable function Δ witnessing $[\![P_0 \vee P_1]\!]^2_X \leq^1_1 [\![Q_0 \vee Q_1]\!]^2_X$. Set $\Delta(\langle\rangle)$. Fix $\sigma \in (2 \times \mathbb{N})^{<\mathbb{N}}$ and assume that $\Delta(\sigma^-)$ has been already defined. For each $i < 2$, we define $\eta_i \in \mathbb{N}^{<\mathbb{N}}$ as follows. If $\Gamma_i(\mathtt{pr}_i(\sigma)) \supsetneq \Gamma_i(\mathtt{pr}_i(\sigma^-))$, say $\Gamma_i(\mathtt{pr}_i(\sigma)) = \Gamma_i(\mathtt{pr}_i(\sigma^-))^\frown\eta$, then we set $\eta_i = \mathtt{write}(i, \eta)$ for such a string η, where recall $\mathtt{write}(i, \eta) = i^{|\eta|} \oplus \eta = \langle(i, \eta(0)), (i, \eta(1)), \dots, (i, \eta(|\eta| - 1))\rangle$. If $\Gamma_i(\mathtt{pr}_i(\sigma)) = \Gamma_i(\mathtt{pr}_i(\sigma^-))$, then we set $\eta_i = \langle\rangle$. Then we define $\Delta(\sigma) = \Delta(\sigma^-)^\frown\eta_0{}^\frown\eta_1$. Note that $\eta_0 = \langle\rangle$ or $\eta_1 = \langle\rangle$, since $\mathtt{pr}_i(\sigma) = \mathtt{pr}_i(\sigma^-)$ for either $i < 2$. Therefore, $\mathtt{mc}(\Delta(g)) = \mathtt{mc}(g)$ for any $g \in \mathbb{N}^\mathbb{N}$. Furthermore, for any $g \in \mathbb{N}^\mathbb{N}$, we have $\mathtt{pr}_i(\Delta(g)) = \Gamma_i(\mathtt{pr}_i(g))$ for each $i < 2$. Thus, $\Delta(g) \in [\![P_0 \vee P_1]\!]^2_X$ for any $g \in [\![Q_0 \vee Q_1]\!]^2_X$.

Next we consider the backtrack-tape model. Assume that $P_0 \leq^1_1 Q_0$ and $P_1 \leq^1_1 Q_1$ via computable functions Γ_0 and Γ_1, respectively. We construct a computable function Θ witnessing $[\![P_0 \vee P_1]\!]^3_X \leq^1_1$

$[\![Q_0 \vee Q_1]\!]^3_X$. Set $\Theta(\langle\rangle)$. Fix $\sigma \in (2\times\mathbb{N})^{<\mathbb{N}}$ and assume that $\Theta(\tau)$ has been already defined for each $\tau \subsetneq \sigma$. If $\sigma = \sigma^{--\frown}\langle m, n\rangle$ for some $m, n \in \mathbb{N}$, then we have $\Gamma_{\mathtt{tail}(\sigma;0)}(\mathtt{tail}(\sigma)^{\leftharpoonup 1}) = \Gamma_{\mathtt{tail}(\sigma;0)}(\mathtt{tail}(\sigma)^{\leftharpoonup 1})^\frown\eta$ for some $\eta \in \mathbb{N}^{<\mathbb{N}}$, and we define $\Theta(\sigma) = \Theta(\sigma^-)^\frown\eta$. If $\sigma = \sigma^{--\frown}\langle\sharp, i\rangle$ for some $i < 2$, i.e., $\mathtt{tail}(\sigma; 0) = i$, then define $\Theta(\sigma) = \Theta(\sigma^-)^\frown\langle\sharp, i\rangle$. Otherwise, we set $\Theta(\sigma) = \Theta(\sigma^-)$. Note that $\#\{n \in \mathbb{N} : \Theta(g; n) = \sharp\} = \#\{n \in \mathbb{N} : g(n) = \sharp\}$ for any $g \in \mathbb{N}^{\mathbb{N}}$. Furthermore, $\mathtt{tail}(\Theta(g); 0) = \mathtt{tail}(g; 0)$, and $\mathtt{tail}(\Theta(g))^{\leftharpoonup 1} = \Gamma_{\mathtt{tail}(g;0)}(\mathtt{tail}(g)^{\leftharpoonup 1})$ for any $g \in [\![Q_0 \vee Q_1]\!]^3_X$. Hence, $\Theta(g) \in [\![P_0 \vee P_1]\!]^3_X$ for any $g \in [\![Q_0 \vee Q_1]\!]^3_X$. □

Remark. The one-tape disjunctions are hard to analyze and are useless in our study, because they are badly behaved in the sense of Proposition 14. Indeed, by analyzing the proofs of Proposition 53 and Corollary 54, it is easy to see that there is a Π^0_1 subset P of Cantor space such that $[\![P \vee P]\!]^1_X \not\equiv^1_1 [\![P \vee P^*]\!]^1_X$ for each $X \in \{\mathsf{LCM}, \mathsf{CL}\}$, where $P^* = \{g \oplus f : f \oplus g \in P\}$. Though the original limit-BHK interpretation of the disjunctive notion seems to be a one-tape notion, we will observe that the two-tape notions and the backtrack notions exhibit amazing and fascinating behaviors as operations on the subsets of Baire space. While the one-tape models are almost statistic, the two-tape models can be understood as learning proof models with *bounded-errors*, and the backtrack tape models can be understood as learning proof models with no predetermined bound for errors. In section 3, we adopt the two-tape notions expect for the classical one-tape disjunction $\cup$, since the two-tape notions (the bounded-errors learning models) is useful to clarify differences among the classes $[\mathfrak{C}_T]^1_1, [\mathfrak{C}_T]^1_{<\omega}, [\mathfrak{C}_T]^1_{\omega|<\omega}, [\mathfrak{C}_T]^{<\omega}_1$ which are defined (as certain classes of bounded-errors functions) later. In section 4, we adopt dynamic generalizations of the backtrack tape models since such models turn out to be a strong tool to establish many theorems.

We now note that the straightforward two-tape (backtrack) notion is hard to use, since, for instance, $[\![P \vee Q]\!]^2_{\mathsf{LCM}[2]}$ may not belong to Π^0_1 even if P and Q are Π^0_1. This observation prompts us to define *consistent* two-tape disjunctions. *The consistency set* $\mathrm{Con}(T_i)_{i\in I}$ *for* $\{T_i\}_{i\in I}$ is defined as follows.

$$\mathrm{Con}(T_i)_{i\in I} = \{f \in (I \times \mathbb{N})^{\mathbb{N}} : (\forall i \in I)(\forall n \in \mathbb{N})\ \mathtt{pr}_i(f \upharpoonright n) \in T_i\}.$$

Then we use the following modified definitions.

Definition 15. Let P_0 and P_1 denote Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$.

$$P_0 \triangledown_\omega P_1 = [\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}} \cap \mathrm{Con}(T_{P_0}, T_{P_1}).$$
$$P_0 \triangledown_n P_1 = [\![P_0 \vee P_1]\!]^2_{\mathsf{LCM}[n]} \cap \mathrm{Con}(T_{P_0}, T_{P_1}).$$
$$P_0 \triangledown_\infty P_1 = [\![P_0 \vee P_1]\!]^2_{\mathsf{CL}} \cap \mathrm{Con}(T_{P_0}, T_{P_1}).$$

Here T_{P_0} and T_{P_1} are corresponding trees for P_0 and P_1, respectively.

Proposition 16. *Let P and Q be Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$.*

1. $P \triangledown_n Q \equiv^1_1 [\![P \vee Q]\!]^2_{\mathsf{LCM}[n]}$ *for each $n \in \mathbb{N}$.*
2. $P \triangledown_\omega Q \equiv^1_1 [\![P \vee Q]\!]^2_{\mathsf{LCM}}$.
3. $P \triangledown_\infty Q \equiv^1_1 [\![P \vee Q]\!]^2_{\mathsf{CL}}$.

Proof. For each item, clearly $P \triangledown_* Q \geq^1_1 [\![P \vee Q]\!]^2_*$. Thus, it suffices to construct a computable functional Φ witnessing $P \triangledown_* Q \leq^1_1 [\![P \vee Q]\!]^2_*$. Let T_0 and T_1 denote the corresponding computable trees for P and Q respectively. Set $\Phi(\langle\rangle) = \langle\rangle$. Fix $\sigma \in (2 \times \mathbb{N})^{<\mathbb{N}}$. Assume that $\Phi(\sigma^-)$ has already been defined, and $\sigma = \sigma^{-\frown}\langle(i, k)\rangle$ for some $i < 2$ and $k \in \mathbb{N}$. If $\mathtt{pr}_i(\sigma) \in T_i$, then we set $\Phi(\sigma) = \Phi(\sigma^-)^\frown\langle(i, k)\rangle$. Otherwise, we set $\Phi(\sigma) = \Phi(\sigma^-)$. Clearly, Φ is a computable function, since T_i is computable for each $i < 2$. For any $g \in (2 \times \mathbb{N})^{\mathbb{N}}$, clearly $\mathtt{mc}(\Phi(g)) \leq \mathtt{mc}(g)$. Fix $g \in [\![P \vee Q]\!]^2_*$, where $* \in \{\mathsf{LCM}, \mathsf{LCM}[n], \mathsf{CL}\}$. Then $\mathtt{pr}_i(g) \in P_i$ for some $i < 2$, where $P_0 = P$ and $P_1 = Q$. Therefore, $\Phi(g)$ is total, and $\mathtt{pr}_i(\Phi(g)) \in P_i$ for such $i < 2$. □

Proposition 17. *Let P and Q be Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$.*

1. $P \triangledown_n Q$ *is Π^0_1, for any $n \in \mathbb{N}$.*

(2) *$P\triangledown_\omega Q$ is Σ^0_2.*
(3) *$P\triangledown_\infty Q$ is Π^0_1.*

Proof. Let T_0 and T_1 denote the corresponding computable trees for P and Q respectively. We consider the following computable tree:

$$T_{P,Q,n} = \{\sigma \in (2\times\mathbb{N})^{<\mathbb{N}} : (\forall i<2)\ \mathtt{pr}_i(\sigma)\in T_i \ \&\ \mathtt{mc}(\sigma)<n\}.$$

Note that $T_{P,Q,n}$ is uniformly computable in n, since $\mathtt{pr}_i(\sigma)$ and $\mathtt{mc}(\sigma)$ are computable uniformly in $\sigma\in\mathbb{N}^{<\mathbb{N}}$. Clearly, $P\triangledown_n Q\subseteq[T_{P,Q,n}]$. Moreover, for any $g\in[T_{P,Q,n}]$, $\mathtt{pr}_i(g)$ is total for some $i<2$. Then, $\mathtt{pr}_i(g)\in[T_i]$ for such i, and $\mathtt{mc}(g)\le n$, since the relation $\mathtt{mc}(f)\le n$ is equivalent to $(\forall k)\ \mathtt{mc}(f\upharpoonright k)\le n$. Thus, $g\in P\triangledown_n Q$. Consequently, $P\triangledown_n Q=[T_{P,Q,n}]$ is Π^0_1. Hence, $P\triangledown_\omega Q=\bigcup_n[T_{P,Q,n}]$ is Σ^0_2. The items (3) also follows from the similar argument. ◻

Definition 18. Pick any $*\in\mathbb{N}\cup\{\omega\}\cup\{\infty\}$. For each disjunctive notions $\triangledown_*$ and sets $P_0,P_1\subseteq\mathbb{N}^\mathbb{N}$, fix any trees $T_{P_i}\subseteq\mathbb{N}^{<\mathbb{N}}$ for $i<2$, and $T_*\subseteq\mathbb{N}^{<\mathbb{N}}$ such that $[T_{P_i}]$ is the closure of P_i for each $i<2$, and $[T_*]$ is also the closure of $P_0\triangledown_* P_1$, under the Baire topology. Then *the heart of $P_0\triangledown_* P_1$* is defined by $T^\heartsuit_*=\{\sigma\in T_*:(\forall i<2)\ \mathtt{pr}_i(\sigma)\in T^{ext}_{P_i}\}$.

Proposition 19. *$T^\heartsuit_*\subseteq T^{ext}_*$.*

Proof. $T^\heartsuit_*\subseteq\{\sigma\in T_*:(\exists i<2)\ \mathtt{pr}_i(\sigma)\in T^{ext}_{P_i}\}=T^{ext}_*$. ◻

This notion is not quite useful when either P_0 or P_1 is not closed. The heart plays an important role only when P_0 and P_1 are Π^0_1 subsets of $2^\mathbb{N}$, with the corresponding computable tree $T_*\subseteq 4^{<\mathbb{N}}$.

2.3. Concatenations versus Disjunction Operations.

Definition 20. Let L_P denote the set of all leaves of the corresponding tree for a closed set P. Then *the (non-commutative) concatenation of P and Q* is defined as follows.

$$P^\frown Q=P\cup\bigcup_{\rho\in L_P}\rho^\frown Q.$$

The commutative concatenation of P and Q is defined by $P\triangledown Q=(P^\frown Q)\oplus(Q^\frown P)$.

Remark. On the study of Wadge degrees of finite level of Borel hierarchy, Duparc [30] introduced various operators such as $P^\longrightarrow Q=P\cup\bigcup_{\rho\in\mathbb{N}^{<\mathbb{N}}}\rho^\frown\langle\sharp\rangle^\frown Q$. The following proposition indicates that our non-commutative concatenation is essentially same as Duparc's operation $P^\longrightarrow Q$.

Proposition 21. *Let P,Q be Π^0_1 subsets of Baire space $\mathbb{N}^\mathbb{N}$. Then, the concatenation $P^\frown Q$ is $(1,1)$-equivalent to the set $P^\to Q$ of all infinite paths of the tree $\{\sigma^\frown\langle\sharp\rangle^\frown\tau:\sigma\in T_P\ \&\ \tau\in T_Q\}$.*

Proof. To see $P^\to Q\le^1_1 P^\frown Q$, we inductively define a total computable function $\mathtt{cut}:\mathbb{N}^\mathbb{N}\to\mathbb{N}^\mathbb{N}$. First set $\mathtt{cut}(\langle\rangle)=\langle\rangle$, and fix $\sigma=\sigma^{-\frown}\langle n\rangle\in\mathbb{N}^{<\mathbb{N}}$. We assume that $\mathtt{cut}(\sigma^-)$ has been already defined. If $\sigma=\sigma^{-\frown}\langle n\rangle\in L_P$, then we set $\mathtt{cut}(\sigma)=\mathtt{cut}(\sigma^-)^\frown\langle n,\sharp\rangle$. Otherwise, we set $\mathtt{cut}(\sigma)=\mathtt{cut}(\sigma^-)^\frown\langle n\rangle$. Then, $\mathtt{cut}$ is computable, since P is Π^0_1 and then T_P is computable. Moreover, we can see the following.

$$\mathtt{cut}(f)=\begin{cases} f & \text{if } f\in P,\\ (f\upharpoonright k)^\frown\langle\sharp\rangle^\frown f^{\leftarrow k} & \text{if } (\exists k\in\mathbb{N})\ f\upharpoonright k\in L_P.\end{cases}$$

Clearly, $P^\to Q\le^1_1 P^\frown Q$ via the computable function $\mathtt{cut}$.

Conversely, we consider the computable function $\mathtt{leaf}:\mathbb{N}^{<\mathbb{N}}\to\mathbb{N}^{<\mathbb{N}}$ which maps σ to the least leaf of L_P extending σ. Then, we inductively define a computable function Γ witnessing $P^\frown Q\le^1_1 P^\to Q$ as follows. First set $\Gamma(\langle\rangle)=\langle\rangle$, and fix $\sigma=\sigma^{-\frown}\langle n\rangle\in(\mathbb{N}\cup\{\sharp\})^{<\mathbb{N}}$. We assume that $\Gamma(\sigma^-)$ has been already defined. If $n\neq\sharp$, then we set $\Gamma(\sigma)=\Gamma(\sigma^-)^\frown\langle n\rangle$. If $n=\sharp$, then we set $\Gamma(\sigma)=\mathtt{leaf}(\Gamma(\sigma^-))$. It is easy to see that $P^\frown Q\le^1_1 P^\to Q$ via Γ. ◻

Remark.

1. A variation of the non-commutative concatenation was used by Binns-Simpson [8] to embed the free Boolean algebra into the Muchnik lattice of Π^0_1 subsets of Cantor space, and such a variation was called *a recursive meet*. The concatenation can be extended to the following infinitary concatenation: $P^\frown\{Q_i\}_{i\in\mathbb{N}} = P\cup\bigcup_n \rho_n{}^\frown Q_n$, where ρ_n is the n-th leaf of the corresponding computable tree of P. Then the recursive meet of $\{Q_i\}_{i\in\mathbb{N}}$ is defined by $\overrightarrow{\bigoplus}_{i\in\mathbb{N}} Q_i = \mathsf{CPA}^\frown\{Q_i\}_{i\in\mathbb{N}}$, where CPA is the set of all *complete consistent extensions of Peano Arithmetic*. Note that CPA is a *Medvedev complete* Π^0_1 subset of $2^{\mathbb{N}}$. That is to say, for any nonempty Π^0_1 subset $P\subseteq 2^{\mathbb{N}}$, a computable function $\Phi:\mathsf{CPA}\to P$ exists. The significance of the recursive meet $\overrightarrow{\bigoplus}_i Q_i$ is that it is a Π^0_1 subset of the compact space $2^{\mathbb{N}}$, while $\bigoplus_i Q_i$ is a subset of $\mathbb{N}^{\mathbb{N}}$, and, indeed, is not compact. Actually, Simpson [61] used the recursive meet to construct an embedding of the Muchnik degrees of Σ^0_2 subsets of $2^{\mathbb{N}}$ into the Muchnik degrees of Π^0_1 subsets of $2^{\mathbb{N}}$.
2. More explicitly, the non-commutative concatenation has been used by Cenzer-Kihara-Weber-Wu [19]. A Π^0_1 set $P\subseteq 2^{\mathbb{N}}$ is *tree-immune* if the Π^0_1 tree $\{\sigma\in 2^{<\mathbb{N}} : P\cap[\sigma]\neq\emptyset\}$ includes no infinite computable subtree. Cenzer-Kihara-Weber-Wu [19] showed that $\mathsf{CPA}^\frown\mathsf{CPA}$ has a greatest Medvedev degree of Π^0_1 subsets of $2^{\mathbb{N}}$ with no tree-immune.

Proposition 22. *Let P, Q be Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$.*

1. $P\triangledown P\equiv^1_1 P^\frown P$.
2. $P\triangledown Q\equiv^1_1 \llbracket P\vee Q\rrbracket^2_{\mathsf{LCM}[2]}$.

Proof. (1) $P\triangledown P = (P^\frown P)\oplus(P^\frown P)\equiv^1_1 P^\frown P$. (2) By Proposition 16 (1), we have $P\triangledown_2 Q\equiv^1_1 \llbracket P\vee Q\rrbracket^2_{\mathsf{LCM}[2]}$. Then, $P\triangledown_2 Q\leq^1_1 P\triangledown Q$ is witnessed by the following reduction Δ.

$$\Delta(f)=\begin{cases}\mathtt{write}(f(0),f^{\leftarrow 1}), & \text{if } f^{\leftarrow 1}\in[T_{\sigma(0)}],\\ \mathtt{write}(f(0),f^{\leftarrow 1}\upharpoonright k)^\frown\mathtt{write}(1-f(0),f^{\leftarrow k+1}), & \text{if } (\exists k\in\mathbb{N})\ f^{\leftarrow 1}\in L_{\sigma(0)}.\end{cases}$$

Here, T_0 and T_1 are the corresponding computable trees for P and Q respectively, and L_i is the set of all leaves of T_i for each $i<2$. Clearly, Δ is computable. Fix $\langle i\rangle^\frown g\in P\triangledown Q$. Obviously, $\mathtt{mc}(\langle i\rangle^\frown g)<2$. If $g\in[T_i]$ then $\mathtt{pr}_i(\Delta(\langle i\rangle^\frown g)) = g\in[T_i]$, and if $g=\sigma^\frown h$ for some $\sigma\in L_i$ and $h\in[T_{1-i}]$ then $\mathtt{pr}_i(\Delta(\langle i\rangle^\frown\sigma^\frown h)) = h\in[T_{1-i}]$. Hence, $\Delta(\langle i\rangle^\frown g)\in P\triangledown_2 Q$.

To see $P\triangledown Q\leq^1_1 P\triangledown_2 Q$, it suffices to construct a computable functional Γ witnessing $(P^\rightarrow Q)\oplus(Q^\rightarrow P)\leq^1_1 P\triangledown_2 Q$ by Proposition 21. Set $\Gamma(\langle\rangle)=\langle\rangle$, and $\Gamma(\langle(i,n)\rangle)=\langle i,n\rangle$ for any $i<2$ and $n\in\mathbb{N}$. Fix $\sigma=\sigma^{---\frown}\langle(i,m),(j,n)\rangle\in(2\times\mathbb{N})^{<\mathbb{N}}$, and assume that $\Gamma(\sigma^-)$ is already defined. If $i\neq j$, then set $\Gamma(\sigma)=\rho^\frown\langle\sharp,n\rangle$. Otherwise, set $\Gamma(\sigma)=\Gamma(\sigma^-)^\frown\langle n\rangle$. Clearly Γ is computable. Fix $g\in P\triangledown_2 Q$. If $\mathtt{mc}(g)=0$, then $\Gamma(g)=\langle i\rangle^\frown\mathtt{pr}_i(g)\in P\oplus Q\subseteq(P^\rightarrow Q)\oplus(Q^\rightarrow P)$, where $i=(g(0))_0$. If $\mathtt{mc}(g)=1$, then $\mathtt{pr}_i(g)$ is a finite string, where $i=(g(0))_0$. In this case, we can easily see $\Gamma(g)=\langle i\rangle^\frown\mathtt{pr}_i(g)^\frown\langle\sharp\rangle^\frown\mathtt{pr}_{1-i}(g)\in(P^\rightarrow Q)\oplus(Q^\rightarrow P)$. □

In the case of $P\triangledown P$, we use the non-commutative concatenation $P^\frown P$ to simplify our proof without mentioning.

Remark. The commutative concatenation can be viewed as a kind of *consistent conservative Popperian learning* (see [38]).

- The term “*consistent*” means: the scientist should modify his hypothesis whenever it was found to be refuted.
- The term “*conservative*” means: the scientist changes his hypothesis only when it was found to be refuted.
- The term “*Popperian*” means: the scientist can test whether his hypothesis is currently consistent or refuted.

The notion of *Popperian learning* is introduced by Case and Ngo-Manguelle [17] based on Gold’s theory of “identification in the limit” [33]. A learner (a scientist) is a computable function $\Psi:\mathbb{N}^{<\mathbb{N}}\to\mathbb{N}$, and a natural phenomenon is a computable function $f:\mathbb{N}\to\mathbb{N}$. Then the formula $\Psi(f\upharpoonright n)=e$ means the

Table 1. Hierarchy of Disjunctions

$P \oplus Q$	Intuitionistic disujunction ($= P \triangledown_1 Q$)
$P \cup Q$	Classical one-tape disjunction
$P \triangledown Q$	Commutative concatenation ($= P \triangledown_2 Q$)
$P \triangledown_n Q$	LCM disjunction with mind-changes-bound n
$P \triangledown_\omega Q$	LCM disjunction
$P \triangledown_\infty Q$	Classical disjunction

following situation: the scientist Ψ predicts that a rule generating the phenomenon f can be explained by a word (a formula, or an algorithm) e (i.e., $f = \Phi_e$) when he observes $f(0), \dots f(n-1)$. We say that Ψ *learns* f if $\Phi_{\lim_n \Psi(f \upharpoonright n)} = f$. The learner Ψ is *Popperian* if $\Phi_{\Psi(\sigma)}$ is total for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$. The learner Ψ is *consistent* at $\sigma \in \mathbb{N}^{<\mathbb{N}}$ if $\Phi_{\Psi(\sigma)} \upharpoonright |\sigma| = \sigma$. The learner Ψ is *conservative* if, for any $\sigma \in \mathbb{N}^{<\mathbb{N}}$, $\Psi(\sigma) = \Psi(\sigma^-)$ whenever $\Phi_{\Psi(\sigma^-)} \upharpoonright |\sigma| = \sigma$. Note that, for every Popperian learner Ψ, he can algorithmically determine whether Ψ is consistent at σ or not, for a given $\sigma \in \mathbb{N}^{<\mathbb{N}}$. The terminology "*Popperian*" derives from Popper's refutability principle in philosophy of science. In this context, the non-commutative concatenation $P^\frown Q$ of P and Q represents the following situation: if, and only if, a choice on the first hypothesis P is refuted, then the scientist proposes the second (refutable) hypothesis Q and start verifying it. The reason why $P^\frown Q$ is Popperian will be discussed later.

Proposition 23. *For* Π^0_1 *sets* $P, Q \subseteq \mathbb{N}^\mathbb{N}$ *and* $n \in \mathbb{N}$,

$$\llbracket P \vee Q \rrbracket^2_{\mathsf{LCM}} \leq^1_1 \llbracket P \vee Q \rrbracket^2_{\mathsf{LCM}[n+2]} \leq^1_1 \llbracket P \vee Q \rrbracket^1_{\mathsf{CL}} \leq^1_1 \llbracket P \vee Q \rrbracket^1_{\mathsf{Int}}.$$

Proof. It suffices to show $P \triangledown Q \leq^1_1 P \cup Q$, since $\llbracket P \vee Q \rrbracket^1_{\mathsf{CL}} \equiv^1_1 P \cup Q$ by Proposition 11 (5) and $\llbracket P \vee Q \rrbracket^2_{\mathsf{LCM}[2]} \equiv^1_1 P \triangledown Q$ by Proposition 22 (2). Indeed, we can show that $(P^\frown Q) \otimes (Q^\frown P) \leq^1_1 P \cup Q$. We construct a computable functional Φ witnessing $P^\frown Q \leq^1_1 P \cup Q$. If $\sigma \in T_P$, then set $\Phi(\sigma) = \sigma$. If $\sigma \notin T_P$, then pick a unique $\rho \subseteq \sigma$ such that $\rho \in L_P$, and set $\Phi(\sigma) = \rho^\frown \sigma$ for such ρ, where L_P is the set of all leaves of T_P. Clearly Φ is computable, and note that $\Phi(\sigma) \subseteq \Phi(\tau)$ whenever $\sigma \subseteq \tau$. If $g \in P$, then $\Phi(g) = g \in P$. If $g \in Q \setminus P$, then there is a unique $\rho \subset g$ such that $\rho \in L_P$, and $\Phi(g) = \rho^\frown g \in P^\frown Q$. Thus, $P^\frown Q \leq^1_1 P \cup Q$ via Φ. ◻

Remark. These LCM-disjunction operations $\triangledown_2$ (the concatenation operation) and $\triangledown_\omega$ are related to the sequential disjunction [40] and the toggling disjunction [41] in *Computability Logic* [39], respectively.

Remark. In Theorem 161 (1), we will see that the non-commutative concatenation $P^\frown Q$ can be viewed as *the* Σ^0_1*-law-of-excluded-middle operation*. Here the arithmetical hierarchy for the law of excluded middle is studied by Akama-Berardi-Hayashi-Kohlenbach [1].

2.4. Nonuniformly Computable Discontinuous Functions.

Proposition 24. *For every partial computable function* $\Psi :\subseteq \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$, *we can effectively find a total computable function* $\widehat{\Psi} : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$ *such that, if* $\lim_s \Psi(f \upharpoonright s)$ *converges, then* $\lim_s \widehat{\Psi}(f \upharpoonright s)$ *converges to* $\lim_s \Psi(f \upharpoonright s)$.

Proof. Without loss of generality, we can assume $\Psi(\langle\rangle) = 0$. For given $\sigma \in \mathbb{N}^{<\mathbb{N}}$, put $\sigma^* = \max\{\tau \subseteq \sigma : \Psi(\tau)[|\sigma|] \downarrow\}$. Then define $\widehat{\Psi}(\sigma) = \Psi(\sigma^*)$ for every $\sigma \in \mathbb{N}^{<\mathbb{N}}$. If $\lim_s \Psi(f \upharpoonright s)$ converges then clearly $\lim_s \widehat{\Psi}(f \upharpoonright s)$ also converges. ◻

For the e-th partial computable function $\Phi_e :\subseteq \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$, $\widehat{\Phi_e}$ is denoted by Ψ_e. The set $\{\Psi_e\}_{e \in \mathbb{N}}$ is referred as *the effective enumeration of all learners*, and Ψ_e is called *the* e*-th learner*.

Table 2. Seven Classes of Nonuniformly Computable Functions

$[\mathfrak{C}_T]^1_1$	(Uniformly) computable
$[\mathfrak{C}_T]^1_{<\omega}$	Learnable with bounded mind changes
$[\mathfrak{C}_T]^1_{\omega\|<\omega}$	Learnable with bounded errors
$[\mathfrak{C}_T]^1_\omega$	Learnable
$[\mathfrak{C}_T]^{<\omega}_1$	k-wise computable
$[\mathfrak{C}_T]^{<\omega}_\omega$	Learnable by a team
$[\mathfrak{C}_T]^\omega_1$	Nonuniformly computable

Remark. We urge the reader not to confuse the notions $\Psi(\sigma)$ and $\Phi(\sigma)$ for a leaner Ψ and a computable function Φ (on $\mathbb{N}^\mathbb{N}$). In the former case, $\Psi(\sigma)$ simply denotes the output (the inference) of the learner Ψ based on the current input σ. In the latter case, however, we use σ as an initial segment of some oracle information, and so really $\Phi(\sigma)$ denotes a string $\langle\Phi(\sigma;0),\Phi(\sigma;1),\Phi(\sigma;2),\dots\rangle$.

We now introduce various subclasses of nonuniformly computable functions on $\mathbb{N}^\mathbb{N}$ based on Learning Theory. For an ordinal α, if α is a successor ordinal $\beta+1$, then α^- denotes β, and if α is a limit ordinal, then α^- denotes the symbol $<\alpha$.

Definition 25. Let D be a subset of Baire space $\mathbb{N}^\mathbb{N}$, and $\alpha,\beta,\gamma \leq \omega+1$ be ordinals. A function $\Gamma : D \to \mathbb{N}^\mathbb{N}$ is $(\alpha^-,\beta^-|\gamma^-)$-*computable* if there are $a<\alpha$, $b<\beta$, $c<\gamma$ and $I \subseteq \mathbb{N}$ of cardinality a such that, for any $g \in D$, there is $e \in I$ satisfying the following three conditions.

(1) (Learnability) $\lim_n \Psi_e(g \upharpoonright n)$ converges, and $\Phi_{\lim_n \Psi_e(g\upharpoonright n)}(g) = \Gamma(g)$.
(2) (Mind-Change Condition) $\#\{n \in \mathbb{N} : \Psi_e(g \upharpoonright n+1) \neq \Psi_e(g \upharpoonright n)\} < b$.
(3) (Error Condition) $\#\{\Psi_e(g \upharpoonright n) : n \in \mathbb{N}\} \leq c$.

If $\gamma = \omega$, then we simply say that Γ is (α,β)-*computable* for $(\alpha,\beta|\gamma)$-computable function Γ. Let $[\mathfrak{C}_T]^\alpha_\beta$ (resp. $[\mathfrak{C}_T]^\alpha_{\beta|\gamma}$) denote the set of all (α,β)-computable (resp. $(\alpha,\beta|\gamma)$-computable) functions.

Remark. Some of $(\alpha,\beta|\gamma)$-computability notions are related to learnability notions: Every $(1,<\omega)$-computable function is *learnable with bounded mind-changes*; every $(1,\omega|<\omega)$-computable function is *learnable with bounded errors*; every $(1,\omega)$-computable function is *learnable*; every $(<\omega,1)$-computable function is *k-wise computable*; and every $(<\omega,\omega)$-computable function is *team-learnable*. The concept of learnability in the context of real number computation has been studied by some researchers including Chadzelek-Hotz [23], Ziegler [79, 78], and de Brecht-Yamamoto [26, 27]. The notion of k-wise computability has been also studied by, for example, Ziegler [81].

Proposition 26. *Let Γ be a partial function on Baire space $\mathbb{N}^\mathbb{N}$.*

(1) *If Γ is $(\alpha_0^-,\beta_0^-|\gamma_0^-)$-computable, $\alpha_0 \leq \alpha_1, \beta_0 \leq \beta_1$, and $\gamma_0 \leq \gamma_1$, then Γ is $(\alpha_1^-,\beta_1^-|\gamma_1^-)$-computable.*
(2) *Γ is $(\alpha,1)$-computable if and only if Γ is $(\alpha,\beta|1)$-computable.*
(3) *Γ is (α,β)-computable if and only if Γ is $(\alpha,\beta|\beta)$-computable.*
(4) *Γ is $(1,1)$-computable if and only if Γ is computable.*
(5) *Γ is $(\omega,1)$-computable if and only if Γ is (ω,ω)-computable if and only if Γ is nonuniformly computable, i.e., $\Gamma(g) \leq_T g$ for any $g \in \mathrm{dom}(\Gamma)$, where recall that $\leq_T$ denotes the Turing reducibility.*

Proof. The items (1) and (2) easily follow from definition. The item (3) follows from $\#\{\Psi_e(g \upharpoonright n) : n \in \mathbb{N}\} - 1 \leq \#\{n : \Psi_e(g \upharpoonright n+1) \neq \Psi_e(g \upharpoonright n)\}$.

(4) If Γ is computable via Φ_e, then Γ is $(1,1)$-computable via the singleton $\{i(e)\}$, where $\Gamma_{i(e)}(\sigma) = e$ for any $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Assume that Γ is $(1,1)$-computable via a singleton $\{e\}$. Then $\Psi_e(\sigma) = \Psi_e(\langle\rangle)$ for any σ extendible to an element of $\mathrm{dom}(\Gamma)$, since $\#\{n \in \mathbb{N} : \Psi_e(g \upharpoonright n+1) \neq \Psi_e(g \upharpoonright n)\} = 0$ for any $g \in \mathrm{dom}(\Gamma)$. Therefore, Γ is computable via $\Phi_{\Psi_e(\langle\rangle)}$.

(5) If Γ is nonuniformly computable, then Γ is $(\omega, 1)$-computable via $\{i(e)\}_{e\in\mathbb{N}}$, where $\Gamma_{i(e)}(\sigma) = e$ for any $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Assume that Γ is (ω, ω)-computable via I. For any $g \in \mathrm{dom}(\Gamma)$, there is $e \in I$ such that $\lim_n \Psi_e(g \upharpoonright n)$ converges to some value $p \in \mathbb{N}$, and $\Phi_p(g) = \Gamma(g)$. Thus, $\Gamma(g) \leq_T g$ via Φ_p. □

Proposition 27. *For each $m, n \in \mathbb{N}$, every $(m, \omega|n)$-computable function is $(m \cdot n, 1)$-computable.*

Proof. Assume that $\Gamma : D \to \mathbb{N}^{\mathbb{N}}$ is $(m, \omega|n)$-computable with m-learners $\{\Psi^e\}_{e<m}$ with n-errors. Now, we define an algorithm Φ^e_k for any $e < m$ and $k < n$, and we ensure the following property:

$$(\forall g \in D)(\exists e < m)(k < n)\ \Phi^e_k(g) = \Gamma(g).$$

The algorithm Φ^e_k proceeds as follows for g. Let $B^e_s(g)$ be the set of all indices occurring in hypothesis of the learner Ψ^e by stage s, i.e., $B^e_n(g) = \{e \in \mathbb{N} : (\exists m \leq s)\ e = \Psi(g \upharpoonright m)\}$. Put $p^e_k(g) = \mu s[|B^e_s(g)| = k]$, and let $d^e_k(g)$ be the unique element of $B^e_{p^e_k(g)}(g) \setminus B^e_{p^e_k(g)-1}(g)$, where $B^e_{-1}(g) = \emptyset$. Clearly, $d^e_k :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}$ is partially computable. Then, we set $\Phi^e_k(g) = \Phi_{d^e_k(g)}(g)$. For any $g \in D$, there is $e < m$ such that $\lim_s \Psi^e(g \upharpoonright s)$ converges to some correct computation d of $\Gamma(g)$, i.e., $\Phi_d(g) = \Gamma(g)$. Since $\#\{\Psi^e(g \upharpoonright s) : s \in \mathbb{N}\} < n$, we have $d^e_k(g) = d$ for some $k < n$. Thus, for any $g \in D$, there are $e < m$ and $k < n$ such that $\Phi^e_k(g) = \Gamma(g)$. Therefore, if i^e_k is an index of Φ^e_k for each $e < m$ and $k < n$, then Γ is $(m \cdot n, 1)$-computable via an upper bound $\max |\{i^e_k : e < m \ \&\ k < n\}|$. □

Corollary 28. *Γ is $(< \omega, \omega| < \omega)$-computable if and only if Γ is $(< \omega, 1)$-computable.*

Proof. Every $(< \omega, \omega| < \omega)$-computable function Γ is $(m, \omega|n)$-computable for some $m, n < \omega$. Therefore, by Proposition 27, Γ is $(m \cdot n, 1)$-computable. In particular, Γ is $(< \omega, 1)$-computable, since $m \cdot n < \omega$. □

Proposition 29. *For each $i < 2$, let Γ_i be a partial $(\alpha_i, \beta_i|\gamma_i)$-computable function on Baire space $\mathbb{N}^{\mathbb{N}}$, where $\alpha_i, \beta_i, \gamma_i \leq \omega$ are ordinals. Then $\Gamma_1 \circ \Gamma_0$ is $(\alpha_0 * \alpha_1, \beta_0 * \beta_1|\gamma_0 * \gamma_1)$-computable, where $*$ is the multiplication as the cardinals, or equivalently, $\kappa * \lambda = \min\{\kappa \cdot \lambda, \omega\}$ for ordinals $\kappa, \lambda \leq \omega$.*

Proof. For each $i < 2$, since Γ_i is $(\alpha_i, \beta_i|\gamma_i)$-computable, there is a collection of learners, $\{\Psi^i_j\}_{j<\alpha_i}$ and a finite cover $\{U^i_j\}_{j<\alpha_i}$ of $\mathrm{dom}(\Gamma_i)$ such that $\Gamma_i(f) = \Phi_{\lim_n \Psi^i_j(f\upharpoonright n)}(f \upharpoonright n)$ and $\#\{n \in \mathbb{N} : \Psi^i_j(f \upharpoonright n + 1) \neq \Psi^i_j(f \upharpoonright n)\} < \beta_i$ and $\#\{\Psi^i_j(f \upharpoonright n) : n \in \mathbb{N}\} < \gamma_i$, for any $j < \alpha_1$ and $f \in U^i_j$. Fix $j < \alpha_0$ and $k < \alpha_1$. Then $\Psi^*_{j,k}(\sigma)$ is defined as follows. Let $E(\sigma)$ be the set of all strings $\rho \subseteq \sigma$ satisfying $\Psi^0_j(\rho) = \Psi^0_j(\sigma)$. Let $\rho(\sigma)$ be the shortest string of $E(\sigma)$, and set $E^*(\sigma) \setminus \{\rho(\sigma)\}$. If $\#\{\tau \in E^*(\sigma) : \Psi^1_k(\tau) \neq \Psi^1_k(\tau^-)\} < \beta_1$ and $\#\{\Psi^1_k(\tau) : \tau \in E(\sigma)\} < \gamma_1$, then put $\Psi^*_{j,k}(\sigma) = \Psi^1_k(\Phi_{\Psi^0_j(\sigma)}(\sigma))$. Otherwise, put $\Psi^*_{j,k}(\sigma) = \Psi^*_{j,k}(\sigma^-)$. For given σ, we compute an index $\Psi_{j,k}(\sigma)$, where $\Phi_{\Psi_{j,k}(\sigma)}(f) = \Phi_{\Psi^*_{j,k}(\sigma))}(\Phi_{\Psi^0_j(\sigma)}(f))$ for any f.

Note that $f \in \mathrm{dom}(\Gamma_1 \circ \Gamma_0)$ if and only if $f \in \mathrm{dom}(\Gamma_0)$ and $\Gamma_0(f) \in \mathrm{dom}(\Gamma_1)$. Therefore, for such f, there are $j < \alpha_0$ and $k < \alpha_1$ such that $f \in U^0_j$ and $\Gamma_0(f) \in U^1_k$. Assume that $f \in \mathrm{dom}(\Gamma_1 \circ \Gamma_0) \cap U^0_j$ and $\Gamma_0(f) \in U^1_k$. It is easy to see that $\Psi^*_{j,k}$ is computable, $\#\{n \in \mathbb{N} : \Psi^*_{j,k}(f \upharpoonright n+1) \neq \Psi^*_{j,k}(f \upharpoonright n)\} < \beta_0 * \beta_1$ and $\#\{\Psi^*_{j,k}(f \upharpoonright n) : n \in \mathbb{N}\} < \gamma_0 * \gamma_1$. Moreover, there exist s and e_0 such that $\Psi^0_j(f \upharpoonright t) = \Psi^0_j(f \upharpoonright s) = e_0$ for any $t \geq s$. Fix such s. Since $\Phi_{e_0}(f) = \Gamma_0(f) \in U^1_k$, for any $t \geq s$, $\#\{\tau \in E^*(f \upharpoonright t) : \Psi^1_k(\tau) \neq \Psi^1_k(\tau^-)\} < \beta_1$ and $\#\{\Psi^1_k(\tau) : \tau \in E(f \upharpoonright t)\} < \gamma_1$, since $E(f \upharpoonright t) = E(f \upharpoonright s)$ and by our choice of Ψ^1_k. Therefore, $\lim_n \Psi^*_{j,k}(f \upharpoonright n)$ converges to $\lim_n \Psi^1_k(\Gamma_0(f \upharpoonright n))$. However, there exist $u \geq s$ and e_1 such that $\Psi^1_k(\Gamma_0(f \upharpoonright v)) = \Psi^1_k(\Gamma_0(f \upharpoonright u)) = e_1$ for any $v \geq u$, since $\{\Gamma_0(f \upharpoonright u)\}_{u\geq s}$ is an increasing sequence of strings and $\Gamma_0(f) \in \mathrm{dom}(\Gamma_1)$. Here $\Phi_{e_1}(\Gamma_0(f)) = \Gamma_1(\Gamma_0(f))$. Thus,

$$\Phi_{\lim_n \Psi_{j,k}(f\upharpoonright n)}(f) = \Phi_{\lim_n \Psi^*_{j,k}(f\upharpoonright n)}(\Phi_{\lim_n \Psi^0_j(f\upharpoonright n)}(f)) = \Phi_{\lim_n \Psi^1_k(\Gamma_0(f))}(\Gamma_0(f)) = \Gamma_1(\Gamma_0(f)).$$

Consequently, $\Gamma_1 \circ \Gamma_0$ is $(\alpha_0 * \alpha_1, \beta_0 * \beta_1|\gamma_0 * \gamma_1)$-computable, via $\{\Psi_{j,k}\}_{j<\alpha_0,k<\alpha_1}$. □

Corollary 30. *$[\mathfrak{C}_T]^{\alpha}_{\beta|\gamma}$ forms a monoid under composition, for any $\alpha, \beta, \gamma \in \{1, < \omega, \omega\}$.*

Proof. Straightforward from Proposition 29. □

Proposition 31. *$[\mathfrak{C}_T]^1_{<\omega}$ is the smallest monoid including $[\mathfrak{C}_T]^1_2$; $[\mathfrak{C}_T]^1_{\omega|<\omega}$ is the smallest monoid including $[\mathfrak{C}_T]^1_{\omega|2}$. $[\mathfrak{C}_T]^{<\omega}_1$ is the smallest monoid including $[\mathfrak{C}_T]^2_1$; $[\mathfrak{C}_T]^{<\omega}_\omega$ is the smallest monoid including $[\mathfrak{C}_T]^2_\omega$.*

Proof. We first show that every $(1, n+1)$-computable function Γ can be represented as $\Gamma = \Gamma_1 \circ \Gamma_0$ for some $(1,n)$-computable function Γ_0 and $(1,2)$-computable function Γ_1. Let Ψ be a learner for Γ. We define a learner Ψ_0 for Γ_0 and a learner Ψ_1 for Γ_1. For a given string $\sigma \in \mathbb{N}^{<\mathbb{N}}$, let $\sigma^* \subseteq \sigma$ be the longest initial segment of σ satisfying $\#\{\tau \subseteq \sigma^* : \Psi(\tau) \neq \Psi(\tau^-)\} < n$. Then, on σ, the learner Ψ_0 guesses an index of the partial computable function $g \mapsto g \oplus \Phi_{\Psi(\sigma^*)}(g)$, i.e., $\Gamma_0(g) = \Phi_{\Psi_0(\sigma)}(g) = g \oplus \Phi_{\Psi(\sigma^*)}(g)$ for any $g \in \mathbb{N}^\mathbb{N}$. Note that $\{n \in \mathbb{N} : \Psi_0(g \upharpoonright n+1) \neq \Psi(g \upharpoonright n)\} < n$ for any $g \in \mathbb{N}^\mathbb{N}$. Therefore, Γ_0 is $(1,n)$-computable. For $\sigma \oplus \tau \in \mathbb{N}^\mathbb{N}$, if $\sigma^* = \sigma$ then the learner Ψ_1 guesses an index of the partial computable function $g \oplus h \mapsto h$. If $\sigma^* \neq \sigma$, then Ψ_1 guesses an index of the partial computable function $g \oplus h \mapsto \Phi_{\Psi(\sigma)}(g)$, i.e., $\Phi_{\Psi_1(\sigma\oplus\tau)}(g \oplus h) = \Phi_{\Psi(\sigma)}(g)$. Since Γ is $(1, n+1)$-computable, and by the definition of σ^*, it is easy to see that Γ_1 is $(1,2)$-computable. For $g \in \mathbb{N}^\mathbb{N}$, if $\#\{n \in \mathbb{N} : \Psi(g \upharpoonright n+1) \neq \Psi(g \upharpoonright n)\} < n$, then

$$\Gamma_1(\Gamma_0(g)) = \Gamma_1(g \oplus \Gamma(g)) = \Gamma(g).$$

If $\#\{n \in \mathbb{N} : \Psi(g \upharpoonright n+1) \neq \Psi(g \upharpoonright n)\} = n$, then

$$\Gamma_1(\Gamma_0(g)) = \Gamma_1(g \oplus \Phi_{\Psi(g^*)}(g)) = \Gamma(g).$$

Consequently, $\Gamma = \Gamma_1 \circ \Gamma_0$ as desired.

We next show that every $(1, \omega|n+1)$-computable function Γ can be represented as $\Gamma = \Gamma_1 \circ \Gamma_0$ for some $(1, \omega|n)$-computable function Γ_0 and $(1, \omega|2)$-computable function Γ_1. Assume that Ψ is a learner for Γ, and we enumerate $\{\Psi(\sigma \upharpoonright n) : n \leq |\sigma|\}$ as $\{i^\sigma_m\}_{m \leq |\sigma|}$. Here, if $m < n$ then Ψ guesses i^σ_m before Ψ guesses i^σ_n on some initial segment of σ. Note that, if $\sigma \subseteq \tau$ and i^σ_m is defined, then $i^\sigma_m = i^\tau_m$. On $\sigma \in \mathbb{N}^{<\mathbb{N}}$, if $\Psi(\sigma) \neq i^\sigma_n$, then Ψ_0 guesses an index of the partial computable function $g \mapsto g \oplus \Phi_{\Psi(\sigma)}(g)$. Otherwise, Ψ_0 guesses an index of the partial computable function $g \mapsto g \oplus \Phi_{i^\sigma_0}(g)$. Then, the partial function Γ_0 identified by the learner Ψ_0 is $(1, \omega|n)$-computable. On $\sigma \oplus \tau \in \mathbb{N}^{<\mathbb{N}}$ if $\Psi(\sigma) \neq i^\sigma_n$, then Ψ_1 guesses an index of the partial computable function $g \oplus h \mapsto h$. Otherwise, Ψ_1 guesses an index of partial computable function $g \oplus h \mapsto \Phi_{\Psi(\sigma)}(g)$.

We show that every $(n+1, 1)$-computable function Γ can be represented as $\Gamma = \Gamma_1 \circ \Gamma_0$ for some $(n, 1)$-computable function Γ_0 and $(2,1)$-computable function Γ_1. Assume that Γ is $(n+1,1)$-computable via a collection $\{\Delta_i\}_{i\leq n}$ of partial computable functions. For $g \in \mathbb{N}^\mathbb{N}$, if $\Gamma(g) = \Delta_i(g)$ for some $i < n$, then $\Gamma_0(g) = g \oplus \Delta_i(g)$. Otherwise, we set $\Gamma_0(g) = g \oplus \Delta_0(g)$. Then, clearly Γ_0 is $(n, 1)$-computable via $\{\lambda g. g \oplus \Delta_i(g)\}_{i<n}$. For $g \oplus h \in \mathbb{N}^\mathbb{N}$, if $\Gamma(g) = \Delta_i(g)$ for some $i < n$, then $\Gamma_1(g \oplus h) = h$. Otherwise, we set $\Gamma_1(g \oplus h) = \Delta_n(g)$. Clearly, Γ_1 is $(2,1)$-computable. Note that, if $g \in \mathrm{dom}(\Gamma)$, then $\Gamma(g) = \Delta_i(g)$ for some $i \leq n$. If $\Gamma(g) = \Delta_i(g)$ for some $i < n$, then $\Gamma_1(\Gamma_0(g)) = \Gamma_1(g \oplus \Delta_i(g)) = \Delta_i(g)$. If $\Gamma(g) = \Delta_n(g)$, then $\Gamma_1(\Gamma_0(g)) = \Gamma_1(g \oplus \Delta_0(g)) = \Delta_n(g)$. Therefore, $\Gamma(g) = \Gamma_1 \circ \Gamma_0(g)$ for any $g \in \mathrm{dom}(\Gamma)$. By the similar way, it is easy to see that every $(n+1, \omega)$-computable function Γ can be represented as $\Gamma = \Gamma_1 \circ \Gamma_0$ for some (n, ω)-computable function Γ_0 and $(2, \omega)$-computable function Γ_1. □

Corollary 32. *$\{[\mathfrak{C}_T]^{\alpha^-}_{\beta^-|\gamma^-} : \alpha, \beta, \gamma \leq \omega+1\}$ contains just seven monoids, $[\mathfrak{C}_T]^1_1$, $[\mathfrak{C}_T]^1_{<\omega}$, $[\mathfrak{C}_T]^1_{\omega|<\omega}$, $[\mathfrak{C}_T]^{<\omega}_1$, $[\mathfrak{C}_T]^1_\omega$, $[\mathfrak{C}_T]^{<\omega}_\omega$, and $[\mathfrak{C}_T]^\omega_1$.*

Proof. By Proposition 26, we have $[\mathfrak{C}_T]^1_{1|1} = [\mathfrak{C}_T]^1_{1|<\omega} = [\mathfrak{C}_T]^1_{1|\omega} = [\mathfrak{C}_T]^1_{<\omega|1} = [\mathfrak{C}_T]^1_{\omega|1}$; $[\mathfrak{C}_T]^1_{<\omega|<\omega} = [\mathfrak{C}_T]^1_{1|<\omega}$; and $[\mathfrak{C}_T]^\omega_{1|1} = [\mathfrak{C}_T]^\omega_{\beta|\gamma}$ for any $\beta, \gamma \in \{1, <\omega, \omega\}$. Moreover, by Proposition 26 and Proposition 27, $[\mathfrak{C}_T]^{<\omega}_{1|1} = [\mathfrak{C}_T]^{<\omega}_{\beta|\gamma}$ whenever $\langle \beta, \gamma \rangle \neq \langle \omega, \omega \rangle$. Therefore, by Proposition 29 and 31, we have just seven monoids, $[\mathfrak{C}_T]^1_1$, $[\mathfrak{C}_T]^1_{<\omega}$, $[\mathfrak{C}_T]^1_{\omega|<\omega}$, $[\mathfrak{C}_T]^{<\omega}_1$, $[\mathfrak{C}_T]^1_\omega$, $[\mathfrak{C}_T]^{<\omega}_\omega$, and $[\mathfrak{C}_T]^\omega_1$. □

When the classical disjunction $P \cup Q$ (resp. $P \triangledown_\infty Q$) is valid, say $f \in P \cup Q$ (resp. $f \in P \triangledown_\infty Q$), then either $f \mapsto 0^\frown f$ or $f \mapsto 1^\frown f$ (resp. either $f \mapsto 0^\frown \mathtt{pr}_0(f)$ or $f \mapsto 1^\frown \mathtt{pr}_1(f)$) witness the validity of $P \oplus Q$

$$
\begin{array}{ccccccccccc}
 & & & & & \rotatebox{30}{$\subsetneq$} & [\mathfrak{C}_T]^{<\omega}_1 = [\mathfrak{C}_T]^{<\omega}_{\omega|<\omega} & \rotatebox{-30}{$\subsetneq$} & & & \\
[\mathfrak{C}_T]^1_1 & \subseteq & [\mathfrak{C}_T]^1_{<\omega} & \subseteq & [\mathfrak{C}_T]^1_{\omega|<\omega} & & & & [\mathfrak{C}_T]^{<\omega}_{\omega} & \subseteq & [\mathfrak{C}_T]^{\omega}_1 = [\mathfrak{C}_T]^{\omega}_{\omega} \\
 & & & & & \rotatebox{-30}{$\subsetneq$} & [\mathfrak{C}_T]^1_{\omega} & \rotatebox{30}{$\subsetneq$} & & &
\end{array}
$$

TABLE 3. Seven monoids of nonuniformly computable functions

intuitionistically. That is to say, the one-tape (resp. two-tape) intuitionistic disjunction is reducible to the classical disjunction via two total (resp. partial) algorithms. In general, we will see that each one-tape disjunction is decomposed by a total (i.e., truth-table, or Popperian) learners.

Definition 33. Let D be a subset of Baire space $\mathbb{N}^{\mathbb{N}}$, and $\alpha, \beta, \gamma \leq \omega + 1$ be ordinals. A function $\Gamma : D \to \mathbb{N}^{\mathbb{N}}$ is $(\alpha^-, \beta^-|\gamma^-)$-*truth-table* if there are bounds $a < \alpha$, $z < \min\{\beta, \gamma\}$, a set $I \subseteq \mathbb{N}$ of cardinality a, and a collection $\{p(e,k) : e \in I \ \&\ k < z\}$ of indices of *truth-table functionals* (i.e., $\mathrm{dom}(\Phi_{p(e,k)}) = \mathbb{N}^{\mathbb{N}}$) such that

1. (Popperian Condition) for any $e \in I$, there is $k < z$ such that $\Psi_e(\sigma) = p(e,k)$.
2. Γ is $(\alpha^-, \beta^-|\gamma^-)$-computable via the family $\{\Psi_e\}_{e \in I}$.

Here, we do not assume the uniform computability of the collection $\{p(e,k) : e \in I \ \&\ k < z\}$. If $\gamma = \omega$, then we simply say that Γ is (α, β)-*truth-table* for $(\alpha, \beta|\gamma)$-truth-table function Γ. Let $[\mathfrak{C}_{tt}]^{\alpha}_{\beta}$ (resp. $[\mathfrak{C}_{tt}]^{\alpha}_{\beta|\gamma}$) denote the set of all (α, β)-truth-table (resp. $(\alpha, \beta|\gamma)$-truth-table) functions.

Proposition 34. $\aleph_0 = \#[\mathfrak{C}_r]^1_1 = \#[\mathfrak{C}_r]^1_{<\omega} = \#[\mathfrak{C}_r]^1_{\omega|<\omega} = \#[\mathfrak{C}_r]^1_{\omega} < \#[\mathfrak{C}_r]^{<\omega}_1 = \#[\mathfrak{C}_r]^{<\omega}_{\omega} = \#[\mathfrak{C}_r]^{\omega}_1 = 2^{2^{\aleph_0}}$, *for each* $r \in \{tt, T\}$.

Proof. Every learner Ψ determines just one learnable function $\Gamma \in [\mathfrak{C}_T]^1_{\omega}$. Therefore, $[\mathfrak{C}_T]^1_{\omega}$ is countable. For non-uniform computability, we first see $\#[\mathfrak{C}_T]^{\omega}_1 \leq 2^{2^{\aleph_0}}$ since $\#(\mathbb{N}^{\mathbb{N}})^{\mathbb{N}^{\mathbb{N}}} = 2^{2^{\aleph_0}}$ by cardinal arithmetic. On the other hand, every function $\Gamma : \mathbb{N}^{\mathbb{N}} \to \{0^{\mathbb{N}}, 1^{\mathbb{N}}\}$ is $(<\omega, 1)$-truth-table via two constant truth-table functionals $\Gamma_0(f) = 0^{\mathbb{N}}$ and $\Gamma_1(f) = 1^{\mathbb{N}}$ for any $f \in \mathbb{N}^{\mathbb{N}}$. Therefore, $\#[\mathfrak{C}_{tt}]^{<\omega}_1 \geq 2^{2^{\aleph_0}}$. □

Proposition 35. *Let P, Q be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

1. *There is a $(1, n|2)$-truth-table function $\Gamma : [\![P \vee Q]\!]^1_{\mathsf{LCM}[n]} \to P \oplus Q$.*
2. *There is a $(1, n|2)$-computable function $\Gamma : [\![P \vee Q]\!]^2_{\mathsf{LCM}[n]} \to P \oplus Q$.*
3. *There is a $(1, n)$-computable function $\Gamma : [\![P \vee Q]\!]^3_{\mathsf{LCM}[n]} \to P \oplus Q$.*
4. *There is a $(1, \omega|2)$-truth-table function $\Gamma : [\![P \vee Q]\!]^1_{\mathsf{LCM}} \to P \oplus Q$.*
5. *There is a $(1, \omega|2)$-computable function $\Gamma : [\![P \vee Q]\!]^2_{\mathsf{LCM}} \to P \oplus Q$.*
6. *There is a $(1, \omega)$-computable function $\Gamma : [\![P \vee Q]\!]^3_{\mathsf{LCM}} \to P \oplus Q$.*
7. *There is a $(2, 1)$-truth-table function $\Gamma : [\![P \vee Q]\!]^1_{\mathsf{CL}} \to P \oplus Q$.*
8. *There is a $(2, 1)$-computable function $\Gamma : [\![P \vee Q]\!]^2_{\mathsf{CL}} \to P \oplus Q$.*

Proof. For the items (1), (4), and (7), we consider the truth-table functionals $\Delta_0 : f \oplus g \mapsto 0^\frown g$ and $\Delta_1 : f \oplus g \mapsto 1^\frown g$. By the definition of $[\![P \vee Q]\!]^1_{\mathsf{CL}}$, obviously $\Delta_0(f \oplus g) \in P \oplus Q$ or $\Delta_1(f \oplus g) \in P \oplus Q$ for any $f \oplus g \in [\![P \vee Q]\!]^1_{\mathsf{CL}}$. Let e_0 and e_1 be indices of Δ_0 and Δ_1, respectively. On $\sigma \oplus \tau \in (2 \times \mathbb{N})^{<\mathbb{N}}$, we set $\Psi(\sigma \oplus \tau) = e_{\sigma(|\sigma|-1)}$. Note that the partial function Γ identified by the learner Ψ is $(1, n|2)$-truth-table on $[\![P \vee Q]\!]^1_{\mathsf{LCM[n]}}$, and $(1, \omega|2)$-truth-table on $[\![P \vee Q]\!]^1_{\mathsf{LCM}}$. Moreover, clearly $\Gamma(f \oplus g) = (\lim_s f(s))^\frown g \in P \oplus Q$ for every $f \oplus g \in [\![P \vee Q]\!]^1_{\mathsf{LCM}}$.

For the items (2), (5), and (8), we consider the partial computable functions $\Delta_0 : f \mapsto 0^\frown \mathtt{pr}_0(f)$ and $\Delta_1 : f \mapsto 1^\frown \mathtt{pr}_1(f)$. By the definition of $[\![P \vee Q]\!]^2_{\mathsf{CL}}$, obviously $\Delta_0(f) \in P \oplus Q$ or $\Delta_1(f) \in P \oplus Q$ for any $f \in [\![P \vee Q]\!]^2_{\mathsf{CL}}$. Let e_0 and e_1 be indices of Δ_0 and Δ_1, respectively. On $\sigma \in (2 \times \mathbb{N})^{<\mathbb{N}}$, we set $\Psi(\sigma) = e_{(\sigma(|\sigma|-1))_0}$. Note that the partial function Γ identified by the learner Ψ is $(1, n|2)$-computable on $[\![P \vee Q]\!]^2_{\mathsf{LCM[n]}}$, and $(1,\omega|2)$-computable on $[\![P \vee Q]\!]^2_{\mathsf{LCM}}$. Moreover, clearly $\Gamma(f) \in P \oplus Q$ for every $f \in [\![P \vee Q]\!]^2_{\mathsf{LCM}}$.

For the items (3) and (6), on $\sigma \in (\mathbb{N} \cup \{\sharp\})^{<\omega}$, $\Psi(\sigma)$ guesses an index of the partial computable function $g \mapsto g^{\leftharpoonup t(\sigma)}$, where $t(\sigma) = \max\{n : \sigma(n) = \sharp\} + 1$ if such n exists; otherwise, $t(\sigma) = 0$. Note that the partial function Γ identified by the learner Ψ is $(1,n)$-computable on $[\![P \vee Q]\!]^3_{\mathsf{LCM[n]}}$, and $(1,\omega)$-computable on $[\![P \vee Q]\!]^3_{\mathsf{LCM}}$. Moreover, clearly $\Gamma(f) \in P \oplus Q$ for every $f \in [\![P \vee Q]\!]^3_{\mathsf{LCM}}$. ◻

Definition 36. Let P, Q be subsets of $\mathbb{N}^{\mathbb{N}}$, and fix $\alpha, \beta, \gamma \in \{1, <\omega, \omega\}$. Then, *$P$ is $(\alpha,\beta|\gamma)$-reducible to Q* (written $P \leq^{\alpha}_{\beta|\gamma} Q$) if there is $\Gamma \in [\mathfrak{C}_T]^{\alpha}_{\beta|\gamma}$ such that $\Gamma : Q \to P$. We say that *P is $(\alpha,\beta|\gamma)$-equivalent to Q* (written $P \equiv^{\alpha}_{\beta|\gamma} Q$) if $P \leq^{\alpha}_{\beta|\gamma} Q$ and $Q \leq^{\alpha}_{\beta|\gamma} P$. We also say that *$P$ is $(\alpha,\beta|\gamma)$-truth-table-reducible to Q* (written $P \leq^{\alpha}_{tt,\beta|\gamma} Q$) if there is $\Gamma \in [\mathfrak{C}_{tt}]^{\alpha}_{\beta|\gamma}$ such that $\Gamma : Q \to P$, and that *P is $(\alpha,\beta|\gamma)$-truth-table-equivalent to Q* (written $P \equiv^{\alpha}_{tt,\beta|\gamma} Q$) if $P \leq^{\alpha}_{tt,\beta|\gamma} Q$ and $Q \leq^{\alpha}_{tt,\beta|\gamma} P$.

Proposition 37. *Fix $\alpha, \beta, \gamma \in \{1, <\omega, \omega\}$. Then, $\leq^{\alpha}_{\beta|\gamma}$ forms a preordering over $\mathcal{P}(\mathbb{N}^{\mathbb{N}})$.*

Proof. By Corollary 30. ◻

Proposition 38. *The preorderings $\leq^1_1$ and $\leq^{\omega}_1$ are equivalent to the Medvedev reducibility and the Muchnik reducibility, respectively.*

Proof. By Proposition 26 (4) and (5). ◻

Definition 39. Fix $\alpha, \beta, \gamma \in \{1, <\omega, \omega\}$. Let $\mathcal{D}^{\alpha}_{\beta|\gamma}$, $\mathcal{D}^{\alpha}_{tt,\beta|\gamma}$, $\mathcal{P}^{\alpha}_{\beta|\gamma}$, and $\mathcal{P}^{\alpha}_{tt,\beta|\gamma}$ denote the quotient sets $\mathcal{P}(\mathbb{N}^{\mathbb{N}})/\equiv^{\alpha}_{\beta|\gamma}$, $\mathcal{P}(\mathbb{N}^{\mathbb{N}})/\equiv^{\alpha}_{tt,\beta|\gamma}$, $\Pi^0_1(2^{\mathbb{N}})/\equiv^{\alpha}_{\beta|\gamma}$, and $\Pi^0_1(2^{\mathbb{N}})/\equiv^{\alpha}_{tt,\beta|\gamma}$, respectively. Here, $\Pi^0_1(2^{\mathbb{N}})$ denotes the set of all nonempty Π^0_1 subsets of $2^{\mathbb{N}}$. For $P \in \mathcal{P}(\mathbb{N}^{\mathbb{N}})$, the equivalence class $\{Q \subseteq \mathbb{N}^{\mathbb{N}} : Q \equiv^{\alpha}_{\beta|\gamma} P\} \in \mathcal{D}^{\alpha}_{\beta|\gamma}$ is called *the $(\alpha,\beta|\gamma)$-degree* of P.

Proposition 40. *For each $\alpha, \beta, \gamma \in \{1, <\omega, \omega\}$, the order structures $\mathcal{D}^{\alpha}_{\beta|\gamma}$, $\mathcal{D}^{\alpha}_{tt,\beta|\gamma}$, $\mathcal{P}^{\alpha}_{\beta|\gamma}$, and $\mathcal{P}^{\alpha}_{tt,\beta|\gamma}$ form lattices. Moreover, the disjunction operations $[\![\cdot \vee \cdot]\!]^1_{\mathsf{LCM[2]}}$, $[\![\cdot \vee \cdot]\!]^2_{\mathsf{LCM[2]}}$, $[\![\cdot \vee \cdot]\!]^1_{\mathsf{LCM}}$, $[\![\cdot \vee \cdot]\!]^2_{\mathsf{LCM}}$, $[\![\cdot \vee \cdot]\!]^3_{\mathsf{LCM}}$, $[\![\cdot \vee \cdot]\!]^1_{\mathsf{CL}}$, and $[\![\cdot \vee \cdot]\!]^2_{\mathsf{CL}}$ are the infimum operations on the structures $\mathcal{D}^1_{tt,<\omega}$, $\mathcal{D}^1_{<\omega}$, $\mathcal{D}^1_{tt,\omega|<\omega}$, $\mathcal{D}^1_{\omega|<\omega}$, $\mathcal{D}^1_{\omega}$, $\mathcal{D}^{<\omega}_{tt,1}$, and $\mathcal{D}^{<\omega}_{1}$, respectively. In particular, $\cup$, $\triangledown$, $\triangledown_{\omega}$, and $\triangledown_{\infty}$ are the infimum operations on the structures $\mathcal{P}^{<\omega}_{tt,1}$, $\mathcal{P}^1_{<\omega}$, $\mathcal{P}^1_{\omega|<\omega}$, and $\mathcal{P}^{<\omega}_1$, respectively.*

Proof. By Proposition 35. ◻

Remark. Later we provide a characterization of our degree structures by iterations of the disjunction operations, in Theorem 156.

Proposition 41. *The degree structures $\mathcal{D}^1_{\omega}$ and $\mathcal{D}^1_{tt,\omega}$ are Brouwerian.*

Proof. Set $B(P,Q) = \{R \subseteq \mathbb{N}^{\mathbb{N}} : P \leq^1_{\omega} Q \otimes R\}$. We need to a construct a function $\beta : \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \times \mathcal{P}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$ such that $\beta(P,Q) = \min B(P,Q)$ for any $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$. Let Λ_e denote the e-th $(1,\omega)$-computable function, i.e., $\Lambda_e(g) = \Phi_{\lim_n \Psi_e(g \upharpoonright n)}(g)$ for any $g \in \mathrm{dom}(\Lambda_e)$. Define β as follows.

$$\beta(P,Q) = \{e^\frown g \in \mathbb{N}^{\mathbb{N}} : (\forall f \in Q)\ \Lambda_e(f \oplus g) \in P\}.$$

It is easy to see that $\beta(P,Q) \in B(P,Q)$ for any $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$. If $R \in B(P,Q)$, say $\Lambda_e : Q \otimes R \to P$, then clearly $e^\frown g \in \beta(P,Q)$ for any $g \in R$. Thus, $\beta(P,Q) \leq^1_1 R$. ◻

Remark. In contrast, later we will show that *neither $\mathcal{D}^1_{<\omega}$, nor $\mathcal{D}^1_{\omega|<\omega}$, nor $\mathcal{D}^{<\omega}_1$, nor $\mathcal{D}^{<\omega}_{\omega}$* form Brouwer algebras. See also Theorem 194 and 196. In the meantime, by Proposition 49, it is shown that there are Brouwerian modifications of $\mathcal{D}^1_{<\omega}$, $\mathcal{D}^1_{\omega|<\omega}$, $\mathcal{D}^{<\omega}_1$, and $\mathcal{D}^{<\omega}_{\omega}$ which look more natural than our original definitions, from the viewpoint of constructive mathematics.

For a sequence $\{\sigma_n\}_{n\in\mathbb{N}} \in (\mathbb{N}^{<\mathbb{N}})^{\mathbb{N}}$ of strings, $\lim_n \sigma_n$ is defined by $(\lim_n \sigma_n)(m) = \lim_n(\sigma_n(m))$. If $\lim_n \sigma_n : \mathbb{N} \to \mathbb{N}$ is total, say $\lim_n \sigma_n = h \in \mathbb{N}^{\mathbb{N}}$, then we say that $\lim_n \sigma_n \in \mathbb{N}^{\mathbb{N}}$ *converges to* h.

Proposition 42. *Fix an ordinal* $\alpha \leq \omega$. *A partial function* $\Gamma :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ *is* $(1,\alpha)$*-computable if and only if there is a total computable function* $\psi : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$ *such that* $\lim_n \psi(g \upharpoonright n)$ *converges to* $\Gamma(g)$, *and* $\#\{n \in \mathbb{N} : \psi(g \upharpoonright n+1) \not\supseteq \psi(g \upharpoonright n)\} < \alpha$, *for any* $g \in \mathrm{dom}(\Gamma)$.

Proof. Assume that Γ is $(1,\alpha)$-computable via a learner Ψ. We put $\psi(\sigma) = \Phi_{\Psi(\sigma)}(\sigma)$ for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Then the condition $\#\{n \in \mathbb{N} : \Psi(g \upharpoonright n+1) \neq \Psi(g \upharpoonright n)\} < \alpha$ implies $\#\{n \in \mathbb{N} : \psi(g \upharpoonright n+1) \not\supseteq \psi(g \upharpoonright n)\} < \alpha$, for any $g \in \mathrm{dom}(\Gamma)$. Because if $\Psi(g \upharpoonright n+1) = \Psi(g \upharpoonright n)$, then $\psi(g \upharpoonright n) = \Phi_{\Psi(g\upharpoonright n)}(g \upharpoonright n) \subseteq \Phi_{\Psi(g\upharpoonright n)}(g \upharpoonright n+1) = \psi(g \upharpoonright n+1)$. Thus, clearly, $\lim_n \psi(g \upharpoonright n)$ converges to $\Phi_{\lim_n \Psi(g\upharpoonright n)}(g) = \Gamma(g)$.

Assume that $\Gamma(g) = \lim_n \psi(g \upharpoonright n)$ for any $g \in \mathrm{dom}(\Gamma)$ for some ψ satisfying the condition in Proposition 42. We define a computable function $\Phi_{e(\sigma)} : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$. For any $g \in \mathbb{N}^{\mathbb{N}}$, put $\Phi_{e(\sigma)}(g; n) = \psi(g \upharpoonright s)(n)$ for each $n \in \mathbb{N}$, where $s \geq |\sigma|$ is the least number such that $\psi(g \upharpoonright s)(n)$ is defined. Clearly, $\Phi_{e(\sigma)}$ is partial computable, and indeed, we can compute an index $e(\sigma)$ of $\Phi_{e(\sigma)}$ uniformly in $\sigma \in \mathbb{N}^{<\mathbb{N}}$. Then, we define a learner Ψ inductively. Put $\Psi(\langle\rangle) = e(\langle\rangle)$. Fix $\sigma \in \mathbb{N}^{<\mathbb{N}}$, and assume that $\Psi(\sigma^-)$ has already been defined. If $\psi(\sigma) \supseteq \psi(\sigma^-)$, then set $\Psi(\sigma) = \Psi(\sigma^-)$. If $\psi(\sigma) \not\supseteq \psi(\sigma^-)$, then set $\Psi(\sigma) = e(\sigma)$. Clearly, the condition $\#\{n \in \mathbb{N} : \psi(g \upharpoonright n+1) \not\supseteq \psi(g \upharpoonright n)\} < \alpha$ implies $\#\{n \in \mathbb{N} : \Psi(g \upharpoonright n+1) \neq \Psi(g \upharpoonright n)\} < \alpha$, for any $g \in \mathrm{dom}(\Gamma)$. In particular, $\lim_n \Psi(g \upharpoonright n)$ converges to some index $e(\sigma)$ for any $g \in \mathrm{dom}(\Gamma)$. Hence, $\Phi_{\lim_n \Psi(g\upharpoonright n)}(g) = \bigcup_{n\geq|\sigma|} \psi(g \upharpoonright n) = \lim_{n\in\mathbb{N}} \psi(g \upharpoonright n) = \Gamma(g)$, since $\{\psi(g \upharpoonright n)\}_{n\geq|\sigma|}$ is an increasing sequence of strings. □

Corollary 43 (de Brecht-Yamamoto [26])**.** *A partial function* $\Gamma :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ *is* $(1,\omega)$*-computable if and only if there is a computable sequence* $\{\Gamma_n\}_{n\in\mathbb{N}}$ *of partial computable functions which converges pointwise to* Γ *on* $\mathrm{dom}(\Gamma)$ *with respect to the discrete topology on* $\mathbb{N}^{\mathbb{N}}$.

Proof. By Proposition 42. □

Remark. Proposition 42 suggests that every $(1,\omega)$-computable function $\Gamma :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is also of Baire class 1 (indeed, effectively Σ^0_2-measurable) with respect to both the discrete topology and the standard Baire topology on $\mathbb{N}^{\mathbb{N}}$. Here, the effective Σ^0_2-measurability is also known as the limit computability [10]. However, on Baire topology, not every effectively Σ^0_2-measurable function is $(1,\omega)$-computable. In fact, we can easily construct a computable function with a $\mathbf{0}'$-oracle (which must be $\Sigma^{0,\mathbf{0}'}_1$-measurable) which maps a computable point to a $\mathbf{0}'$-computable point, whereas every $(1,\omega)$-computable function Γ is non-uniformly computable, i.e., $\Gamma(x) \leq_T x$ for any $x \in \mathrm{dom}(\Gamma)$, as seen in Proposition 26. Moreover, not every $(1,\omega)$-computable function is $\mathbf{0}'$-computable, since we can also easily construct a discontinuous $(1,\omega)$-computable function while every function computable relative to some oracle must be continuous. Therefore, in the context of computability on Baire topology, $(1,\omega)$*-computability (learnability) is quite different from* $\mathbf{0}'$*-computability and limit computability*. But, by this observation, the notion of learnability is rather expected to be a useful tool to the study of effective discontinuous functions.

2.5. **Nonuniform Computability on Falsifiable Problems.**

In this paper, we will mainly pay attention to the behavior of nonuniform computability on Π^0_1 *subsets of Cantor space* $2^{\mathbb{N}}$. We first see that, if we restrict our attention to Π^0_1 sets, then some reducibility notions collapse.

Proposition 44. *Let* P *be a* Π^0_1 *subset of* $\mathbb{N}^{\mathbb{N}}$, *and* X *be any subset of* $\mathbb{N}^{\mathbb{N}}$.

(1) $X \leq^1_{tt,1} P$ *if and only if* $X \leq^1_1 P$.
(2) $X \leq^1_{tt,<\omega} P$ *if and only if* $X \leq^1_{<\omega} P$.
(3) $X \leq^1_{tt,\omega} P$ *if and only if* $X \leq^1_{\omega} P$.
(4) $P \leq^1_{tt,<\omega} X$ *if and only if* $P \leq^{<\omega}_{tt,\omega|<\omega} X$.
(5) $P \leq^1_{tt,\omega} X$ *if and only if* $P \leq^{<\omega}_{tt,\omega} X$.

Proof. (1) See Simpson [61].

(2,3) Assume that $X \leq^1_\omega Q$ via a learner Ψ. From Ψ, we construct a Popperian learner $\Psi^* : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$, i.e., $\Psi(\sigma)$ is an index of truth-table functional for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$. We may assume that $\Psi(\sigma)$ is defined, by Proposition 24. Let T_P be the corresponding computable tree for P. If $\sigma \notin T_P$, then $\Psi^*(\sigma)$ returns an index of the constant function $f \mapsto 0^{\mathbb{N}}$. If $\sigma \in T_P$, then let $\Psi^*(\sigma)$ be an index of the following computation procedure. Given $f \in \mathbb{N}^{\mathbb{N}}$, at stage $s \in \mathbb{N}$, if $\sigma \not\subset f$, then returns $0^{\mathbb{N}}$. If $f \upharpoonright s \in T_P$ extends σ, and $\Psi(f \upharpoonright t) = \sigma$ for any $|\sigma| \leq t \leq s$, then simulate the computation of $\Phi_{\Psi(\sigma)}(f \upharpoonright s)$. Otherwise, for the least such stage s, returns $\Phi_{\Psi(\sigma)}(f \upharpoonright s - 1)^\frown 0^{\mathbb{N}}$. Clearly, $\Phi_{\Psi^*(\sigma)}(f)$ defines an element of $\mathbb{N}^{\mathbb{N}}$, for any $f \in \mathbb{N}^{\mathbb{N}}$. Moreover, Ψ^* agrees with Ψ on P, i.e., $\Phi_{\lim_n \Psi^*(f \upharpoonright n)}(f) = \Phi_{\lim_n \Psi(f \upharpoonright n)}(f)$ for any $f \in P$.

(4,5) Assume that $P \leq^{<\omega}_{tt,\omega|<\omega} X$ via n Popperian learners, $\{\Psi_i\}_{i<n}$. Given $g \in X$, on the first challenge, our leaner Δ guesses that $\Psi_0(g \upharpoonright 0)$ is a correct algorithm. As each Ψ_i is Popperian, and P is Π^0_1, the predicate $\Phi_{\Psi_0(g \upharpoonright 0)}(g) \in P$ is Π^0_1. Therefore, whenever $\Phi_{\Psi_0(g \upharpoonright 0)}(g) \in P$ is incorrect, the learner Δ is able to understand that his guess is refuted. If it happens, the learner goes to the next challenge. On the $(ns + i)$-th challenge, Δ guesses that $\Psi_i(g \upharpoonright s)$ is correct. By continuing this procedure, eventually Δ learns a collect algorithm to solve the problem P. Note that, if an (n, b, c)-computable function exists from X to P, then the learning procedure of Δ is stabilized before the (nc)-th challenge starts, i.e., Δ determines a $(1, nc)$-truth-table computable function. □

Corollary 45. *$\{\leq^\alpha_{\beta|\gamma} \cap (\Pi^0_1 \times \Pi^0_1), \leq^\alpha_{tt,\beta|\gamma} \cap (\Pi^0_1 \times \Pi^0_1) : \alpha, \beta, \gamma \in \{1, <\omega, \omega\}\}$ contains at most nine preorderings: $\leq^1_1$, $\leq^{<\omega}_{tt,1}$, $\leq^1_{<\omega}$, $\leq^1_{\omega|<\omega}$, $\leq^{<\omega}_1$, $\leq^1_\omega$, $\leq^{<\omega}_{<\omega}$, $\leq^\omega_{tt,1}$, and $\leq^\omega_1$.* □

Remark. We may introduce the notion of Popperian learning on Baire space $\mathbb{N}^{\mathbb{N}}$. Let $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, and $f : Q \to P$ is a learnable function via a learner Ψ. The procedure of the learner Ψ is described as follows. On an observation $\sigma \in \mathbb{N}^{<\mathbb{N}}$ with $[\sigma] \cap Q \neq \emptyset$, a learner Ψ conjectures that e is a correct algorithm computing a solution of P from σ, that is, $\Phi_{\Psi(\sigma)}(f) = \Phi_e(f) \in P$ for any future observation $f \in Q \cap [\sigma]$. If Q is Π^0_1, Proposition 44 (3) suggests that we may assume that e is an index of a total computable function. Then, the learner Ψ can find mistakes of his hypothesis on P if P is also a Π^0_1 subset of the Baire space $\mathbb{N}^{\mathbb{N}}$. Therefore, restricting to Π^0_1 subsets is expected to be an analogy of *Popperian learning*. In this context, the usual Popperian learning on total computable functions could be regarded as a learning process on Π^0_1 singletons.

Definition 46. Let D be a subset of Baire space $\mathbb{N}^{\mathbb{N}}$, and $\alpha, \beta, \gamma \leq \omega + 1$ be ordinals, or $\mathtt{eff}$, where $\mathtt{eff}^- = \mathtt{eff}$. We generalize the $(\alpha^-, \beta^-, \gamma^-)$-computability as follows. If $\alpha = \mathtt{eff}$, then we revise the word "for any $g \in D$, there is $e \in I$" to the word "there is a partial computable function $B_0 :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ such that, for any $g \in D$, for $e = B_0(g)$". If $\beta = \mathtt{eff}$, then we revise the mind change condition as $\#\{n \in \mathbb{N} : \Psi_e(g \upharpoonright n + 1) \neq \Psi_e(g \upharpoonright n)\} < B_1(g)$, where B_1 is a computable function from $\mathbb{N}^{\mathbb{N}}$ to $\mathbb{N}$. If $\gamma = \mathtt{eff}$, then we revise the error condition as $\#\{\Psi_e(g \upharpoonright n) : n \in \mathbb{N}\} < B_2(g)$, where B_2 is a computable function from $\mathbb{N}^{\mathbb{N}}$ to $\mathbb{N}$. For new notions, $\leq^\alpha_{\beta|\gamma}$, $\mathcal{D}^\alpha_{\beta|\gamma}$, and $\mathcal{P}^\alpha_{\beta|\gamma}$ are also defined as the usual way.

Proposition 47. *Suppose that, if $\tau = \mathtt{eff}$, then let τ^* mean the symbol $<\omega$, and otherwise, set $\tau^* = \tau$. Then, every (α, β, γ)-computable function with a compact domain is $(\alpha^*, \beta^*, \gamma^*)$-computable.*

Proof. By continuity of B_0, B_1, and B_2 in Definition 46, $\{B_i^{-1}(\{e\})\}_{e \in \mathbb{N}}$ for each $i < 3$ is an open cover of D. Hence, by compactness of D, we have the desired condition. □

Corollary 48. *$\mathcal{P}^1_{\mathtt{eff}} = \mathcal{P}^1_{<\omega}$; $\mathcal{P}^1_{\omega|\mathtt{eff}} = \mathcal{P}^1_{\omega|<\omega}$; $\mathcal{P}^{\mathtt{eff}}_1 = \mathcal{P}^{<\omega}_1$; and $\mathcal{P}^{\mathtt{eff}}_\omega = \mathcal{P}^{<\omega}_\omega$.* □

That is to say, for Π^0_1 subsets of Cantor space $2^{\mathbb{N}}$, no new reducibility notion is constructed from Definition 46. However, from the perspective of intuitionistic caluculus, our new notions in Definition 46 have nice features.

Proposition 49. *$\mathcal{D}^1_{\mathtt{eff}}$, $\mathcal{D}^1_{\omega|\mathtt{eff}}$, $\mathcal{D}^{\mathtt{eff}}_1$, and $\mathcal{D}^{\mathtt{eff}}_\omega$ are Brouwerian.*

Proof. Fix $\alpha, \beta, \gamma \in \{1, <\omega, \mathtt{eff}, \omega\}$, and set $B(P, Q) = \{R \subseteq \mathbb{N}^\mathbb{N} : P \leq^{\alpha}_{\beta|\gamma} Q \otimes R\}$. We need to construct a function $\beta : \mathcal{P}(\mathbb{N}^\mathbb{N}) \times \mathcal{P}(\mathbb{N}^\mathbb{N}) \to \mathcal{P}(\mathbb{N}^\mathbb{N})$ such that $\beta(P, Q) = \min B(P, Q)$ for any $P, Q \subseteq \mathbb{N}^\mathbb{N}$. Let Λ_e denote the e-th $(1, \omega)$-computable function, and Θ_e be the e-th partial computable function from $\mathbb{N}^\mathbb{N}$ to $\mathbb{N}$. Put $\mathrm{change}_e(g) = \#\{n \in \mathbb{N} : \Lambda_e(g \upharpoonright n+1) \neq \Lambda_e(g \upharpoonright n)\}$, and $\mathrm{error}_e(g) = \#\{\Lambda_e(g \upharpoonright n) : n \in \mathbb{N}\}$. Then,

$$\beta(P, Q) = \begin{cases} \{(e,d)^\frown g : (\forall f \in Q)\ \Lambda_e(f \oplus g) \in P \ \&\ \mathrm{change}_e(f \oplus g) < \Theta_d(f \oplus g)\}, \\ \hfill \text{if } (\alpha, \beta, \gamma) = (1, \mathtt{eff}, \omega), \\ \{(e,d)^\frown g : (\forall f \in Q)\ \Lambda_e(f \oplus g) \in P \ \&\ \mathrm{error}_e(f \oplus g) < \Theta_d(f \oplus g)\}, \\ \hfill \text{if } (\alpha, \beta, \gamma) = (1, \omega, \mathtt{eff}), \\ \{d^\frown g : (\forall f \in Q)(\exists e < \Theta(f \oplus g))\ \Phi_e(f \oplus g) \in P\}, \\ \hfill \text{if } (\alpha, \beta, \gamma) = (\mathtt{eff}, 1, \omega), \\ \{d^\frown g : (\forall f \in Q)(\exists e < \Theta(f \oplus g))\ \Lambda_e(f \oplus g) \in P\}, \\ \hfill \text{if } (\alpha, \beta, \gamma) = (\mathtt{eff}, \omega, \omega), \end{cases}$$

It is easy to see that $\beta(P, Q) \in B(P, Q)$ for any $P, Q \subseteq \mathbb{N}^\mathbb{N}$. For the minimality, if $R \in B(P, Q)$, we have suitable d and e such that $(d, e)^\frown g \in \beta(P, Q)$ for any $g \in R$. Thus, $\beta(P, Q) \leq^1_1 R$. □

Remark. Unfortunately, neither $\mathcal{P}^1_{\mathtt{eff}}$ nor $\mathcal{P}^1_{\omega|\mathtt{eff}}$ nor $\mathcal{P}^{\mathtt{eff}}_1$ nor $\mathcal{P}^{\mathtt{eff}}_\omega$ form Brouwer algebra. See Theorem 194 and 196.

The Medvedev reducibility $\leq^1_1$, the learnable reducibility $\leq^1_\omega$, and the Muchnik reducibility $\leq^\omega_1$ could be viewed as analogues of the intuitionistic implication, the LCM implication and the classical implication, respectively. In this way, $\leq^1_1$ is interpreted as algorithmical reducibility with no mind-changes. Then, how the classical reducibility $\leq^\omega_1$ behave as a learning procedure?

Proposition 50. *For any Π^0_1 set $P \subseteq \mathbb{N}^\mathbb{N}$ and any set $Q \subseteq \mathbb{N}^\mathbb{N}$, $P \leq^\omega_1 Q$ if and only if*

$$(\exists \Psi)(\forall f \in Q)\ \Phi_{\liminf_n \Psi(f \upharpoonright n)}(f) \in P.$$

Here Ψ ranges over all learners (i.e., computable functions from $\mathbb{N}^{<\mathbb{N}}$ to $\mathbb{N}$).

Proof. The "only if" part is obvious. For the "if" part, we will inductively define $\Psi(\sigma)$ and $l(\sigma, e)$ for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $e \in \mathbb{N}$. Let T_P denote the corresponding tree for P. First, put $\Psi(\langle\rangle) = 0$ and $l(\langle\rangle, e) = 0$ for each e. Now assume that, for any $\tau \in \mathbb{N}^{<\mathbb{N}}$ with $|\tau| < |\sigma|$, we have already defined $\Psi(\tau)$, and $l(\tau, e)$ for each $e \in \mathbb{N}$. Then, we define $\Psi(\sigma)$ and $l(\sigma, e)$ for each e as follows:

$$\Psi(\sigma) = \begin{cases} \mu e < |\sigma|\ [\Phi_e(\sigma) \upharpoonright (l(\sigma^-, e) + 1) \in T_P] & \text{if such } e \text{ exists,} \\ |\sigma| & \text{otherwise} \end{cases}$$

$$l(\sigma, e) = \begin{cases} l(\sigma^-, e) + 1 & \text{if } e = \Psi(\sigma), \\ l(\sigma^-, e) & \text{otherwise.} \end{cases}$$

By our assumption $P \leq^\omega_1 Q$, $\liminf_n \Psi(f \upharpoonright n)$ exists for all $f \in P$. Thus, the desired condition $\Phi_{\liminf_n \Psi(f \upharpoonright n)}(f) \in Q$ holds. □

Remark. Recall that a subset of $2^\mathbb{N}$ is Π^0_1 if and only if it is the set of all infinite paths through a computable subtree of $2^{<\mathbb{N}}$. Thus, in our model of inductive inference, each learner tries to learn a program for an infinite branch of T from a given infinite branch of another tree T^*. Another model of *branch learning* has been studied by Kummer-Ott [47], and Ott-Stephan [56] in which each learner tries to learn a program for an infinite *computable* branch of T from the global information about T. They pointed out that the concept of branch learning is equivalent to learning winning strategies for closed computable Gale-Stewart games, since the class of Π^0_1 subsets of $2^\mathbb{N}$ correspond exactly to the class of winning strategies for such games (see also Cenzer-Remmel [22]). Case-Ott-Sharma-Stephan [18] explains the concept of branch learning by using a temperature controller trying to keep the temperature in a room between 18°C and 22°C. In their model, each learner tries to learn a program for an infinite computable branch of T from the global information about T *with an additional information about one infinite branch of* T, i.e., the leaner may watch a human *master*. A k-wise variation for branch learning called *weak k-search problem* has been studied by Kaufmann-Kummer [46].

3. Degrees of Difficulty of Disjunctions

To separate the strength of superintuitionistic disjunctions, a rather exotic Π^0_1 set constructed by Jockusch and Soare is useful. We say that a set $A \subseteq \mathbb{N}^\mathbb{N}$ is *an antichain* if it is an antichain with respect to the Turing reducibility $\leq_T$. In other words, f is Turing incomparable with g, for any two distinct elements $f, g \in A$. A nonempty closed set $A \subseteq \mathbb{N}^\mathbb{N}$ is *perfect* if it has no isolated point.

Theorem 51 (Jockusch-Soare [45])**.** *There exists a perfect* Π^0_1 *antichain in* $2^\mathbb{N}$.

A stronger condition is sometimes required. For a set $P \subseteq \mathbb{N}^\mathbb{N}$ and an element $g \in \mathbb{N}^\mathbb{N}$, let $P^{\leq_T g}$ denote the set of all element of P which are Turing reducible to g. Then, a set $A \subseteq \mathbb{N}^\mathbb{N}$ is antichain if and only if $A^{\leq_T g} = \{g\}$ for every $g \in A$. A set $P \subseteq \mathbb{N}^\mathbb{N}$ is *independent* if $P^{\leq_T \bigoplus D} = D$ for every finite subset $D \subset P$.

Theorem 52 (see Binns-Simpson [8])**.** *There exists a perfect independent* Π^0_1 *subset of* $2^\mathbb{N}$.

On the other hand, in the later section, we will see that our hierarchy of disjunctions collapses for homogeneous sets, which may be regarded as an opposite notion to antichains and independent sets.

3.1. The Disjunction $\oplus$ versus the Disjunction $\cup$.

Recall that a set $P \subseteq \mathbb{N}^\mathbb{N}$ is *special* if it is nonempty and it contains no computable points.

Lemma 53. *Let P_0, P_1 be Π^0_1 subsets of $2^\mathbb{N}$, and let Q be a special Π^0_1 subset of $2^\mathbb{N}$. Assume that there exist $f \in P_0$ and $g \in P_1$ with $Q^{\leq_T f \oplus g} = Q^{\leq_T f} \cup Q^{\leq_T g}$ such that $Q^{\leq_T f}$ and $Q^{\leq_T g}$ are separated by open sets. Then $Q \not\leq^1_1 (P_0 \otimes 2^\mathbb{N}) \cup (2^\mathbb{N} \otimes P_1)$.*

Proof. Suppose that $Q \leq^1_1 (P_0 \otimes 2^\mathbb{N}) \cup (2^\mathbb{N} \otimes P_1)$ via a computable functional Φ. Then $f \oplus g \in (P_0 \otimes 2^\mathbb{N}) \cup (2^\mathbb{N} \otimes P_1)$. By our choice of f and g, $\Phi(f \oplus g)$ must belong to $Q^{\leq_T f \oplus g} = Q^{\leq_T f} \cup Q^{\leq_T g}$. By our assumption, $Q^{\leq_T f}$ and $Q^{\leq_T g}$ are separated by two disjoint open sets $U, V \subseteq 2^\mathbb{N}$. That is, $Q^{\leq_T f} \subseteq U$, $Q^{\leq_T g} \subseteq V$, and $U \cap V = \emptyset$. Therefore, either $\Phi(f \oplus g) \in Q \cap U$ or $\Phi(f \oplus g) \in Q \cap V$ holds. In any case, there exists an open neighborhood $[\sigma] \ni \Phi(f \oplus g)$ such that $[\sigma] \subseteq U$ or $[\sigma] \subseteq V$. Without loss of generality, we can assume $[\sigma] \subseteq U$. We pick initial segments $\tau_0 \subset f$ and $\tau_1 \subset g$ with $\Phi(\tau_0 \oplus \tau_1) \supseteq \sigma$. Then $(\tau_0{}^\frown 0^\mathbb{N}) \oplus g \in (P_0 \otimes 2^\mathbb{N}) \cup (2^\mathbb{N} \otimes P_1)$, and it is Turing equivalent to g. However this is impossible because $\Phi(\tau_0{}^\frown 0^\mathbb{N} \oplus g) \in [\sigma]$, and $[\sigma] \cap Q^{\leq_T g} \subseteq U \cap Q^{\leq_T g} = \emptyset$. □

Corollary 54. (1) *There are Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \cup Q <^1_1 P \oplus Q$.*
(2) *There are Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \equiv^{<\omega}_{tt,1} Q$ and $P <^1_1 Q$.*

Proof. (1) Let R be a perfect independent Π^0_1 subset of $2^\mathbb{N}$. Set $P = 2^\mathbb{N} \otimes R$ and $Q = R \otimes 2^\mathbb{N}$. Note that $P \oplus Q \equiv^1_1 R$. Pick $f, g \in R$ such that $f \neq g$. Then $R^{\leq_T f} = \{f\}$, $R^{\leq_T g} = \{g\}$, and $R^{\leq_T f \oplus g} = R^{\leq_T f} \sqcup R^{\leq_T g} = \{f, g\}$. Since R is Hausdorff, two points f and g are separated by open sets. Thus, $P \oplus Q \equiv^1_1 R \not\leq^1_1 P \cup Q$ by Lemma 53. (2) $P \oplus Q \equiv^{<\omega}_{tt,1} P \cup Q <^1_1 P \oplus Q$. □

3.2. The Disjunction $\cup$ versus the Disjunction $\triangledown$.

Lemma 55. *Let P_0, P_1 be Π^0_1 subsets of $2^\mathbb{N}$, and let Q be a special Π^0_1 subset of $2^\mathbb{N}$. Assume that there exist $f \in P_0$ and $g \in P_1$ such that any $h \in Q^{\leq_T f}$ and $Q^{\leq_T g}$ are separated by open sets. Then $Q \not\leq^1_1 P_0{}^\frown P_1$.*

Proof. Suppose that $Q \le^1_1 P_0{}^\frown P_1$ via a computable functional Φ. By our choice of $f \in P_0 \subseteq P_0{}^\frown P_1$, there must exist an open set $U \subseteq 2^{\mathbb{N}}$ such that $\Phi(f) \in Q \cap U$ and $Q^{\le_T g} \cap U = \emptyset$. Since U is open there exists a clopen neighborhood $[\sigma] \ni \Phi(f)$ such that $[\sigma] \cap Q \subseteq U$. We pick an initial segment $\tau \subset f$ with $\Phi(\tau) \supseteq \sigma$. Since $f \in P_0$ holds, we have that $\tau \in T_{P_0}$, and we pick $\rho \in L_{P_0}$ extending τ. Then $\rho^\frown g \in P_0{}^\frown P_1$, and $\rho^\frown g$ is Turing equivalent to g. So, if $Q \le^1_1 P_0{}^\frown P_1$ via Φ, then $\Phi(\rho^\frown g)$ must belong to $Q^{\le_T g}$. However this is impossible because $\Phi(\rho^\frown g) \in [\sigma]$, and $[\sigma] \cap Q^{\le_T g} \subseteq U \cap Q^{\le_T g} = \emptyset$. □

Corollary 56. *There are Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ such that $P^\frown Q <^1_1 P \cup Q <^1_1 P \oplus Q$.*

Proof. Assume that R be a perfect Π^0_1 antichain of $2^{\mathbb{N}}$. Set $P = 2^{\mathbb{N}} \otimes R$ and $Q = R \otimes 2^{\mathbb{N}}$. Pick $f, g \in R$ such that $f \neq g$. Then $R^{\le_T f} = \{f\}$ and $R^{\le_T g} = \{g\}$ since R is antichain. Therefore, $(P \cup Q)^{\le_T X} \subseteq (\{X\} \otimes 2^{\mathbb{N}}) \cup (2^{\mathbb{N}} \otimes \{X\})$ for each $X \in \{f, g\}$. For $h = h_0 \oplus h_1 \in (P \cup Q)^{\le_T f}$, we have $h_0 \neq g$ and $h_1 \neq g$. Thus, $h \notin (2^{\mathbb{N}} \otimes \{g\}) \cup (\{g\} \otimes 2^{\mathbb{N}})$, and note that $(2^{\mathbb{N}} \otimes \{g\}) \cup (\{g\} \otimes 2^{\mathbb{N}})$ is closed. Then, there is an open neighborhood $U \subseteq 2^{\mathbb{N}}$ such that $h \in U$ and $U \cap (P \cup Q)^{\le_T g} = \emptyset$, since $P \cup Q$ is regular, and $(P \cup Q)^{\le_T g} \subseteq (2^{\mathbb{N}} \otimes \{g\}) \cup (\{g\} \otimes 2^{\mathbb{N}})$. Namely, any $h \in (P \cup Q)^{\le_T f}$ and $(P \cup Q)^{\le_T g}$ are separated by some open set. Consequently, by Lemma 55, we have $P \cup Q \not\le^1_1 P^\frown Q$. □

Let $P^{\to}Q$ and $P^{\sqcap}Q$ denote $[\{\sigma^\frown \sharp^\frown \tau : \sigma \in T_P \ \& \ \tau \in T_Q\}]$ and $[\{\sigma^\frown \tau : \sigma \in T_P \ \& \ \tau \in T_Q\}]$, respectively. Recall from Proposition 21 that $P^\frown Q \equiv^1_1 P^{\to}Q$. However, there is a $(1,1)$-difference between $P^\frown Q$ and $P^{\sqcap}Q$.

Proposition 57. *There are Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ such that $P^{\sqcap}Q <^1_1 P^\frown Q$.*

Proof. It is easy to see that $P^{\sqcap}Q \le^1_1 P^{\to}Q$ for any $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$. Let $R \subseteq 2^{\mathbb{N}}$ be a Π^0_1 antichain. Then we divide R into four parts, P_0, P_1, P_2, and P_3. Put $P = P_3$, and $Q = (\langle 0,1\rangle^\frown P_2{}^\frown P_0) \cup (\langle 1\rangle^\frown P_2{}^\frown P_1)$. Without loss of generality, we may assume that $\langle 0\rangle \in T_P$. Suppose that $P^{\to}Q \le^1_1 P^{\sqcap}Q$ via a computable function Φ. Choose $g \in P_2$. Then we have $\langle 0,1\rangle^\frown g \in P^{\sqcap}Q$. Therefore, $\Phi(\langle 0,1\rangle^\frown g) \in P^{\to}Q$ must contain $\sharp$, since $P = P_3$ has no element computable in $g \in P_2$. Thus, there is $n \in \mathbb{N}$ such that $\Phi(\langle 0,1\rangle^\frown (g \upharpoonright n))$ contains $\langle \sharp, i\rangle$ as a substring for some $i < 2$. Fix such i. Then, $\Phi(\langle 0,1\rangle^\frown (g \upharpoonright n)) \in P^{\to}(Q \cap [\langle i\rangle])$. We extend $g \upharpoonright n$ to some leaf ρ of P_2. Choose $h_k \in P_k$ for each $k < 2$. Then, $\langle 0,1\rangle^\frown \rho^\frown h_0 \in Q \subseteq P^{\sqcap}Q$, and $\langle 0,1\rangle^\frown \rho^\frown h_1 \in \langle 0\rangle^\frown Q \subseteq P^{\sqcap}Q$. Thus, $\Phi(\langle 0,1\rangle^\frown \rho^\frown h_k)$ must belongs to $P^{\to}(Q \cap [\langle i\rangle])$, for each $k < 2$. However $P^{\to}(Q \cap [\langle i\rangle])$ has no element computable in $\langle 0,1\rangle^\frown \rho^\frown h_{1-i}$. A contradiction. □

3.3. **The Disjunction $\triangledown$ versus the Disjunction $\triangledown_\omega$.** Let Ψ be a learner (i.e., a total computable function $\Psi : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$). A point $\alpha \in \mathbb{N}^{\mathbb{N}}$ is said to be *an m-changing point of* Ψ if $\#\{s \in \mathbb{N} : \Psi(\alpha \upharpoonright s+1) \neq \Psi(\alpha \upharpoonright s)\} \geq m$. Then, the set of all m-changing point of Ψ is denoted by $\mathrm{Change}_\Psi(\geq m)$. A point $\alpha \in \mathbb{N}^{\mathbb{N}}$ is *anti-Popperian* with respect to Ψ if $\lim_n \Psi(\alpha \upharpoonright n)$ converges, but $\Phi_{\lim_n \Psi(\alpha \upharpoonright n)}(\alpha)$ is partial. The set of all anti-Popperian points of Ψ is denoted by AP_Ψ.

Remark **(Trichotomy).** Let Γ be a $(1,\omega)$-computable function identified by a learner Ψ, and let P be any subset of Baire space $\mathbb{N}^{\mathbb{N}}$. Then $\mathbb{N}^{\mathbb{N}} \setminus \Gamma^{-1}(P)$ is divided into the following three parts: the set $\Gamma^{-1}(\mathbb{N}^{\mathbb{N}} \setminus P)$; the Σ^0_2 set AP_Ψ; and the Π^0_2 set $\bigcap_{m \in \mathbb{N}} \mathrm{Change}_\Psi(\geq m)$.

We say that P_0 and P_1 are *everywhere $(\omega, 1)$-incomparable* if $P_0 \cap [\sigma_0]$ is Muchnik incomparable with $P_1 \cap [\sigma_1]$ (that is, $P_i \cap [\sigma_i] \not\le^\omega_1 P_{1-i} \cap [\sigma_{1-i}]$ for each $i < 2$) whenever $[\sigma_i] \cap P_i \neq \emptyset$ for each $i < 2$.

Theorem 58. *Let P_0, P_1 be everywhere $(\omega, 1)$-incomparable Π^0_1 subsets of $2^{\mathbb{N}}$, and ρ be any binary string. For any $(1,\omega)$-computable function Γ identified by a learner Ψ, the closure of* $\mathrm{Change}_\Psi(\geq m) \cup \Gamma^{-1}(\mathbb{N}^{\mathbb{N}} \setminus P_0 \oplus P_1) \cup \mathrm{AP}_\Psi$ *includes $\rho^\frown(P_0 \triangledown_n P_1)^\heartsuit$ with respect to the relative topology on $\rho^\frown(P_0 \triangledown_{m+n} P_1)^\heartsuit$ (as a subspace of Baire space $\mathbb{N}^{\mathbb{N}}$).*

Proof. Fix a string $\rho^\frown \tau_0$ which is extendible in the heart of $\rho^\frown(P_0 \triangledown_n P_1)$. Then, $\mathtt{pr}_i(\tau_0)$ must be extendible in P_i. Fix $f_i \in P_i \cap [\mathtt{pr}_i(\tau_0)]$ witnessing $P_{1-i} \not\le^\omega_1 P_i$ for each $i < 2$, i.e., P_{1-i} contains no f_i-computable element. Such f_i exists, by everywhere $(\omega, 1)$-incomparability. Assume that $f_i = \mathtt{pr}_i(\tau_0)^\frown f_i^*$ for each

$i < 2$ and that the last declaration along τ_0 is j_0, i.e., $\tau_0 = \tau_0^{-\frown}(j_0, k)$ for some $k < 2$. Then we can proceed the following actions.

- Extend τ_0 to $g_0 = \tau_0{}^\frown\mathtt{write}(j_0, f^*_{j_0}) \in \rho^\frown(P_0 \triangledown_n P_1)$.
- Wait for the least $s_0 > |\tau_0|$ such that $\Phi_{\Psi(g_0 \upharpoonright s_0)}(g_0 \upharpoonright s_0; 0) = j_0$.
- Extend $g_0 \upharpoonright s_0$ to $g_1 = (g_0 \upharpoonright s_0)^\frown\mathtt{write}(j_1, f^*_{j_1}) \in \rho^\frown(P_0 \triangledown_{n+1} P_1)$, where $j_1 = 1 - j_0$.
- Wait for the least $s_1 > s_0$ such that $\Phi_{\Psi(g_1 \upharpoonright s_1)}(g_1 \upharpoonright s_1; 0) = 1 - j_0$.

If both s_0 and s_1 are defined, then this action forces the learner Ψ to change his mind. In other words, $g_1 \in \mathrm{Change}_\Psi(\geq 1)$. Assume that s_l is undefined for some $l < 2$ Note that $g_l \equiv_T f_{j_l}$, since $\mathtt{pr}_{j_l}(g_l) = f_{j_l}$ and $\mathtt{pr}_{1-j_l}(g_l)$ is finite, for each $l < 2$. Therefore, $\Gamma(g_l) \notin (1 - j_l)^\frown P_{1-j_l}$ since P_{1-j_l} has no g_l-computable element. In this case, $g_l \in \Gamma^{-1}(\mathbb{N}^\mathbb{N} \setminus P_0 \oplus P_1)$. Hence, in $\rho^\frown(P_0 \triangledown_{n+1} P_1)^\heartsuit$, the closure of $\mathrm{Change}_\Psi(\geq 1) \cup \Gamma^{-1}(\mathbb{N}^\mathbb{N} \setminus P_0 \oplus P_1) \cup \mathrm{AP}_\Psi$ includes $\rho^\frown(P_0 \triangledown_n P_1)^\heartsuit$. By iterating this procedure, in $\rho^\frown(P_0 \triangledown_{m+n} P_1)^\heartsuit$, we can easily see that the closure of $\mathrm{Change}_\Psi(\geq m) \cup \Gamma^{-1}(\mathbb{N}^\mathbb{N} \setminus P_0 \oplus P_1) \cup \mathrm{AP}_\Psi$ includes $\rho^\frown(P_0 \triangledown_n P_1)^\heartsuit$. □

Corollary 59.

1. *There exists Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \triangledown_\omega Q <^1_{<\omega} P \triangledown Q$.*
2. *There exists Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \equiv^1_{\omega|<\omega} Q$ and $P <^1_{<\omega} Q$.*

Proof. (1) Let P be a perfect Π^0_1 antichain in $2^\mathbb{N}$ of Theorem 51. Fix a clopen set C such that $P_0 = P \cap C \neq \emptyset$, and $P_1 = P \setminus C \neq \emptyset$. Then every $f \in P_0$ and $g \in P_1$ are Turing incomparable. Therefore, P_0 and P_1 are everywhere $(\omega, 1)$-incomparable. Let ρ_n denote the n-th leaf of the tree T_{CPA} of a Medvedev complete Π^0_1 set $\mathsf{CPA} \subseteq 2^\mathbb{N}$. Fix a $(1, m)$-computable function Γ identified by a learner Ψ. By Theorem 58, $\rho_{m+1}{}^\frown(P_0 \triangledown_{m+1} P_1)$ intersects with $\mathrm{Change}_\Psi(\geq m+1) \cup \Gamma^{-1}(\omega^\omega \setminus P_0 \oplus P_1)$. Thus, $P_0 \oplus P_1 \not\leq^1_{<\omega} \overrightarrow{\bigoplus_n}(P_0 \triangledown_n P_1)$. Additionally, we easily have $P_0 \triangledown_\omega P_1 \leq^1_{<\omega} \overrightarrow{\bigoplus_n}(P_0 \triangledown_n P_1)$. (2) $P = \overrightarrow{\bigoplus_n}(P_0 \triangledown_n P_1)$ and $Q = P_0 \oplus P_1$ are Π^0_1. □

3.4. The Disjunction $\triangledown_\omega$ versus the Disjunction $\triangledown_\infty$.

Theorem 60. *Let P_0, P_1 be everywhere $(\omega, 1)$-incomparable Π^0_1 subsets of $2^\mathbb{N}$. For any $(1, \omega)$-computable function Γ, the complement of $\Gamma^{-1}(P_0 \oplus P_1)$ is dense in $(P_0 \triangledown_\infty P_1)^\heartsuit$ (as a subspace of Baire space $\mathbb{N}^\mathbb{N}$).*

Proof. Fix a learner Ψ which identifies the $(1, \omega)$-computable function Γ. Fix any clopen set $[\tau]$ intersecting with the heart of $(P_0 \triangledown_\infty P_1)$. Assume that $[\tau] \cap (P_0 \triangledown_\infty P_1)^\heartsuit$ contains no element of $\Gamma^{-1}(\mathbb{N}^\mathbb{N} \setminus P_0 \oplus P_1) \cup \mathrm{AP}_\Psi$. By Theorem 58, $\mathrm{Change}_\Psi(\geq n)$ is dense and open in the heart of $(P_0 \triangledown_\infty P_1) \cap [\tau] = \tau^\frown((P_0 \cap [\mathtt{pr}_0(\tau)]) \triangledown_\infty (P_1 \cap [\mathtt{pr}_1(\tau)]))$. As $[\tau] \cap (P_0 \triangledown_\infty P_1)^\heartsuit$ is Π^0_1, the intersection $\bigcap_{n \in \mathbb{N}} \mathrm{Change}_\Psi(\geq n)$ is dense in $[\tau] \cap (P_0 \triangledown_\infty P_1)^\heartsuit$, by Baire Category Theorem. Hence, $\mathbb{N}^\mathbb{N} \setminus \Gamma^{-1}(P_0 \oplus P_1)$ intersects with any nonempty clopen set $[\tau]$ with $[\tau] \cap (P_0 \triangledown_\infty P_1)^\heartsuit \neq \emptyset$. □

Corollary 61.

1. *There exist Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \equiv^{<\omega}_1 Q$ holds but $Q \not\leq^1_\omega P$ holds.*
2. *There exist Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$ such that $P \equiv^{<\omega}_\omega Q$ holds but $P <^1_\omega Q$ holds.*

Proof. Let P be a perfect Π^0_1 antichain in $2^\mathbb{N}$ of Theorem 51. Fix a clopen set C such that $P_0 = P \cap C \neq \emptyset$, and $P_1 = P \setminus C \neq \emptyset$. Then P_0 and P_1 are everywhere $(\omega, 1)$-incomparable. Fix a $(1, \omega)$-computable function Γ identified by a learner Ψ. By Theorem 60, $\mathbb{N}^\mathbb{N} \setminus (P_0 \oplus P_1)$ is dense in $(P_0 \triangledown_\infty P_1)^\heartsuit$. For Π^0_1 sets $P_0, P_1 \subseteq 2^\mathbb{N}$, both $P_0 \triangledown_\infty P_1$ and $P_0 \oplus P_1$ are Π^0_1, and $P_0 \triangledown_\omega P_1 \leq^1_1 P_0 \oplus P_1$. □

3.5. Infinite-Tape Models.

This subsection is concerned with degrees of difficulty of *infinitary* disjunctive notions between *the intuitionistic infinitary disjunction* and *the classical infinitary disjunction*.

Definition 62. Fix a collection $\{P_i\}_{i \in I}$ of subsets of Baire space $\mathbb{N}^\mathbb{N}$.



FIGURE 1. The two-tape (bounded-errors) model of disjunctions for independent Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$.

(1) $[\![\bigvee_{i\in I} P_i]\!]_{\mathsf{Int}} = \{f \in (I\times\mathbb{N})^{\mathbb{N}} : ((\exists i \in I)\ \mathtt{pr}_i(f) \in P_i)\ \&\ \mathtt{mc}(f) = 0\}$.
(2) $[\![\bigvee_{i\in I} P_i]\!]_{\mathsf{LCM}} = \{f \in (I\times\mathbb{N})^{\mathbb{N}} : ((\exists i \in I)\ \mathtt{pr}_i(f) \in P_i)\ \&\ \mathtt{mc}(f) < \infty\}$.
(3) $[\![\bigvee_{i\in I} P_i]\!]_{\mathsf{CL}} = \{f \in (I\times\mathbb{N})^{\mathbb{N}} : (\exists i \in I)\ \mathtt{pr}_i(f) \in P_i\}$.

Proposition 63. *Let $\{P_n\}_{n\in\mathbb{N}}$ be an infinite collection of subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) $[\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{Int}} \equiv^1_1 \bigoplus_{n\in\mathbb{N}} P_n$*, where* $\bigoplus_{n\in\mathbb{N}} P_n = \{\langle n\rangle{}^\frown f : f \in P_n\}$*.*
(2) $[\![\bigvee_{i,n} P_{i,n}]\!]_{\mathsf{LCM}} \equiv^1_1 [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$*, where $P_{i,n} = P_i$ for each $i < 2$ and $n \in \mathbb{N}$.*

Proof. (1) $[\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{Int}} \geq^1_1 \bigoplus_{n\in\mathbb{N}} P_n$ is witnessed by $f \mapsto (f(0))_0{}^\frown\mathtt{pr}_{(f(0))_0}(f)$, and $[\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{Int}} \leq^1_1 \bigoplus_{n\in\mathbb{N}} P_n$ is witnessed by $f \mapsto \mathtt{write}(f(0), f^{\leftharpoonup 1})$, where recall that $\mathtt{write}(f(0), f^{\leftharpoonup 1}) = (f(0))^{\mathbb{N}} \oplus (\lambda n.f(n+1))$ denotes the infinite sequence $\langle (f(0), f(1)), (f(0), f(2)), \dots, (f(0), f(n)), (f(0), f(n+1)), \dots\rangle$ which indicates the instruction to writing the infinite word $f^{\leftharpoonup 1}$ on the $f(0)$-th tape.

(2) We first construct a computable function Ξ witnessing $[\![\bigvee_{i,n} P_{i,n}]\!]_{\mathsf{LCM}} \geq^1_1 [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$. For $((i,n),v) \in (2\times\mathbb{N})\times\mathbb{N}$, we first set $\Xi(\langle((i,n),v)\rangle) = \langle((i,n),v)\rangle$. For each string $\sigma \in ((2\times\mathbb{N})\times\mathbb{N})^{<\mathbb{N}}$ inductively assume that $\Xi(\sigma^-)$ has been already defined. Assume $\sigma = \sigma^{--\frown}\langle((i,n),v),((j,m),w)\rangle$. If $(i,n) = (j,m)$, then we set $\Xi(\sigma) = \Xi(\sigma^-)^\frown\langle w\rangle$. Otherwise, we set $\Xi(\sigma) = \Xi(\sigma^-)^\frown\langle\sharp, j, w\rangle$. For any $f \in [\![\bigvee_{i,n} P_{i,n}]\!]_{\mathsf{LCM}}$, the backtrack symbol $\sharp$ occurs in $\Xi(f)$ finitely often, since $\mathtt{mc}(f) < \infty$. Therefore, $\mathtt{tail}(\Xi(f))$ converges, and $\mathtt{tail}(\Xi(f))^{\leftharpoonup 1} = \mathtt{pr}_{i,m}(f) \in P_i$ for some $i < 2$ and $m \in \mathbb{N}$. Thus, $\Xi(f) \in [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$.

We next construct a computable function Ξ^* witnessing $[\![\bigvee_{i,n} P_{i,n}]\!]_{\mathsf{LCM}} \leq^1_1 [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$. Set $\Xi^*(\langle\rangle) = \langle\rangle$. For $\sigma = \sigma^{--\frown}\langle v, w\rangle \in (\mathbb{N}\cup\{\sharp\})^{<\mathbb{N}}$, inductively assume that $\Xi^*(\sigma^-)$ has been already defined. To define $\Xi^*(\sigma)$, recall the definition $\mathtt{count}(\sigma) = \#\{n < |\sigma| : \sigma(n) = \sharp\}$. In the case that $v \neq \sharp$ and $w \neq \sharp$, we let $\Xi^*(\sigma)$ be the concatenation of $\Xi^*(\sigma^-)$ and $\langle((\mathtt{tail}(\sigma;0), \mathtt{count}(\sigma)), w)\rangle$. Otherwise, we set $\Xi^*(\sigma) = \Xi^*(\sigma^-)$. For any $f \in [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$, we have $\mathtt{mc}(\Xi^*(f)) < \infty$, since $\mathtt{count}(f) = \#\{k \in \mathbb{N} : f(k) = \sharp\}$ is finite. Therefore, we have $\mathtt{pr}_{(\mathtt{tail}(f;0),\mathtt{count}(f))}(\Xi^*(f)) = \mathtt{tail}(f)^{\leftharpoonup 1} \in P_{\mathtt{tail}(f;0)}$. Thus, $\Xi^*(f) \in [\![\bigvee_{i,n} P_{i,n}]\!]_{\mathsf{LCM}}$. □

We again use the consistent modifications of infinitary models, $\left[\bigtriangledown_\omega\right]_{n\in\mathbb{N}} P_n = [\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{LCM}} \cap \mathrm{Con}(T_{P_n})_{n\in\mathbb{N}}$, and $[\bigtriangledown_\infty]_{n\in\mathbb{N}} P_n = [\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{CL}} \cap \mathrm{Con}(T_{P_n})_{n\in\mathbb{N}}$.

Proposition 64. *Let $\{P_n\}_{n\in\mathbb{N}}$ be a computable collection of Π^0_1 subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) $[\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{LCM}} \equiv^1_1 \left[\bigtriangledown_\omega\right]_{n\in\mathbb{N}} P_n$*.*
(2) $[\![\bigvee_{n\in\mathbb{N}} P_n]\!]_{\mathsf{CL}} \equiv^1_1 [\bigtriangledown_\infty]_{n\in\mathbb{N}} P_n$*.*

Proof. As in the proof of Proposition 16. □

Recall that $\overrightarrow{\bigoplus}_n P_n$ is the recursive meet of computable collection $\{P_n\}_{n\in\mathbb{N}}$ of Π^0_1 subsets of $2^{\mathbb{N}}$ defined by $\overrightarrow{\bigoplus}_n P_n = \mathsf{CPA}^\frown\{P_n\}_{n\in\mathbb{N}} = \mathsf{CPA} \cup \bigcup_n \rho_n{}^\frown P_n$, where ρ_n is the length-lexicographically n-th leaf of the corresponding computable tree of the Π^0_1 set $\mathsf{CPA} \subseteq 2^{\mathbb{N}}$ consisting of complete consistent extensions of Peano Arithmetic.

Proposition 65. *For any computable sequence $\{P_n\}_{n\in\mathbb{N}}$ of nonempty Π^0_1 subsets of $2^{\mathbb{N}}$, $\overrightarrow{\bigoplus}_{n\in\mathbb{N}} P_n \equiv^1_{<\omega} \bigoplus_{n\in\mathbb{N}} P_n$.*

Proof. The condition $\overrightarrow{\bigoplus}_{n\in\mathbb{N}} P_n \leq^1_1 \bigoplus_{n\in\mathbb{N}} P_n$ is witnessed by a computable function $n^\frown g \mapsto \rho_n{}^\frown g$. We will construct a learner witnessing $\overrightarrow{\bigoplus}_{n\in\mathbb{N}} P_n \geq^1_{<\omega} \bigoplus_{n\in\mathbb{N}} P_n$. Fix a computable function $\Phi_e : \mathsf{CPA} \to 0^\frown P_0$. Such Φ_e exists, since every nonempty Π^0_1 subset of $2^{\mathbb{N}}$ is $(1,1)$-reducible to CPA. We also fix a partial computable function $\Phi_{i(n)} : \rho_n{}^\frown g \mapsto n^\frown g$, for each $n \in \mathbb{N}$. For $\sigma \in 2^{<\mathbb{N}}$, if $\sigma \in T_{\mathsf{CPA}}$ then set $\Psi(\sigma) = e$. If $\sigma \notin T_{\mathsf{CPA}}$, then $\rho_n \subseteq \sigma$ for some n. For such n, we set $\Psi(\sigma) = i(n)$. The function Γ identified by the learner Ψ is clearly $(1,2)$-computable, and $\Gamma(g) \in \bigoplus_{n\in\mathbb{N}} P_n$ for any $g \in \overrightarrow{\bigoplus}_{n\in\mathbb{N}} P_n$. ☐

When comparing these infinitary disjunctions, it quickly becomes clear that the bounded reducibility notions $\leq^1_{<\omega}$, $\leq^1_{\omega|<\omega}$, and $\leq^{<\omega}_1$ are useless. Moreover, it is easy to see the following.

Proposition 66. *Let $\{P_n\}_{n\in\mathbb{N}}$ be an infinite collection of subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) $\bigoplus_n P_n \equiv^1_\omega \left[\bigtriangledown_\omega\right]_n P_n$.
(2) $\bigoplus_n P_n \equiv^\omega_1 [\bigtriangledown_\infty]_n P_n$.

Proof. The computable function $n^\frown g \mapsto \mathtt{write}(n,g) = n^{\mathbb{N}} \oplus g$ witnesses $\bigoplus_n P_n \geq^1_1 \left[\bigtriangledown_\omega\right]_n P_n$ and $\bigoplus_n P_n \geq^1_1 [\bigtriangledown_\infty]_n P_n$. Fix a partial computable function $\Phi_{i(n)} : g \mapsto n^\frown \mathtt{pr}_n(g)$, for each $n \in \mathbb{N}$. Clearly, for any $g \in [\bigtriangledown_\infty]_n P_n$, there is $n \in \mathbb{N}$ such that $\Phi_{i(n)} \in \bigoplus_n P_n$. We construct a learner Ψ witnessing $\bigoplus_n P_n \leq^1_\omega \left[\bigtriangledown_\omega\right]_n P_n$. For $\sigma = \sigma^{-\frown}\langle(n,w)\rangle \in (\mathbb{N}\times\mathbb{N})^{<\mathbb{N}}$, set $\Psi(\sigma) = i(n)$. For any $g \in \left[\bigtriangledown_\omega\right]_n P_n$, the learner Ψ changes his mind finitely often along g, and $\Phi_{\lim_s \Psi(g\upharpoonright s)}(g) \in \bigoplus_n P_n$. ☐

Therefore, the only comparison that is left is between the intuitionistic infinitary disjunction $\bigoplus$ and the classical infinitary disjunction $\bigtriangledown_\infty$ under the $(1,\omega)$-reducibility. To understand the difference between degrees of difficulty of $\bigoplus$ and $\bigtriangledown_\infty$, the notion of a learning process *by a team of learners* is useful. Note that the infinitary disjunctions $\bigoplus$, $\bigtriangledown_\omega$, and $\bigtriangledown_\infty$ are not subsets of $2^{\mathbb{N}}$ (indeed, not compact), even if all P_n, $n \in \mathbb{N}$, are Π^0_1 subsets of $2^{\mathbb{N}}$. Therefore, it may be rather suitable to compare two compact Π^0_1 sets $\overrightarrow{\bigoplus}_t P_t$ and $\overrightarrow{\bigoplus}_t (P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$, for Π^0_1 subsets $P_t \subseteq 2^{\mathbb{N}}$, $t \in \mathbb{N}$. When one think of $\bigoplus_n P_n$ (resp. $[\bigtriangledown_\infty]_n P_n$) as an intuitionistic (resp. classical) interpretation of arithmetical existential quantifier $(\exists n)P_n$, then $\bigoplus_t (P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$ can be interpreted as calculating a bound t such that $(\exists n \leq t)P_n$ is classically valid.

Proposition 67. *Let $\{P_n\}_{n\in\mathbb{N}}$ be a collection of Π^0_1 subsets of $2^{\mathbb{N}}$. Then, the set $\overrightarrow{\bigoplus}_t (P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$ is computably homeomorphic to a Π^0_1 subset of $2^{\mathbb{N}}$.*

Proof. It suffices to show that $\overrightarrow{\bigoplus}_t (P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$ is computably bounded, that is to say, there is a computable function $h : \mathbb{N} \to \mathbb{N}$ such that $g(n) < h(n)$ for any element $g \in \overrightarrow{\bigoplus}_t (P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$. Note that $P_0 \triangledown_\omega \dots \triangledown_\omega P_t \subseteq ((t+1)\times 2)^{\mathbb{N}}$, and $|\rho_{2^t}| \geq t$. Hence, we have $g(t) < 2^{t+2}$ for any $t \in \mathbb{N}$. ☐

3.6. The Disjunction $\bigoplus$ versus the Disjunction $\bigtriangledown_\infty$.

A sequence $\{x_i\}_{i\in\mathbb{N}}$ of elements of $\mathbb{N}^{\mathbb{N}}$ is *Turing independent* if x_i is not computable in $\bigoplus_{j\neq i} x_j$ for each $i \in \mathbb{N}$. A collection $\{P_i\}_{i\in I}$ of subsets of $\mathbb{N}^{\mathbb{N}}$ is *pairwise everywhere independent* if, for any collection $\{[\sigma_i]\}_{i\in I}$ of clopen sets with $P_i \cap [\sigma_i] \neq \emptyset$ for each $i \in I$, there is a choice $\{x_i\}_{i\in I} \in \prod_{i\in I}(P_i \cap [\sigma_i])$ such that P_i has no element computable in $\bigoplus_{j\in I\setminus\{i\}} x_j$ for each $i \in I$.

Theorem 68. *Let $\{P_i\}_{i<2^t}$ be a pairwise everywhere independent collection of Π^0_1 subsets of $2^{\mathbb{N}}$, and let ρ be any binary string. For any (t,ω)-computable function Γ, the complement of $\Gamma^{-1}(P_0 \oplus \dots \oplus P_{2^t-1})$ is* *dense in the heart of* $\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})$ *(as a subspace of Baire space* $\mathbb{N}^\mathbb{N}$*). Indeed, for any nonempty interval* I *in the heart of* $\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})$*, there is* $g\in\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})^\heartsuit\cap I\setminus\Gamma^{-1}(P_0\oplus\dots\oplus P_{2^t-1})$ *which is computable in some* $g^*\in\bigotimes_{k<2^t-1}P_k$*.*

Proof. Assume that the (t,ω)-computable function Γ is identified by a team $\{\Psi_i\}_{i<t}$ of learners. Fix a string $\rho^\frown\tau_0$ which is extendible in the heart of $\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})$. Then, $\mathtt{pr}_i(\tau_0)$ must be extendible in P_i for each $i<2^t$. Fix $\{f_i\}_{i<2^t}\in\prod_{i<2^t}(P_i\cap[\mathtt{pr}_i(\tau_0)])$ witnessing the independence of $\{P_i\}_{i<2^t}$, i.e., P_i contains no $\bigoplus_{j\neq i}f_j$-computable element. Assume that $f_i=\mathtt{pr}_i(\tau_0)^\frown f_i^*$ for each $i<2^t$ and that the last declaration along τ_0 is $j_0<2^t$, i.e., $\tau_0=\tau_0^{-\frown}(j_0,k)$ for some $k<2$. Fix a computable function δ mapping $j<2^t$ to a unique binary string $\delta(j)$ satisfying $j=\sum_{e=0}^{t-1}2^e\cdot\delta(j;e)$. Let E^e_k denote the set $\{j<2^t:\delta(j;e)=k\}$. Then we can proceed the following actions.

- Extend τ_0 to $g_0=\tau_0{}^\frown\mathtt{write}(j_0,f^*_{j_0})\in\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})$.
- Wait for the least $s_0>|\tau_0|$ such that $\Phi_{\Psi_e(g_0\upharpoonright s_0)}(g_0\upharpoonright s_0;0)\in E^e_{\delta(j_0;e)}$ for some $e<2^t$.
- If such s_0 exists, then enumerate all such $e<2^t$ into an auxiliary set Ch_0, and define $\delta(j_1)$ as follows:
$$\delta(j_1;e)=\begin{cases}\delta(j_0;e) & \text{if } e\notin\mathrm{Ch}_0,\\ 1-\delta(j_0;e) & \text{if } e\in\mathrm{Ch}_0.\end{cases}$$
- Extend $g_0\upharpoonright s_0$ to $g_1=(g_0\upharpoonright s_0)^\frown\mathtt{write}(j_1,f^*_{j_1})\in\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})$, where $j_1=\sum_{e=0}^{t-1}2^e\cdot\delta(j_1;e)$.

These actions force each learner Ψ_e with $e\in\mathrm{Ch}_0$ to change his mind whenever the learner Ψ_e want to have an element of $\bigoplus_{i<2^t}P_i$. Fix $u\in\mathbb{N}$. Assume that j_u, g_u, s_u, and Ch_u has been already defined, and the following induction hypothesis at stage u is satisfied.

- $\mathtt{pr}_e(g_u)\subseteq f_e$ for any $e<2^t$, hence, $g_u\in\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})^\heartsuit\cap[\rho^\frown\tau_0]$.
- $\{s_v\}_{v\leq u}$ is strict increasing, and $\mathrm{Ch}_u\neq\emptyset$.
- For each $e\in\mathrm{Ch}_u$, if $\Phi_{\Psi_e(g_u\upharpoonright s_u)}(g_u\upharpoonright s_u;0)$ converges to some value $k<2^t$, then $k\in E^e_{\delta(j_u;e)}$

It is easy to see that $u=0$ satisfies the induction hypothesis. At stage $u+1\in\mathbb{N}$, we proceeds the following actions.

- Define $\delta(j_{u+1})$ as follows:
$$\delta(j_{u+1};e)=\begin{cases}\delta(j_u;e) & \text{if } e\notin\mathrm{Ch}_u,\\ 1-\delta(j_u;e) & \text{if } e\in\mathrm{Ch}_u.\end{cases}$$
- Extend $g_u\upharpoonright s_u$ to $g_{u+1}=(g_u\upharpoonright s_u)^\frown\mathtt{write}(j_{u+1},f^{(u+1)}_{j_{u+1}})$, where $j_{u+1}=\sum_{e=0}^{t-1}2^e\cdot\delta(j_{u+1};e)$, and f^{u+1} satisfies $\mathtt{pr}_{j_{u+1}}(g_{u+1})=\mathtt{pr}_{j_{u+1}}(g_u\upharpoonright s_u)^\frown f^{(u+1)}_{j_{u+1}}=\rho^\frown f_{j_{u+1}}$.
- Wait for the least $s_{u+1}>s_u$ such that $\Phi_{\Psi_e(g_{u+1}\upharpoonright s_{u+1})}(g_{u+1}\upharpoonright s_{u+1};0)\in E^e_{\delta(j_{u+1};e)}$ for some $e<2^t$.
- If such s_{u+1} exists, then enumerate all such $e<2^t$ into Ch_{u+1},

By our action, it is easy to see that $u+1$ satisfies the induction hypothesis. As $\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})^\heartsuit$ is closed (with respect to the Baire topology) and $\{s_u\}_{u\in\mathbb{N}}$ is strictly increasing, the sequence $\{g_u\}_{u\in\mathbb{N}}$ converges to some $g\in\rho^\frown(P_0\triangledown_\infty\dots\triangledown_\infty P_{2^t-1})^\heartsuit$. Let $I(g)\subseteq 2^t$ be the set of all $e<2^t$ such that $\mathtt{pr}_e(g)$ is total.

Claim. $g\leq_T\bigoplus_{e\in I(g)}f_e$.

Note that $g=g[f_0,\dots,f_{2^t-1}]$ is effectively constructed uniformly in a given collection $\{f_k\}_{k<2^t}$. In other words, there is a (uniformly) computable function Θ mapping $\{f_k\}_{k<2^t}$ to $\Theta(\{f_k\}_{k<2^t})=g=g[f_0,\dots,f_{2^t-1}]$. Then, it is easy to see that the function Θ maps $\{f_e\}_{e\in I(g)}\cup\{\mathtt{pr}_e(g)^\frown 0^\mathbb{N}\}_{e\in 2^t\setminus I(g)}$ to g. Hence, $g\leq_T\bigoplus_{e\in I(g)}f_e\oplus\bigoplus_{e\in 2^t\setminus I(g)}\mathtt{pr}_e(g)^\frown 0^\mathbb{N}$. Therefore, $g\leq_T\bigoplus_{e\in I(g)}f_e$ as desired, since $\mathtt{pr}_e(g)^\frown 0^\mathbb{N}$ is computable for any $e\in 2^t\setminus I(g)$.

Let Γ_e denote the $(1,\omega)$-computable function identified by Ψ_e, that is, $\Gamma_e(\alpha) = \Phi_{\lim_n \Psi(\alpha\upharpoonright n)}(\alpha)$ for any $\alpha \in \mathbb{N}^{\mathbb{N}}$. We consider the following two cases.

Case 1 ($e \in \mathrm{Ch}_u$ for finitely many $u \in \mathbb{N}$). Fix u such that $e \notin \mathrm{Ch}_v$ for any $v > u$. For each $v > u$, $\Phi_{\Psi_e(g\upharpoonright s_u)}(g \upharpoonright s_u; 0)$ does not converges to an element of $E^e_{\delta(j_v;e)} = E^e_{\delta(j_u;e)}$. By our definition, for each $k \notin E^e_{\delta(j_u;e)}$, $\mathtt{pr}_k(g) \subset \rho^\frown f_k$ is finite. By previous claim, $g \leq_T \bigoplus_{e\neq k} f_e$. Thus, by independence, P_k has no g-computable element. If $\Phi_{\Psi_e(g\upharpoonright s_u)}(g \upharpoonright s_u; 0)\uparrow$ for any $u \in \mathbb{N}$, then $g \in \mathrm{AP}_{\Psi_e}$. If $\lim_n \Psi_e(g \upharpoonright n)$ does not converge, then $g \in \bigcap_{m\in\mathbb{N}} \mathrm{Change}_{\Psi_e}(\geq m)$. Otherwise, $\Phi_{\lim_n \Psi_e(g\upharpoonright n)}(g; 0)$ converges to some value $k \notin E^e_{\delta(j_u;e)}$. As $\Phi_{\lim_n \Psi_e(g\upharpoonright n)}(g)$ is g-computable, we see $\Phi_{\lim_n \Psi_e(g\upharpoonright n)}(g) \notin k^\frown P_k$. Consequently, $g \in \mathbb{N}^{\mathbb{N}} \setminus \Gamma_e^{-1}(\bigoplus_{k<2^t} P_k)$.

Case 2 ($e \in \mathrm{Ch}_u$ for infinitely many $u \in \mathbb{N}$). We enumerate an infinite increasing sequence $\{u[n]\}_{n\in\mathbb{N}}$, where $u[n]$ is the n-th element such that $e \in \mathrm{Ch}_u$. As $e \in \mathrm{Ch}_{u[n]}$, we have $\Phi_{\Psi_e(g\upharpoonright u[n])}(g \upharpoonright u[n]; 0) \in E^e_{\delta(j_{u[n]};e)}$. By our action, $\delta(j_{u[n+1]}; e) = \delta(j_{u[n]+1}; e) \neq \delta(j_{u[n]}; e)$. This implies $E^e_{\delta(j_{u[n+1]};e)} \cap E^e_{\delta(j_{u[n]};e)} = \emptyset$. However, we must have $\Phi_{\Psi_e(g\upharpoonright u[n+1])}(g \upharpoonright u[n + 1]; 0) \in E^e_{\delta(j_{u[n+1]};e)}$, since $e \in \mathrm{Ch}_{u[n+1]}$. This forces the learner Ψ_e to change his mind. By iterating this procedure, we eventually obtain $g \in \bigcap_{m\in\mathbb{N}} \mathrm{Change}_{\Psi_e}(\geq m)$.

Consequently, $g \in \bigcap_{e\in\mathbb{N}}(\mathbb{N}^{\mathbb{N}} \setminus \Gamma_e^{-1}(\bigoplus_{k<2^t} P_k))$. Thus, $g \in \mathbb{N}^{\mathbb{N}} \setminus \Gamma_e^{-1}(\bigoplus_{k<2^t} P_k)$. For any τ_0 such that $\rho^\frown\tau_0$ which is extendible in the heart of $\rho^\frown(P_0 \triangledown_\infty \dots \triangledown_\infty P_{2^t-1})$, we can construct such g extending τ_0. Therefore, $\mathbb{N}^{\mathbb{N}} \setminus \Gamma_e^{-1}(\bigoplus_{k<2^t} P_k)$ intersects any nonempty interval in $\rho^\frown(P_0 \triangledown_\infty \dots \triangledown_\infty P_{2^t-1})^\heartsuit$. In other words, $\mathbb{N}^{\mathbb{N}} \setminus \Gamma_e^{-1}(P_0 \oplus \dots \oplus P_{2^t-1})$ is dense in $\rho^\frown(P_0 \triangledown_\infty \dots \triangledown_\infty P_{2^t-1})^\heartsuit$ as desired. □

The following theorem by Jockusch-Soare [45, Theorem 4.1] is important.

Theorem 69 (Jockusch-Soare [45]). *There is a computable sequence* $\{\prod_n P^i_n\}_{i\in\mathbb{N}}$ *of nonempty homogeneous* Π^0_1 *subsets of* $2^{\mathbb{N}}$ *such that* $\{x_i\}_{i\in\mathbb{N}}$ *is Turing independent for any choice* $x_i \in \prod_n P^i_n$, $i \in \mathbb{N}$.

Clearly any such Π^0_1 set contains no element of a PA degree, a Turing degree of a complete consistent extension of Peano Arithmetic. Accordingly, every element of such a Π^0_1 set computes no element of a Medvedev complete Π^0_1 set CPA.

Corollary 70. *There are* Π^0_1 *sets* $P_n \subseteq 2^{\mathbb{N}}$, $n \in \mathbb{N}$, *such that* $\overrightarrow{\bigoplus}_t(P_0 \triangledown_\infty \dots \triangledown_\infty P_t) <^{<\omega}_{\omega} \overrightarrow{\bigoplus}_t P_t$.

Proof. Fix the computable sequence $\{P_i\}_{i\in\mathbb{N}}$ of Theorem 69. Then $\{P_i\}_{i\in\mathbb{N}}$ is pairwise everywhere independent. Assume that $\overrightarrow{\bigoplus}_t P_t \leq^{<\omega}_{\omega} \overrightarrow{\bigoplus}_t(P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$ via a (t,ω)-computable function Γ. Let ρ_n denote the n-th leaf of the tree T_{CPA} of a Medvedev complete Π^0_1 subset of $2^{\mathbb{N}}$. By Theorem 68, $\Gamma^{-1}(\bigcup_{k<2^t} \rho_k{}^\frown P_k)$ is dense in the heart of $\rho_{2^t}{}^\frown(P_0 \triangledown_\infty \dots \triangledown_\infty P_{2^t-1})$. In particular, there is an infinite path $g \in \rho_{2^t}{}^\frown(P_0 \triangledown_\infty \dots \triangledown_\infty P_{2^t-1})$ such that $\Gamma(g) \notin \bigcup_{k<2^t} \rho_k{}^\frown P_k$ which is computable in some $g^* \leq_T \bigotimes_{k<2^t} P_k$. By our choice of $\{P_i\}_{i\in\mathbb{N}}$, $\Gamma(g)$ computes no element of $\bigcup_{k\geq 2^t} P_k \cup \mathsf{CPA}$. Thus, $\Gamma(g) \notin \overrightarrow{\bigoplus}_t P_t$. □

Corollary 71.

(1) *There exists a computable sequence* $\{P_n\}_{n\in\mathbb{N}}$ *of* Π^0_1 *subsets of Cantor space* $2^{\mathbb{N}}$, *such that the condition* $[\bigtriangledown_\infty]_n P_n <^{<\omega}_{\omega} \bigoplus_n P_n$ *is satisfied.*
(2) *There exist* Π^0_1 *sets* $P, Q \subseteq 2^{\mathbb{N}}$ *such that* $P \equiv^{\omega}_1 Q$ *holds but* $P <^{<\omega}_{\omega} Q$ *holds.*

Proof. (1) By Corollary 70. the condition $[\bigtriangledown_\infty]_n P_n \leq^1_1 \overrightarrow{\bigoplus}_t(P_0 \triangledown_\infty \dots \triangledown_\infty P_t) <^{<\omega}_{\omega} \overrightarrow{\bigoplus}_t P_t \leq^1_1 \bigoplus_n P_n$ is satisfied. (2) Put $P = [\bigtriangledown_\infty]_n P_n \leq^1_1 \overrightarrow{\bigoplus}_t(P_0 \triangledown_\infty \dots \triangledown_\infty P_t)$ and $Q = \overrightarrow{\bigoplus}_t P_t$. Then P and Q are $(1,1)$-equivalent to Π^0_1 subsets, by Proposition 67. By Theorem 70 and Proposition 66, $P <^{<\omega}_{\omega} Q \equiv^{\omega}_1 P$. □

4. Contiguous Degrees and Dynamic Infinitary Disjunctions

4.1. When the Hierarchy Collapses.

We have already observed the following hierarchy, for pairwise independent Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$.

$$P \oplus Q >^1_1 P \cup Q >^1_1 P \triangledown Q >^1_{<\omega} P \triangledown_\omega Q >^1_{\omega|<\omega} P \triangledown_\infty Q \equiv^{<\omega}_1 P \oplus Q.$$

Homogeneity is an opposite notion of antichain (and independence). Recall that $S \subseteq \mathbb{N}^{\mathbb{N}}$ is *homogeneous* if $S = \prod_n S_n$ for some $S_n \subseteq \mathbb{N}$, $n \in \mathbb{N}$. Every antichain is degree-non-isomorphic everywhere. On the other hand, every homogeneous set S is *degree-isomorphic everywhere*, that is to say, $S \cap C$ is degree-isomorphic to $S \cap D$ for any clopen sets $C, D \subseteq \mathbb{N}^{\mathbb{N}}$ with $S \cap C \neq \emptyset$ and with $S \cap D \neq \emptyset$. The next observation is that our hierarchy collapses for homogeneous Π^0_1 sets, modulo the $(1, <\omega)$-equivalence.

Theorem 72. *For every homogeneous Π^0_1 set $S \subseteq 2^{\mathbb{N}}$ and for any Π^0_1 set $Q \subseteq 2^{\mathbb{N}}$, if $S \leq^{<\omega}_1 Q$ then $S \leq^1_{<\omega} Q$.*

Proof. Let $S = \prod_x F_x$ for some Π^0_1 sets $F_x \subseteq \{0, 1\}$. Assume $S \leq^{<\omega}_1 Q$ via the bound b. That is, for every $g \in Q$ there exists an index $e < b$ such that $\Phi_e(g) \in S$. Let us begin defining a learner Ψ who changes his mind at most finitely often. Fix $g \in Q$. Ψ first sets $A_0 = \{e \in \mathbb{N} : e < b\}$. By our assumption, we have $\Phi_e(g) \in S$ for some $e \in A_0$. Then the learner Ψ challenges to predict the solution algorithm $e < b$ such that $\Phi_e(g) \in S$ by using an observation $g \in Q$. He begins the 1-st challenge. On *the* $(s + 1)$*-th challenge of* Ψ, inductively assume that, the learner have already defined a set $A_s \subseteq A_0$. Let v be a stage at which the $s + 1$-th challenge of Ψ on g begins. In this challenge, the learner Ψ uses the two following computable functionals Γ and Δ.

- For a given argument x, $\Gamma(x, s + 1)$ outputs the least $\langle e(x), t(x)\rangle$ such that $e(x) \in A_s$ and $\Phi_{e(x)}(g \upharpoonright t(x); x) \downarrow$ if such $\langle e(x), t(x)\rangle$ exists.
- If $\Gamma(x, s + 1) = \langle e(x), t(x)\rangle$, then $\Delta(g; x, s + 1) = \Phi_{e(x)}(g \upharpoonright t(x); x)$.

Set $\Delta_{s+1}(g; x) = \Delta(g; x, s + 1)$. Clearly, an index $d(s + 1)$ of Δ_{s+1} is calculated from $s + 1$. Then the learner $\Psi(g \upharpoonright v)$ outputs $d(s + 1)$ on the $(s + 1)$-th challenge. Hence $\Phi_{\Psi(g\upharpoonright v)}(g; x) = \Phi_{d(s+1)}(g; x) = \Phi_{e(x)}(g \upharpoonright t(x); x)$ for any x. He does not change his mind until the beginning stage v' of the next challenge, i.e., $\Phi_{\Psi(g\upharpoonright v'')}(g) = \Phi_{\Psi(g\upharpoonright k)}(g)$ for $k \leq v'' < v'$. *The next challenge* might begin when it turns out that Ψ's prediction on his $(s + 1)$-th challenge is incorrect, namely:

- $\Phi_{\Psi(g\upharpoonright v)}(g \upharpoonright u) \upharpoonright n \notin T_{S,u}$ for some $n < u$ at some stage $u > v$.

Here T_S is a corresponding computable tree of S. For each $x \in \mathbb{N}$, fix a decreasing approximation $\{F_{x,s}\}_{s\in\mathbb{N}}$ of a Π^0_1 set $F_x \subseteq \{0, 1\}$, uniformly in x. In this case, there exists $x < n$ such that the following condition holds.

$$\Phi_{\Psi(g\upharpoonright v)}(g \upharpoonright u; x) = \Delta_{s+1}(g; x) = \Phi_{e(x)}(g; x) \notin F_{x,s}.$$

For such a least x, the learner removes $e(x)$ from A_s, that is, let $A_{s+1} = A_s \setminus \{e(x)\}$. If $A_{s+1} \neq \emptyset$ then the learner Ψ begins *the* $(s + 2)$*-th challenge* at the current stage u. The construction of the learner Ψ is completed. An important point of this construction is that the learner never uses an index rejected on some challenge. This makes the prediction on $g \in Q$ of the learner Ψ converge.

Claim. Ψ changes his mind at most b times.

Whenever Ψ changes, A_s must decrease. However $\#A_0 = b$.

Claim. For every $g \in Q$ it holds that $\Phi_{\lim_s \Psi(g\upharpoonright s)}(g) \in S$.

For $g \in Q$, let $B^g \subseteq A_0$ be the set of all $e \in A_0$ such that $\Phi_e(g) \in S = \prod_x F_x$. By the definition of A_0, clearly B^g is not empty. Moreover, $B^g \subseteq \bigcap_s A_s$ holds, since e is removed from $\bigcap_s A_s$ only when $\Phi_e(g; x) \notin F_x$ for some x. Thus, $\Phi_{\Psi(g\upharpoonright v)}(g) : \mathbb{N} \to \mathbb{N}$ is total for every stage v. This means that, if $\Phi_{\Psi(g\upharpoonright v)}(g) \notin S$, then the learner Ψ will know his mistake at some stage u, i.e., $\Phi_{\Psi(g\upharpoonright v)}(g \upharpoonright u; x) \notin F_{x,u}$

for some $x < u$. Then some index is removed from $\bigcap_s A_s$. However, this occurs at most b times. Thus, $\Phi_{\lim_s \Psi(g\upharpoonright s)}(g) \in S$. □

Let $\alpha,\beta,\gamma \in \{1,<\omega,\omega\}$. We say that a $(\alpha,\beta|\gamma)$-degree $\mathbf{a}$ of a nonempty Π^0_1 subset of $2^{\mathbb{N}}$ is $(\alpha,\beta|\gamma)$-*complete* if $\mathbf{b} \leq \mathbf{a}$ for every $(\alpha,\beta|\gamma)$-degree $\mathbf{b}$ of a nonempty Π^0_1 subset of $2^{\mathbb{N}}$. If a Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ has an $(\alpha,\beta|\gamma)$-complete $(\alpha,\beta|\gamma)$-degree, then it is also called $(\alpha,\beta|\gamma)$-*complete*.

Corollary 73. *A Π^0_1 subset of $2^{\mathbb{N}}$ is $(1,<\omega)$-complete if and only if it is $(1,\omega|<\omega)$-complete if and only if it is $(<\omega,1)$-complete.*

Proof. Let DNR_2 denote the set of all two-valued diagonally noncomputable functions, where a function $f : \mathbb{N} \to 2$ is *diagonally noncomputable* if $f(e) \neq \Phi_e(e)$ for any index e. This set is clearly homogeneous, and Π^0_1. Moreover, it is $(1,1)$-complete (hence $(\alpha,\beta|\gamma)$-complete for any $\alpha,\beta,\gamma \in \{1,<\omega,\omega\}$). Therefore, we can apply Theorem 72 with $S = \mathsf{DNR}_2$. □

Corollary 74. *There are Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ such that $P \oplus Q \equiv^1_{<\omega} P\triangledown_\infty Q$. Indeed, if P is homogeneous and $Q \equiv^1_1 P$, then $P \oplus Q \equiv^1_{<\omega} P\triangledown_\infty Q$ is satisfied.*

Proof. Let P be any homogeneous Π^0_1 subset of $2^{\mathbb{N}}$. Then $P \oplus P$ is also homogeneous. By Proposition 35, there is a $(2,1)$-computable function from $P\triangledown_\infty P$ to $P \oplus P$, hence $P \oplus P \leq^{<\omega}_1 P\triangledown_\infty P$. Thus, by Theorem 72, $P \oplus P \leq^1_{<\omega} P\triangledown_\infty P$. If $Q \equiv^1_1 P$, then, by Proposition 14, $P \oplus Q \equiv^1_1 P \oplus P \leq^1_{<\omega} P\triangledown_\infty P \equiv^1_1 P\triangledown_\infty Q$. □

It is natural to ask whether our hierarchy of disjunctive notions for homogeneous Π^0_1 sets also collapses *modulo the* $(1,1)$-*equivalence*. The answer is *negative*. We say that a homogeneous set $\prod_n F_n$ is *computably bounded* if there is a computable function $l : \mathbb{N} \to \mathbb{N}$ such that $F_n \subseteq \{0,\dots,l(n)\}$ for any $n \in \mathbb{N}$. Clearly, every homogeneous subset of Cantor space $2^{\mathbb{N}}$ is computably bounded. Cenzer-Kihara-Weber-Wu [19] introduced the notion of immunity for closed sets. A closed subset P of Cantor space $2^{\mathbb{N}}$ is *immune* if T^{ext}_P has no infinite computable subset.

Theorem 75. *Let $P \subseteq 2^{\mathbb{N}}$ be a non-immune Π^0_1 set, and $S_0, S_1, \dots, S_m \subseteq \mathbb{N}^{\mathbb{N}}$ be special computably bounded homogeneous Π^0_1 sets. Then $\bigcup_{i\leq m} S_i \not\leq^1_1 P$.*

Proof. Let V_0 be an infinite c.e. subtree of T^{ext}_P. Assume that $\bigcup_{i\leq m}\prod_n F^i_n \leq^1_1 P$ via a computable functional Φ, where, for each $i < m$, $\{F^i_n\}_{n\in\omega}$ is a uniformly Π^0_1 sequence of subsets of $\{0,1,\dots l_i\}$. Let S^{ext}_i denotes the corresponding Π^0_1 tree of $\prod_n F^i_n$, and let $L_i = \{\rho : (\exists\tau \in S^{ext}_i)(\exists i)\ \rho = \tau^\frown\langle i\rangle \notin S^{ext}_i\}$, for each i. Note that L_i *differs* from the set of leaves of the corresponding *computable* tree of $\prod_n F^i_n$. We first consider the set $L^\Phi_i = \{\rho \in L_i : (\exists\sigma \in V_i)\ \Phi(\sigma) \supseteq \rho\}$, where V_i for $0 < i \leq m$ will be defined in the below construction. Note that L^Φ_0 is computably enumerable. There are three cases:

(1) L^Φ_0 is infinite;
(2) L^Φ_0 is finite, hence $\Phi([V_0])$ is a subset of $\prod_n F^0_n$;
(3) otherwise.

(Case 1): For any n, there exists $\rho \in L^\Phi_0$ of height $> n+1$, and $\rho(n) \in F^0_n$. From any computable enumeration of L^Φ_0 we can calculate a computable path of $\prod_n F^0_n$. This contradicts the specialness of $S_0 = \prod_n F^0_n$.

(Case 2): There exists a finite number k such that, for every string $\sigma \in V_0$ of height $> k$, $\Phi(\sigma)$ belongs to S^{ext}_i. This also contradicts the specialness of $S_0 = \prod_n F^0_n$.

(Case 3): There exists infinitely many strings $\sigma \in V_0$ such that $\Phi(\sigma)$ extends some string of L^Φ_0. Since L^Φ_0 is finite, by the pigeon hole principle, there exists $\rho_0 \in L^\Phi_0$ such that $\Phi(\sigma)$ extends ρ for infinitely many $\sigma \in V_0$. Fix such ρ_0, and let $V_1 = \{\sigma \in V_0 : \Phi(\sigma) \supseteq \rho\}$. Then the downward closure of V_1 is an infinite c.e. subtree of T^{ext}_P, and $\Phi([V_1]) \cap S_0 = \emptyset$.

By iterating this procedure, we win the either of the cases 1 or 2 for some $i \leq m$. The reason is that, if the case 3 occurs for j, then V_{j+1} is defined as an infinite c.e. subtree of T^{ext}_P such that $\Phi([V_1]) \cap$

$(\bigcup_{i\leq j} S_i) = \emptyset$. Since $\bigcup_{i\leq m}\prod_n F^i_n \leq^1_1 P \leq [V_m]$, i.e., $\Phi([V_m]) \subseteq \bigcup_{i\leq m} S_i$, the case 3 does not occur for m. $\square$

Corollary 76. *Let P, Q be any nonempty Π^0_1 subsets of $2^{\mathbb{N}}$, and S, T be computably bounded homogeneous Π^0_1 sets. Then $S \cup T \not\leq^1_1 P \triangledown Q$.*

Proof. Clearly $P \triangledown Q$ is not immune. Thus, Theorem 75 implies $S \cup T \not\leq^1_1 P \triangledown Q$. $\square$

To understand degrees of difficulty of disjunctive notions, and to discover new *easier* (possibly infinitary) disjunctive notions, it is interesting to discuss *contiguous degrees*.

Definition 77. Let $(\alpha,\beta,\gamma), (\alpha^*,\beta^*,\gamma^*) \in \{1, <\omega, \omega\}^3$, and assume that $\leq^{\alpha}_{\beta|\gamma}$ is not stronger than $\leq^{\alpha^*}_{\beta^*|\gamma^*}$. An $(\alpha,\beta|\gamma)$-degree $\mathbf{a}^{\alpha}_{\beta|\gamma}$ is $(\alpha^*,\beta^*|\gamma^*)$-*contiguous* if $\mathbf{a}^{\alpha}_{\beta|\gamma}$ contains at most one $(\alpha^*,\beta^*|\gamma^*)$-degree, that is to say, for any representatives $A, B \in \mathbf{a}^{\alpha}_{\beta|\gamma}$, we have that A is $(\alpha^*,\beta^*|\gamma^*)$-equivalent to B.

Corollary 78.

(1) *There is a $(1, <\omega)$-contiguous $(<\omega, 1)$-degree of Π^0_1 sets of $2^{\mathbb{N}}$.*
(2) *Every $(1, <\omega)$-degree which contains a homogeneous Π^0_1 set or a Π^0_1 antichain is not $(1,1)$-contiguous.*
(3) *Every $(1, \omega| <\omega)$-degree of Π^0_1 antichains is not $(1, <\omega)$-contiguous.*
(4) *Every $(<\omega, 1)$-degree of Π^0_1 antichains is not $(1,\omega)$-contiguous (hence, is not $(1,\omega| <\omega)$-contiguous).*

Proof. (1) This follows from Theorem 72.

(2) If $\mathbf{d}$ is a $(1, <\omega)$-degree of a homogeneous Π^0_1 set S, then $\mathbf{d}$ contains S and $S \triangledown S$, since $S \equiv^1_{<\omega} S \triangledown S$. However, $S \triangledown S <^1_1 S \cup S = S$ by Corollary 76. If $\mathbf{d}$ is a $(1, <\omega)$-degree of a Π^0_1 antichain P, then $\mathbf{d}$ contains $(P\times 2^{\mathbb{N}})\cup(2^{\mathbb{N}}\times P)$ and $P \triangledown P$, since $P \equiv^1_{<\omega} (P\times 2^{\mathbb{N}})\cup(2^{\mathbb{N}}\times P)$. However, $P \triangledown P <^1_1 (P\times 2^{\mathbb{N}})\cup(2^{\mathbb{N}}\times P)$ holds by Lemma 55.

(3) Note that, for any Π^0_1 set P and any clopen set C, it holds that $(P \cap C) \oplus (P \setminus C) \equiv^1_1 P$. Let $\mathbf{d}$ be a $(1, \omega| <\omega)$-degree of a Π^0_1 antichain P. Fix a clopen set C such that $P_0 = P \cap C \neq \emptyset$, and $P_1 = P \setminus C \neq \emptyset$. Then $\mathbf{d}$ contains $P_0 \oplus P_1$ and $\overrightarrow{\bigoplus}_n (P_0 \triangledown_n P_1)$, since $P_0 \oplus P_1 \equiv^1_{\omega|<\omega} \overrightarrow{\bigoplus}_n (P_0 \triangledown_n P_1)$. However, $\overrightarrow{\bigoplus}_n (P_0 \triangledown_n P_1) <^1_{<\omega} P_0 \oplus P_1$ holds by Corollary 59.

(4) Let $\mathbf{d}$ be a $(<\omega, 1)$-degree of a Π^0_1 antichain P. Fix a clopen set C such that $P_0 = P \cap C \neq \emptyset$, and $P_1 = P \setminus C \neq \emptyset$. Then $\mathbf{d}$ contains $P_0 \oplus P_1$ and $P_0 \triangledown_\omega P_1$, since $P_0 \oplus P_1 \equiv^{<\omega}_1 P_0 \triangledown_\omega P_1$. However, $P_0 \triangledown_\omega P_1 <_l P_0 \oplus P_1$ holds by Corollary 61. $\square$

4.2. **Dynamic Proof Models.** In the classical proof process, a verifier Ψ on "P_0 or P_1" may change his mind infinitely often. In the backtrack-tape model, this situation means that Ψ chooses the backtrack symbol $\sharp$ infinitely many often. Then the word on Λ is eventually finite, and it verifies neither P_0 nor P_1. Therefore, in the model, if Ψ succeeds to verify "P_0 or P_1" then the backtrack symbol $\sharp$ occurs on the record Δ at most finitely often. Consequently, in the backtrack-tape model, classical verification coincides with LCM verification. However, we would like to cover the case that infinitely many mind-changes occur. This may be archived by regarding the backtrack-tape model as a kind of infinitary tape model.

The dynamic-tape model: Assume that a directed graph (V, E) is given, where V can be infinite, $E \subseteq V \times V$, and an *initial vertex* $\varepsilon \in V$ is chosen. For any $v \in V$, let $\mathrm{adj}(v) = \{w \in V : (v, w) \in E\}$. When a verifier Ψ tries to prove that "$\bigvee_{v\in V} P_v$", infinite tapes $\square$, and Λ_v for $v \in V$ are given. The tape $\square$ is called *the declaration*, Λ_v is called *the working tape* for each $v \in V$. First the letter ε is written on $\square$, and no word is written on Λ_v for $v \in V$. At each stage s, assume that $v[s]$ is written on $\square$. Then the verifier Ψ executes one or the other of two following actions.

(1) Ψ declares some $w \in \mathrm{adj}(v[s])$, erases all words on $\square$, and writes w on $\square$; or

(2) Ψ writes a letter $k \in \mathbb{N}$ on the working tape $\Lambda_{v[s]}$.

Assume that a verifier Ψ tries to prove that "P_0 or P_1".

- **Intuitionism**: Consider $V = \{\varepsilon, 0, 1\}$, $E = \{(\varepsilon, 0), (\varepsilon, 1)\}$, and $P_\varepsilon = \emptyset$.
- **LCM with ordinal-bounded mind-changes**: For a computable well-founded tree $V = T \subseteq \mathbb{N}^{<\mathbb{N}}$, consider the following.
$$E = E(T) = \{(\sigma, \tau) \in T \times T : (\exists i \in \mathbb{N})\ \tau = \sigma^\frown i\},\ P_\sigma = \begin{cases} P_0, & \text{if } |\sigma| \text{ is even,} \\ P_1, & \text{if } |\sigma| \text{ is odd,} \end{cases}$$
- **LCM**: Consider $V = \mathbb{N}$; $E = \{(n, n+1) : n \in \mathbb{N}\}$; $P_{2n} = P_0$ for any $n \in \mathbb{N}$; and $P_{2n+1} = P_1$ for any $n \in \mathbb{N}$. Moreover, the word written on the declaration $\square$ must converge.
- (V, E)**-relaxed Classical**: $(V, E) = (V_0, V_1, E)$ is a given directed bipartite graph, and $P_\tau = P_i$ for any $\tau \in V_i$ and $i < 2$.

Definition 79 (Consistent Dynamic Disjunctions)**.** Let $G = (V, E)$ be a directed graph, $\{P_v\}_{v \in V}$ be a computable sequence of Π^0_1 subsets of $\mathbb{N}^\mathbb{N}$, and T_{P_v} be the corresponding tree for P_v. For $E \subseteq V^2$, put $\overline{E} = E \cup \{\langle v, v\rangle : v \in V\}$.

(1) $\bigtriangledown_{v \in (V,E)} P_v = \{f \in (V \times \mathbb{N})^\mathbb{N} : (\forall n \in \mathbb{N})\ (\langle (f(n))_0, (f(n+1))_0\rangle \in \overline{E})\ \&\ (\exists v \in V)\ \mathtt{pr}_v(f) \in P_v\} \cap \mathrm{Con}(T_{P_v})_{v \in V}$.

(2) $\blacktriangledown_{v \in (V,E)} P_v = \{f \in (V \times \mathbb{N})^\mathbb{N} : (\forall n \in \mathbb{N})\ (\langle (f(n))_0, (f(n+1))_0\rangle \in \overline{E})\} \cap \mathrm{Con}(T_{P_v})_{v \in V}$.

Here, recall that, for $x = (x_0, x_1)$, the first coordinate x_0 is denoted by $(x)_0$. If $P_v = P$ for any $v \in V$, then we simply write $\bigtriangledown_{v \in V} P$ and $\blacktriangledown_{v \in V} P$ for $\bigtriangledown_{v \in (V,E)} P_v$ and $\blacktriangledown_{v \in (V,E)} P_v$ respectively.

As our dynamic-tape model is an infinitary-tape model, this model may be natural to be regarded as expressing a proof process of an infinitary disjunction $\bigvee_{v \in V} P_v$. Therefore, we refer the model with (V, E) as *an infinitary disjunction along* (V, E). Later we will introduce a more complicated model. It will be called *the nested-tape model*. We first see an upper and lower bound of the degrees of difficulty of these disjunctive notions, and a relationship among various models we have introduced. Let $\widehat{\mathrm{Deg}}(P)$ denote the *Turing upward closure* of P, i.e., $\widehat{\mathrm{Deg}}(P) = \{g : (\exists f \leq_T g)\ f \in P\}$, and $[V]$ denote the set of all infinite paths through a graph (V, E), i.e., $[V] = \{p \in V^\mathbb{N} : (p(n), p(n+1)) \in E\}$.

Proposition 80. *Let* (V, E) *be a computable directed graph, and* $\{P_v\}_{v \in V}$ *be a computable sequence of* Π^0_1 *subsets of* $\mathbb{N}^\mathbb{N}$.

(1) $\widehat{\mathrm{Deg}}\left(\bigoplus_{v \in V} P_v\right) \leq^1_1 \bigtriangledown_{v \in (V,E)} P_v \leq^1_1 \bigoplus_{v \in V} P_v$.

(2) $\widehat{\mathrm{Deg}}\left([V] \oplus \bigoplus_{v \in V} P_v\right) \leq^1_1 \blacktriangledown_{v \in (V,E)} P_v \leq^1_1 [V] \oplus \bigoplus_{v \in V} P_v$.

Proof. (1) $\bigtriangledown_{v \in (V,E)} P_v \leq^1_1 \bigoplus_{v \in V} P_v$ is witnessed by $v^\frown f \mapsto \mathtt{write}(v, f) = v^\mathbb{N} \oplus f$. For any $f \in \bigtriangledown_{v \in (V,E)} P_v$, we have $\mathtt{pr}_v(f) \in P_v$ for some $v \in V$. Thus, we have $\mathtt{pr}_v(f) \leq_T f$, since $\mathtt{pr}_v$ is partially computable, and $f \in \mathrm{dom}(\mathtt{pr}_v)$. Hence, $f \in \widehat{\mathrm{Deg}}(P_v)$.

(2) Fix $f \in [V] \oplus \bigoplus_{v \in V} P_v$. If $f(0) = 1$, then we can show the desired condition as in (1). If f is of the form $f = 0^\frown g$, we have $\lambda n.\langle g(n), 0\rangle \in \blacktriangledown_{v \in (\mathbb{N}, S)} P_v$ since $g \in [V]$. Hence, $\blacktriangledown_{v \in (V,E)} P_v \leq^1_1 [V] \oplus \bigoplus_{v \in V} P_v$. To see $\widehat{\mathrm{Deg}}\left([V] \oplus \bigoplus_{v \in V} P_v\right) \leq^1_1 \blacktriangledown_{v \in (V,E)} P_v$, we inductively define a partial computable function $\mathtt{walk} :\subseteq (V \times \mathbb{N})^\mathbb{N} \to V^\mathbb{N}$ as follows. Set $\mathtt{walk}(\langle\rangle) = \langle\rangle$, and fix $\sigma = \sigma^{--\frown}\langle (u, m), (v, n)\rangle \in (V \times \mathbb{N})^{<\mathbb{N}}$. Assume that $\mathtt{walk}(\sigma^-)$ has been already defined. Then, $\mathtt{walk}(\sigma)$ is defined as follows.

$$\mathtt{walk}(\sigma^{--\frown}\langle (u, m), (v, n)\rangle) = \begin{cases} \mathtt{walk}(\sigma^-)^\frown\langle v\rangle & \text{if } v \neq u, \\ \mathtt{walk}(\sigma^-) & \text{otherwise.} \end{cases}$$

For any $f \in \blacktriangledown_{v \in (V,E)} P_v$, if $\mathtt{pr}_v(f)$ is total for some $v \in V$, then the desired condition follows as in (1). Otherwise, $\mathtt{mc}(f) = \infty$, i.e., there are infinitely many $n \in \mathbb{N}$ such that $(f(n+1))_0 \neq (f(n))_0$. In this case, $\mathtt{walk}(f) = \bigcup_{s \in \mathbb{N}} \mathtt{walk}(f \upharpoonright s)$ is an infinite path through the graph (V, E). In other words, the condition $f \in \blacktriangledown_{v\in(V,E)}P_v$ ensures that $\mathtt{pr}_v(f)$ is total and belongs to P_v for some $v \in V$, or otherwise $\mathtt{walk}(f)$ is total and belongs to $[V]$. Consequently, $f \in \widehat{\mathrm{Deg}}\left([V] \oplus \bigoplus_{v\in V} P_v\right)$, since $\mathtt{pr}_v$ and $\mathtt{walk}$ are partial computable. □

Proposition 81. *Let P, P_0, P_1, P_v, for $v \in V$, be Π^0_1 subsets of $\mathbb{N}^\mathbb{N}$, uniformly.*

1. $\bigtriangledown_{v\in(T,E(T))} P_v = \blacktriangledown_{v\in(T,E(T))}P_v$ *for any well-founded tree* $T \subseteq \mathbb{N}^{<\mathbb{N}}$.
2. $P_0 \oplus P_1 \equiv^1_1 \bigtriangledown_{v\in(V_1,E_1)} P_v \equiv^1_1 \blacktriangledown_{v\in(V_1,E_1)}P_v$, *where* $V_1 = \{\varepsilon, 0, 1\}$, $E_1 = \{(\varepsilon, 0), (\varepsilon, 1)\}$, *and* $P_\varepsilon = \emptyset$.
3. $P_0\triangledown P_1 \equiv^1_1 \bigtriangledown_{v\in(V_2,E_2)} P_v \equiv^1_1 \blacktriangledown_{v\in(V_1,E_1)}P_v$, *where* $P_\varepsilon = \emptyset$, $P_{01} = P_1$, $P_{10} = P_0$, $V_2 = \{\varepsilon, 0, 1, 01, 10\}$, *and* $E_2 = \{(\varepsilon, 0), (\varepsilon, 1), (0, 01), (1, 10)\}$, .
4. $[\![P_0 \vee P_1]\!]^2_{\mathsf{CL}} \equiv^1_1 \bigtriangledown_{v\in(\{0,1\},\{0,1\}^2)} P_v$.
5. $[\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}} \equiv^1_1 \bigtriangledown_{v\in(\mathbb{N},S)} P_v$, *where* $S = \{(n, n + 1) : n \in \mathbb{N}\}$; $P_{2n} = P_0$ *and* $P_{2n+1} = P_1$ *for any* $n \in \mathbb{N}$.
6. $[\![P \vee P]\!]^3_{\mathsf{LCM}} \equiv^1_1 [\![\bigvee_{n\in\mathbb{N}} P]\!]_{\mathsf{LCM}} \equiv^1_1 \bigtriangledown_{v\in(\mathbb{N},S)} P$.
7. $\widehat{\mathrm{Deg}}\left(\bigoplus_{v\in\mathbb{N}} P_v\right) \equiv^1_1 [\![\bigvee_{v\in\mathbb{N}} P_v]\!]_{\mathsf{CL}} \equiv^1_1 \bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v \equiv^1_1 [\bigtriangledown_\infty]_{v\in\mathbb{N}} P_v$.

Proof. (1) By Definition, $\bigtriangledown_v P_v \subseteq \blacktriangledown_v P_v$. On the other hand, any $f \in \blacktriangledown_{v\in(T,E(T))}P_v$ can pass at most finitely many vertices since $(T, E(T))$ has no infinite path. In other words, the set $\{(f(n))_0 : n \in \mathbb{N}\}$ is finite. By Pigeon Hole Principle, there is a vertex $v \in T$ such that $(f(n))_0 = v$ occurs for infinitely many $n \in \mathbb{N}$. Then, $\mathtt{pr}_v(f)$ must be infinite. Therefore, $\mathtt{pr}_v(f) \in [T_{P_v}] = P_v$ since $f \in \mathrm{Con}(T_{P_v})_{v\in V}$. Hence, $f \in \bigtriangledown_{v\in(T,E(T))} P_v$.

(2) The condition $\bigtriangledown_{v\in(V_1,E_1)} P_v \leq^1_1 P_0 \oplus P_1$ follows from Proposition 80 (1). For any $f \in \bigtriangledown_{v\in(V_1,E_1)} P_v$, there is $i < 2$ such that $(f(n))_0 = i$ for any $n \in \mathbb{N}$. Thus, $i^\frown f \in P_0 \oplus P_1$. The $(1, 1)$-equivalence of $\bigtriangledown_{v\in(V_1,E_1)} P_v$ and $\blacktriangledown_{v\in(V_1,E_1)}P_v$ follows from the item (1) since (V_1, E_1) is finite.

(3) Clearly, $P_0\triangledown_2 P_1 \subseteq \bigtriangledown_{v\in(V_2,E_2)} P_v$. Thus, by Proposition 22 (2), $P_0\triangledown P_1 \geq^1_1 \bigtriangledown_{v\in(V_2,E_2)} P_v$. For $f \in \bigtriangledown_{v\in(V_2,E_2)} P_v$, if $|(f(0))_0| = 1$ then $\Phi(f) = f \in P_0\triangledown_2 P_1$. If $|(f(0))_0| = 2$, say $(f(0))_0 = \langle i, j\rangle$, then $\Phi(f) = \mathtt{write}(j, \mathtt{pr}_j(f)) \in P_0\triangledown_1 P_1$. Hence, $P_0\triangledown P_1 \leq^1_1 \bigtriangledown_{v\in(V_2,E_2)} P_v$ via the computable function Φ. The $(1, 1)$-equivalence of $\bigtriangledown_{v\in(V_1,E_1)} P_v$ and $\blacktriangledown_{v\in(V_1,E_1)}P_v$ follows from the item (1) since (V_1, E_1) is finite.

(4) It is easy to see that $\bigtriangledown_{v\in(\{0,1\},\{0,1\}^2)} P_v = P_0\triangledown_\infty P_1$. Moreover, $[\![P_0 \vee P_1]\!]^2_{\mathsf{CL}} \equiv^1_1 P_0\triangledown_\infty P_1$ by Proposition 16.

(5) For each $\sigma = \tau^\frown\langle(i, m), (j, n)\rangle \in (\mathbb{N} \times \mathbb{N})^{<\mathbb{N}}$, we inductively define a computable function $\Xi(\sigma)$ as follows. If $i = j$, then we set $\Xi(\sigma) = \Xi(\tau^\frown\langle(i, m)\rangle)^\frown\langle n\rangle$. Otherwise, we set $\Xi(\sigma) = \Xi(\tau^\frown\langle(i, m)\rangle)^\frown\langle\sharp, j, n\rangle$. Then, $[\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}} \leq^1_1 \bigtriangledown_{v\in(\mathbb{N},S)} P_v$ is witnessed by Ξ. Conversely, to see $\bigtriangledown_{v\in(\mathbb{N},S)} P_v \leq^1_1 [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$, we again inductively define another computable function $\Xi^*(\sigma)$, for each $\sigma \in (\mathbb{N} \cup \{\sharp\})$. Set $\Xi^*(\langle\rangle) = \langle\rangle$, fix $\sigma = \sigma^- - {}^\frown\langle j, k\rangle \in (\mathbb{N} \cup \{\sharp\})^{<\mathbb{N}}$, and assume that $\Xi^*(\sigma^-)$ has been already defined. For $w \geq v + 2$, we consider the instruction $\mathtt{move}(v, w) = \langle(v + 1, 0), (v + 2, 0), \dots, (w - 2, 0), (w - 1, 0)\rangle \in (V \times \mathbb{N})^{w-v-1}$ to move from the tape Λ_v to the tape Λ_w in the dynamic tape model. If $w < v + 2$, then we assume that $\mathtt{move}(v, w)$ is the empty string. Put $p(\sigma) = 2 \cdot \mathtt{count}(\sigma) + \mathtt{tail}(\sigma; 0)$, where recall that $\mathtt{count}(\sigma) = \#\{n < |\sigma| : \sigma(n) = \sharp\}$. If $j \neq \sharp$ and $k \neq \sharp$, then we define $\Xi^*(\sigma) = \Xi^*(\sigma^-)^\frown\mathtt{move}(p(\sigma^*), p(\sigma))^\frown\langle(p(\sigma), k)\rangle$, where σ^* is the last string $\Xi^*(\sigma^*) \supsetneq \Xi^*((\sigma^*)^-)$. Otherwise, we set $\Xi^*(\sigma) = \Xi^*(\sigma^-)$. Then, we have $\langle(\Xi^*(f; n))_0, (\Xi^*(f; n + 1))_0\rangle \in \overline{S}$ for any $f \in [\![P_0 \vee P_1]\!]^3_{\mathsf{LCM}}$. It is easy to verify that $\Xi^*(f) \in \bigtriangledown_{v\in(\mathbb{N},S)} P_v$.

(6) The $(1, 1)$-equivalence of $[\![P \vee P]\!]^3_{\mathsf{LCM}}$ and $[\![\bigvee_{n\in\mathbb{N}} P]\!]_{\mathsf{LCM}}$ follows from Proposition 63 (2). Thus, the desired condition follows from (5).

(7) Clearly, $[\![\bigvee_{v\in\mathbb{N}} P_v]\!]_{\mathsf{CL}} \cap \mathrm{Con}(T_{P_v})_{v\in\mathbb{N}} = \bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v$. Thus, the equivalence $[\![\bigvee_{v\in\mathbb{N}} P_v]\!]_{\mathsf{CL}} \equiv^1_1 \bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v \equiv^1_1 [\bigtriangledown_\infty]_{v\in\mathbb{N}} P_v$ follows from Proposition 16 and 64. $\widehat{\mathrm{Deg}}\,(\bigcup_{v\in\mathbb{N}} P_v) \leq^1_1 \bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v$ follows from Proposition 80 (1). We may assume that $\Phi_e(\langle\rangle) = \langle\rangle$ for each index $e \in \mathbb{N}$. We inductively define a computable function Γ witnessing $\bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v \leq^1_1 \widehat{\mathrm{Deg}}\,(\bigcup_{v\in\mathbb{N}} P_v)$. For each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $e \in \mathbb{N}$, we also inductively define two parameters $\mathtt{act}_e(\sigma) \in \mathbb{N}$ and $\mathtt{rq}_e(\sigma) \in \mathbb{N} \cup \{-1\}$. Here, $\mathtt{act}_e(\sigma)$ will represent the last stage at which the e-th strategy acts along σ, and $\mathtt{rq}_e(\sigma) \geq 0$ will indicate that the e-th strategy



FIGURE 2. The dynamical representations of disjunction operations: (1) $[\![P \lor Q]\!]_{\mathsf{Int}}$ $(P \oplus Q)$; (2) $P^\frown Q$; (3) $[\![P \lor Q]\!]^2_{\mathsf{LCM}[2]}$ $(P \triangledown Q)$; (4) $[\![P \lor Q]\!]^2_{\mathsf{CL}}$ $(P \triangledown_\infty Q)$; (5) $[\![P \lor Q]\!]^3_{\mathsf{LCM}}$; (6) $\widehat{\mathrm{Deg}}(P)$, the Turing upward closure of P.

requires attention. First we set $\mathtt{act}_e(\langle\rangle) = 0$ and $\mathtt{rq}_e(\langle\rangle) = -1$ for each $e \in \mathbb{N}$. Inductively we assume that $\Gamma(\sigma^-)$, $\mathtt{act}_e(\sigma^-)$, and $\mathtt{rq}_e(\sigma^-)$ is already defined. Calculate $r = \min\{\mathtt{rq}_e(\sigma^-) : e < |\sigma| \ \& \ \mathtt{rq}_e(\sigma^-) > 0\}$, and pick the least e such that $\mathtt{rq}_e(\sigma^-) = r$ if such r and e exist. In this case, we say that e *acts*. If there is no such e, we set $\Gamma(\sigma) = \Gamma(\sigma^-)$, $\mathtt{act}_e(\sigma) = \mathtt{act}_e(\sigma^-)$, and $\mathtt{rq}_e(\sigma) = \mathtt{rq}_e(\sigma^-)$. If there is such e, put $\sigma^* = (\Phi_e(\sigma))^{\leftarrow|\Phi_e(\sigma \upharpoonright |\mathtt{act}_e(\sigma)|)|}$, i.e., $\Phi_e(\sigma) = (\Phi_e(\sigma \upharpoonright |\mathtt{act}_e(\sigma)|)|)^\frown\sigma^*$. Then we set $\Gamma(\sigma) = \Gamma(\sigma^-)^\frown\mathtt{write}(e, \sigma^*)$. Then, put $\mathtt{rq}_e(\sigma) = -1$ and $\mathtt{act}_e(\sigma) = |\sigma|$. For each $e^* \in \mathbb{N} \setminus \{e\}$, set $\mathtt{act}_{e^*}(\sigma) = \mathtt{act}_{e^*}(\sigma^-)$. Moreover, if $e^* \leq |\sigma|$, $\mathtt{rq}_{e^*}(\sigma^-) = -1$, and $|\Phi_{e^*}(\sigma \upharpoonright |\mathtt{act}_{e^*}(\sigma)|)| < |\Phi_{e^*}(\sigma)|$, then declare $\mathtt{rq}_{e^*}(\sigma) = |\sigma|$. Otherwise, put $\mathtt{rq}_{e^*}(\sigma) = \mathtt{rq}_{e^*}(\sigma^-)$. Fix $g \in \mathbb{N}^\mathbb{N}$. We claim that $\Phi_e(g)$ act infinitely often whenever $\Phi_e(g)$ is total. Our construction ensures that only finitely many e's require attentions along $g \upharpoonright s$ for each $s \in \mathbb{N}$. Therefore, for $R = \{e \in \mathbb{N} : \mathtt{rq}_e(g \upharpoonright s) > 0\}$, if $e \in R$, then the strategy e acts by stage $s + \#R$, i.e., $\mathtt{act}_e(g \upharpoonright s + \#R) \geq s$. Assume that e act at stage $t \in \mathbb{N}$. Then the algorithm $\Gamma(g \upharpoonright t)$ writes the new information $(g \upharpoonright t)^*$ of $\Phi_e(g)$ on the e-th tape, i.e., $\mathtt{pr}_e(\Gamma(g \upharpoonright t)) = \Phi_e(g \upharpoonright t)$. Thus, eventually, we have $\mathtt{pr}_e(\Gamma(g)) = \Phi_e(g)$. For any $g \in \widehat{\mathrm{Deg}}\,(\bigcup_{v\in\mathbb{N}} P_v)$, there is an index $e \in \mathbb{N}$ such that $\Phi_e(g) \in P_v$ for some $v \in \mathbb{N}$. Consequently, $\Gamma(g) \in \bigtriangledown_{v\in(\mathbb{N},\mathbb{N}^2)} P_v$. □

Proposition 82. *Let (V, E) be a computable directed graph, and $\{P_v\}_{v\in V}$ be a computable sequence of Π^0_1 subsets of $2^\mathbb{N}$. Then we have the following.*

(1) $\bigtriangledown_{v\in(V,E)} P_v$ *is* Σ^0_3.
(2) $\blacktriangledown_{v\in(V,E)} P_v$ *is* Π^0_1.

Proof. Clearly, $\mathrm{Con}(T_{P_v})_{v\in V}$ is Π^0_1. Moreover, the relation $\langle (f(n))_0, (f(n+1))_0\rangle \in \overline{E}$ is computable, uniformly in $f \in (\mathbb{N} \times \mathbb{N})^\mathbb{N}$ and $n \in \mathbb{N}$. Thus, $\blacktriangledown_{v\in(V,E)} P_v$ is Π^0_1. The relation $\mathtt{pr}_v(f) \in P_v$ is Π^0_2 in $v \in V$ and $f \in \mathbb{N}^\mathbb{N}$, since it is equivalent to the following formula.

$$(\forall n \in \mathbb{N})(\exists m \in \mathbb{N})\ |\mathtt{pr}_v(f \upharpoonright m)| > n \ \& \ \mathtt{pr}_v(f \upharpoonright m) \in T_{P_v}.$$

Therefore, $\bigtriangledown_{v\in(V,E)} P_v$ is Σ^0_3. □

4.3. Concatenation, Dynamic Disjunctions, and Contiguous Degrees.

Theorem 83. *For any Π^0_1 set $P, Q \subseteq 2^\mathbb{N}$, $Q^\frown P$ does not $(1,1)$-cup to P. That is to say, for any $R \subseteq \mathbb{N}^\mathbb{N}$, if $P \leq^1_1 (Q^\frown P) \otimes R$ then $P \leq^1_1 R$.*

Proof. Fix a computable tree V_P (resp. V_Q) such that $[V_P] = P$ (resp. $[V_Q] = Q$), and let L_P (resp. L_Q) denote all leaves of V_P (resp. V_Q). For a tree $T \subseteq 2^{<\mathbb{N}}$ and $g \in \mathbb{N}^\mathbb{N}$, we write $T \otimes \{g\}$ for $\{\sigma \oplus \tau : \sigma \in V_P \ \& \ \tau \subset g \ \& \ |\sigma| = |\tau|\}$. For computable trees S and T, we also write $S^\frown T$ for $S \cup \bigcup_{\rho\in L_S} \rho^\frown T$, where L_S denotes the set of all leaves of S. Assume $P \leq^1_1 (Q^\frown P) \otimes R$ via a computable functional Φ. We construct a computable functional Ψ witnessing $P \leq^1_1 R$. Fix $g \in R$. Then we will find a g-c.e. tree $D^g \subseteq V_P$ without dead ends. To this end, we inductively construct a uniformly g-computable sequences $\{D^g_i\}_{i\in\omega}, \{E^g_i\}_{i\in\omega}$ of g-computable trees, as follows.

$$E^g_0 = V_Q \otimes \{g\}; \qquad D^g_0 = \Phi(E^g_0).$$
$$E^g_{i+1} = (V_Q{}^\frown D^g_i) \otimes \{g\}; \qquad D^g_{i+1} = \Phi(E^g_{i+1}).$$

Here $\Phi(E^g_i)$ denotes the image of E^g_i under a functional Φ, namely, $\Phi(E^g_i) = \{\tau \subseteq 2^{<\mathbb{N}} : (\exists \sigma \in E^g_i)\ \tau \subseteq \Phi(\sigma)\}$. Finally, we define a g-c.e. tree $D^g = \bigcup_n D^g_n$. Now, we let W be the tree $V_Q{}^\frown V_P$, and then we observe $[W] = Q^\frown P$ and $V_Q \subseteq W^{ext}$.

Lemma 84. *For any i, $D^g_i \subseteq V^{ext}_P$ and $E^g_i \subseteq W^{ext} \otimes \{g\}$.*

Proof. This lemma is proved by induction. First, our assumption $V_Q \subseteq W^{ext}$ ensures $E^g_0 = V_Q \otimes \{g\} \subseteq W^{ext} \otimes \{g\}$, and we also have $D^g_0 = \Phi(E^g_0) \subseteq V^{ext}_P$ since $\Phi((Q^\frown P) \otimes R) \subseteq [V_P]$ implies $\Phi(W^{ext} \otimes \{g\}) \subseteq V^{ext}_P$ for $g \in R$. Assume the lemma holds for each $j \leq i$. We now show that the lemma also holds for $i+1$. By assumption, $V_Q{}^\frown D^g_i \subseteq V_Q{}^\frown V^{ext}_P = W^{ext}$. So by definition of E^g_{i+1}, we get $E^g_{i+1} \subseteq W^{ext} \otimes \{g\}$. Furthermore, we observe $D^g_{i+1} = \Phi(E^g_{i+1}) \subseteq \Phi(W^{ext} \otimes \{g\}) \subseteq V^{ext}_P$. ◻

Lemma 85. *There is a computable function Γ mapping each $g \in R$ to a g-computable sequence $\Gamma(g) = \{D^g_n\}_{n\in\omega}$ of g-computable trees.*

Proof. We inductively show this lemma. It suffices to show that we can construct D^g_i from E^g_i by uniformly g-computable way, since clearly E^g_0 is computable in g, and $D^g_i \mapsto E^g_{i+1}$ is uniformly g-computable. Assume that $E^g_i \subseteq 2^{<\mathbb{N}} \otimes \{g\}$ is given. For each $\sigma \in 2^{\mathbb{N}}$, if $\sigma \oplus (g \upharpoonright |\sigma|) \in E^g_i$, then put $l(\sigma) = |\Phi(\sigma \oplus (g \upharpoonright |\sigma|))|$. If $\sigma \oplus (g \upharpoonright |\sigma|) \notin E^g_i$, then put $l(\sigma) = \infty$. Note that $l : 2^{<\mathbb{N}} \to \mathbb{N} \cup \{\infty\}$ is g-computable, since the notation $\Phi(\sigma)$ just means the computation of Φ restricted to step $|\sigma|$ with the oracle σ. By Lemma 84, $\lim_n l(f \upharpoonright n) = \infty$ for any $f \in 2^{\mathbb{N}}$. Because, for any f with $f \oplus g \in [E^g_i]$, we have $\Phi(f \oplus g) \in [\Phi(E^g_i)] \subseteq \Phi([W] \otimes \{g\}) \subseteq \Phi((Q^\frown P) \otimes R) \subseteq P$, hence, $f \oplus g \in \mathrm{dom}(\Phi)$. Therefore, by compactness, for each $n \in \mathbb{N}$, there is $h_n \in \mathbb{N}$ such that $l(\sigma) \geq n$ for each $\sigma \in 2^{<\mathbb{N}}$ of length h_n. We can compute $h^g_i(n) = h_n$ with the oracle g, since l is g-computable. Here, we can compute a g-computable index of h^g_i from an index of E^g_i, uniformly in $i \in \mathbb{N}$ and $g \in \mathbb{N}^{\mathbb{N}}$. Thus, the relation $\tau \in D^g_i$ is equivalent to the g-computable condition that $\tau \subseteq \Phi(\sigma)$ for some $\sigma \in E^g_i$ of length $h^g_i(|\tau|)$, uniformly in $i \in \mathbb{N}$ and $g \in \mathbb{N}^{\mathbb{N}}$. Formally, the set $\{(\tau, i, g) \in 2^{<\mathbb{N}} \times \mathbb{N} \times \mathbb{N}^{\mathbb{N}} : \tau \in D^g_i\}$ is computable. ◻

Define L_{D_i} as the set of all leaves of the tree D^g_i, and define L_{E_i} as the set of all leaves of the tree E^g_i for each i.

Lemma 86. *Let X be D or E, and i be any natural number. For any $\rho \in L^g_{X_i}$, there are infinitely many nodes $\tau \in L^g_{X_{i+1}}$ which are extensions of ρ.*

Proof. This lemma is proved by induction. First we pick $\rho \in L_{E_0} = L_Q \otimes \{g\} = \{\sigma \oplus \tau : \sigma \in L_Q \ \&\ \tau \subset g \ \&\ |\sigma| = |\tau|\}$. We note that V_P is an infinite tree since P is special. By using our assumption $P \leq^1_1 (Q^\frown P) \otimes R$ via Φ and the property $[V_Q] \otimes \{g\} \subseteq (Q^\frown P) \otimes R$, the tree $D^g_0 = \Phi(E^g_0)$ has a path, i.e., it is infinite. By definition, we have $E^g_1 = V_P{}^\frown D^g_0 \supseteq \rho^\frown D^g_0$, and so E^g_1 has infinitely many extensions of ρ. Now, we assume this lemma for E and any $j \leq i$. For a given $\rho \in L_{D_i}$, there is a node $\sigma \in E^g_i$ such that $\Phi(\sigma) = \rho$ by our definition of $D^g_i = \Phi(E^g_i)$. Without loss of generality, we can pick σ as a leaf of E^g_i. By induction hypothesis, σ has infinitely many extensions in E^g_{i+1}. By lemma 84, we know $E^g_{i+1} \subseteq W^{ext} \otimes \{g\}$. This implies that $\Phi(E^g_{i+1}(\supseteq \sigma))$ must be infinite whenever $E^g_{i+1}(\supseteq \sigma)$ is infinite. We now remark that, for any $\sigma' \in \Phi(E^g_{i+1}(\supseteq \sigma))$, $\Phi(\sigma') \supseteq \Phi(\sigma) = \rho$. Thus, $\Phi(E^g_{i+1}(\supseteq \sigma))$ gives infinitely many extensions of ρ, and our definition $D^g_{i+1} = \Phi(E^g_{i+1})$ clearly implies the lemma for D and i. Now, we will show the lemma for E and $i+1$. By our definition of $E^g_{i+1} = (V_Q{}^\frown D^g_i) \otimes \{g\}$, every $\rho \in L_{E_{i+1}}$ must be of form

$\rho = (\sigma^\frown\tau) \oplus (g \upharpoonright |\sigma^\frown\tau|)$ for some $\sigma \in L_Q$ and $\tau \in L_{D_i}$. So if $\tau \in L_{D_i}$ has infinitely many extensions in $L_{D_{i+1}}$ then $\rho = (\sigma^\frown\tau) \oplus (g \upharpoonright |\sigma^\frown\tau|)$ has infinitely many extensions in $L_{E_{i+2}}$. Thus, we have established the lemma for E and $i+1$. Now, the lemma follows by induction. □

Lemma 87. *D^g is a g-c.e. subtree of V_P without dead ends for any $g \in P$. Hence, D^g has a g-computable path of V_P.*

Proof. For a given g-c.e. tree $D^g = \{\delta_s\}_{s\in\omega}$, we first compute $\Delta(D^g)[0] = \delta_0$. If $\Delta(D^g)[t]$ has already been defined, then we wait for stage s with $\delta_s \supsetneq \Delta(D^g)[t]$. At this point, we extend $\Delta(D^g)[t+1]$ to σ_s. We eventually reach this stage since V_P has no dead ends. This procedure is g-computable and $\Delta(D^g) = \lim_t \Delta(D^g)[t]$ must be a g-computable path of D^g. □

Then we define a computable functional Ψ as $\Psi(g) = \Delta(\bigcup_n \Gamma(g))$ for any $g \in \mathbb{N}^\mathbb{N}$. This witnesses $P \leq^1_1 R$ as desired. □

Corollary 88. *For every special Π^0_1 set $P \subseteq 2^\mathbb{N}$, there exists a Π^0_1 set $Q \subseteq 2^\mathbb{N}$ such that $Q <^1_1 P$ and $Q \equiv^1_{<\omega} P$.*

Proof. By Theorem 83, we have $P^\frown P <^1_1 P \equiv^1_{<\omega} P$. □

Definition 89. Fix $\alpha, \beta, \gamma \in \{1, <\omega, \omega\}$. An $(\alpha,\beta|\gamma)$-degree $\mathbf{a} \in \mathcal{P}^\alpha_{\beta|\gamma}$ has *the strong anticupping property* if there is a nonzero $(\alpha,\beta|\gamma)$-degree $\mathbf{b} \in \mathcal{P}^\alpha_{\beta|\gamma}$ such that, for any $(\alpha,\beta|\gamma)$-degree $\mathbf{c}$, if $\mathbf{a} \leq \mathbf{b} \vee \mathbf{c}$, then $\mathbf{a} \leq \mathbf{c}$.

Corollary 90. *Every nonzero $\mathbf{a} \in \mathcal{P}^1_1$ has the strong anticupping property.*

Proof. Fix $P \in \mathbf{a}$. Let $\mathbf{b}$ be the $(1,1)$-degree of $P^\frown P$. Then, by Theorem 83, for any $(1,1)$-degree $\mathbf{c}$, if $\mathbf{a} \leq \mathbf{b} \vee \mathbf{c}$, then $\mathbf{a} \leq \mathbf{c}$. □

Definition 91. For a Π^0_1 set P, we define *the weakening of P* as $P^\triangledown = \mathsf{CPA}^\frown P$, where recall that CPA denotes the set of all complete consistent extensions of Peano Arithmetic, and it is a $(1,1)$-complete Π^0_1 subset of $2^\mathbb{N}$.

By previous theorem, the weakening $P^\triangledown$ does not $(1,1)$-cup to P. In particular, we have $P^\triangledown <^1_1 P$.

Remark. The specialness of P may represent that, intuitionistically, P is neither provable nor refutable, i.e., undecidable. Additionally, we can think of $P \otimes Q$ as representing the conjunction of P and Q (both intuitionistic and classical sense). Thus, the previous theorem expresses that, if an undecidable proposition P can be represented as a Π^0_1 subset of $2^\mathbb{N}$, then P is not (intuitionistically) derivable from the conjunction $P^\triangledown \wedge R$ of the weakening $P^\triangledown$ and any proposition R which does not derive P intuitionistically.

Proposition 92. *For any Π^0_1 sets $P_0, P_1, Q_0, Q_1 \subseteq 2^\mathbb{N}$, if $P_0 \leq^1_1 Q_0$ and $P_1 \leq^1_1 Q_1$, then ${P_0}^\frown P_1 \leq^1_1 {Q_0}^\frown Q_1$. Hence, the operator $\triangledown : \mathcal{P}^1_1 \to \mathcal{P}^1_1$ introduced by $(\deg^1_1(P))^\triangledown = \deg^1_1(P^\triangledown)$ is well-defined.*

Proof. Assume that $P_0 \leq^1_1 Q_0$ via a computable function Φ_0, and $P_1 \leq^1_1 Q_1$ via a computable function Φ_1. Then we can construct a computable function Φ^*_0 with $\Phi^*_0(T_{Q_0}) \subseteq T_{P_0}$ by modifying Φ_0 as follows. For each $\sigma \in T_{Q_0}$, if $\Phi_0(\sigma)$ extends a leaf $\rho \in L_{P_0}$, then we set $\Phi^*_0(\sigma) = \rho$. If $\sigma \in L_{Q_0}$, then let $\Phi^*_0(\sigma)$ be the leftmost leaf of L_{P_0} extending $\Phi_0(\sigma)$. Otherwise, we set $\Phi^*_0(\sigma) = \Phi_0(\sigma)$. Then, for each $\sigma = \rho^\frown\tau$ with $\rho \in L_{Q_0}$ and $\tau \in T_{Q_1}$, we put $\Phi(\sigma) = \Phi^*_0(\rho)^\frown\Phi_1(\tau)$. Clearly Φ is computable, and ${P_0}^\frown P_1 \leq^1_1 {Q_0}^\frown Q_1$ via Φ. □

Recall that $\mathcal{P}^1_1(\leq \mathbf{1}^\triangledown) = \{\mathbf{a} \in \mathcal{P}^1_1 : \mathbf{a} \leq \mathbf{1}^\triangledown\}$ is a principal prime ideal consisting of tree-immune-free Medvedev degrees [19]. Here, recall that a Π^0_1 set $P \subseteq 2^\mathbb{N}$ is tree-immune if T^{ext}_P contains no infinite computable subtree. Then, we also observe the following.

Proposition 93. *Fix Π^0_1 sets $P_0, P_1, Q_0, Q_1 \subseteq 2^\mathbb{N}$, and assume that ${P_0}^\frown P_1 \leq^1_1 {Q_0}^\frown Q_1$. Then, either $P_0 \leq^1_1 Q_0$ or $P_1 \leq^1_1 Q_1$ holds. Moreover, if P_0 is tree-immune and Q_0 is nonempty, then $P_1 \leq^1_1 Q_1$.*

Proof. Assume that $P_0{}^\frown P_1 \leq^1_1 Q_0{}^\frown Q_1$ via a computable function Φ. If $\Phi(\rho) \in T^{ext}_{P_0}$ for any leaf $\rho \in L_{Q_0}$, then $\Phi(g) \in [T_{P_0}]$ for any $g \in [Q_0]$, i.e., $P_0 \leq^1_1 Q_0$. If $\Phi(\rho) \notin T^{\rm ext}_{P_0}$ for some leaf $\rho \in L_{Q_0}$, then there are only finitely many strings of T_{P_0} extending $\Phi(\rho)$. Thus, $[T_{P_0}{}^\frown T_{P_1}] \cap [\Phi(\rho)]$ is essentially a sum of finitely many P_1's, hence it is $(1,1)$-equivalent to P_1. Since a computable functional Φ maps $\rho^\frown Q_1$ to the above class, obviously, $P_1 \leq^1_1 Q_1$. If P_0 is tree-immune, then $\Phi(\rho) \notin T^{\rm ext}_{P_0}$ for some leaf $\rho \in L_{Q_0}$, since otherwise the image of T_{Q_0} under Φ is included in T_{P_0}, and clearly it is infinite and computable. Therefore, we must have $P_1 \leq^1_1 Q_1$. □

Corollary 94. *The weakening* $\triangledown : \mathbf{a} \mapsto \mathbf{a}^\triangledown$ *is injective. Hence,* $\triangledown$ *provides an order-preserving self-embedding of the* $(1,1)$*-degrees* $\mathcal{P}^1_1$ *of nonempty* Π^0_1 *subsets of* $2^{\mathbb{N}}$.

Proof. By Cenzer-Kihara-Weber-Wu [19], CPA is tree-immune. Therefore, by Proposition 93, $\mathsf{CPA}^\frown Q = Q^\triangledown \leq^1_1 P^\triangledown = \mathsf{CPA}^\frown P$ implies $Q \leq^1_1 P$. □

4.4. Infinitary Disjunctions along well-Founded Trees.

Definition 95. *The n-weakening* $P^{(n)}$, for each $n \in \mathbb{N}$, is inductively defined as follows: $P^{(1)} = P$, and $P^{(n+1)} = P^\frown P^{(n)}$.

Remark. One may think of $P^{(n)}$ as the mass problem requiring to construct a solution to the original problem P with mind-change-bound n, or as the set of all oracles which tell us the information how we can construct an element of P with mind-change-bound n. On the sequent calculus of Limit Computable Mathematics, the number of times of the use of *the weakening rule* is related to the number of times of mind-changes (see Berardi-Yamagata [6] and Tatsuta-Berardi [71]). This is one of the reasons why we call $P^{(n)}$ the n-weakening of P.

Proposition 96. *For any* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there is a* $(1, n+1)$*-computable function* $\Gamma : P^{(n+1)} \to P$.

Proof. Each $\sigma \in 2^{<\mathbb{N}}$ is represented to be of the form $\sigma = \rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_k{}^\frown\tau$ for $\rho_i \in L_P$, $i \leq k$, and $\tau \in T_P$. Then we consider $\mathtt{cut}(\sigma) = \langle \rho_0, \rho_1, \dots, \rho_k, \tau\rangle$, and $\mathtt{tail}^{\mathtt{cut}}(\sigma) = \mathtt{cut}(\sigma; |\mathtt{cut}(\sigma) - 1|) = \tau$. Moreover, $\mathtt{tail}^{\mathtt{cut}}(g)$ is defined by $\lim_s \mathtt{tail}^{\mathtt{cut}}(g \upharpoonright s)$ if it converges. Then, $\mathtt{tail}^{\mathtt{cut}}$ is learnable via a learner Ψ who guesses an index of $g \mapsto g^{\leftarrow \sum_{i<|\mathtt{cut}(\sigma)-1|} |\mathtt{cut}(\sigma;i)|}$ on σ. Then, clearly $\mathtt{tail}^{\mathtt{cut}} : P^{(n+1)} \to P$. For any $g \in P^{(n+1)}$, the learning procedure of $\mathtt{tail}^{\mathtt{cut}}(g)$ changes at most $n+1$ times. □

Proposition 97. *For any* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$ *and any set* $Q \subseteq \mathbb{N}^{\mathbb{N}}$*, if* $P \leq^1_{tt,<\omega} Q$ *via a* $(1,n)$*-truth-table function, then* $P^{(n)} \leq^1_1 Q$.

Proof. Assume that $\Theta : P^{(n+1)} \otimes Q$ is $(1,n)$-truth-table computable via n many total computable functions $\Theta_0, \dots, \Theta_{n-1}$. We define a computable function $\gamma : n \times 2^{<\mathbb{N}} \to 2^{<\mathbb{N}}$ as follows. If $\Theta_m(\sigma) \in T_P$, then put $\gamma(m,\sigma) = \Theta_m(\sigma)$. If $\Theta_m(\sigma) \supsetneq \rho$ for some $\rho \in L_P$, then we define $\gamma(m,\sigma)$ to be such ρ. Let $z(\sigma) = \min\{m < n : \Theta_m(\sigma) \in T_P\}$. Then, for $\sigma \in 2^{<\mathbb{N}}$, the value $\Phi(\sigma)$ is defined by $\bigsqcap_{m \leq z(\sigma)} \gamma(m,\sigma)$. Then Φ ensures that $P^{(n)} \leq^1_1 Q$. □

Corollary 98. *For any* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$ *and any set* $R \subseteq \mathbb{N}^{\mathbb{N}}$*, if a* $(1,n)$*-truth-table function* $\Theta : P^{(n+1)} \otimes R \to P$ *exists, then a* $(1,n)$*-computable function* $\Theta^* : R \to P$ *also exists.*

Proof. By applying Proposition 97 to $Q = P^{(n+1)} \otimes R$, we have $P^{(n)} \leq^1_1 P^{(n+1)} \otimes R$. So $P^{(n)} \leq^1_1 R$ by Theorem 83, since $P^{(n+1)} = P^\frown P^{(n)}$. Since a $(1,n)$-computable function $P^{(n)} \to P$ exists by Proposition 96, a $(1,n)$-computable function $R \to P$ exists, as desired. □

Definition 99 (Transfinite Weakening). Let $(\mathcal{O}, \leq_{\mathcal{O}})$ denote *Kleene's system of ordinal notations* (see Rogers [57]). Then for each $a \in \mathcal{O}$ we define *the a-weakening* $P^{(a)}$ as follows.

1. $P^{(0)} = P$.
2. If $a = 2^b$, then $P^{(a)} = P^\frown P^{(b)}$.

(3) If $a = 3 \cdot 5^e$, then $P^{(a)} = P^\frown\{P^{(\Phi_e(n))}\}_{n\in\mathbb{N}}$.

Remark. The transfinite levels of mind-change hierarchy of Δ^0_2 reals are well-known as *the Ershov hierachy* introduced in 1968 by Ershov (see Stephan-Yang-Yu [70]). The transfinite levels of mind-change hierarchy in machine learning has been introduced by Freivalds-Smith [32].

Proposition 100. *For any Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ and any notation $a \in \mathcal{O}$, the a-th weakening $P^{(a)}$ is a Π^0_1 subset of $2^{\mathbb{N}}$.*

Proof. Fix $a \in \mathcal{O}$. By our definition, obviously $P^{(a)}$ is a subset of $2^{\mathbb{N}}$. We inductively assume that $\{P^{(b)} : b <_{\mathcal{O}} a\}$ is uniformly Π^0_1. For $a = 2^b$, we can easily compute a Π^0_1 index of $P^{(a)} = P^\frown P^{(b)}$ is from a Π^0_1 index of $P^{(b)}$. For $a = 3 \cdot 5^e$, we can also easily compute a Π^0_1 index of $P^a = P^\frown\{P^{(\Phi_e(n))}\}_{n\in\mathbb{N}}$ from a computable sequence of Π^0_1 indices of $\{P^{(\Phi_e(n))}\}_{n\in\mathbb{N}}$. Thus, $\{P^{(b)} : b \leq_{\mathcal{O}} a\}$ is uniformly Π^0_1. $\square$

Definition 101. For each $a \in \mathcal{O}$, we let $|a|_{\mathcal{O}}$ be the order-type of $\{b \in \mathcal{O} : b <_{\mathcal{O}} a\}$. The ordinal ω_1^{CK} denotes *the Church-Kleene ω_1*, the first non-constructive ordinal, i.e., $\omega_1^{CK} = \sup\{|a|_{\mathcal{O}} : a \in \mathcal{O}\}$. Recall that every computable ordinal is obtained from a computable well-founded tree. For each well-founded tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we inductively define $|\sigma|_T : T \to \omega_1^{CK}$ by $|\sigma|_T = \sup\{|\tau|_T : \tau \in T \;\&\; \tau \supsetneq \sigma\}$, where $\sup\emptyset = 0$. Then, *the order type of T* is defined by $o(T) = |\langle\rangle|_T$.

We may introduce the notion of mass problem with mind-change-bound α, by using a corresponding infinitary disjunction along a well-founded tree of order type α. However, an infinitary disjunction along a well-founded is, generally, not a subset of Cantor space $2^{\mathbb{N}}$. Meanwhile, each a-weakening is a subset of $2^{\mathbb{N}}$ as seen in Proposition 100. We now see that each a-weakening can be think of as a learning procedure with $|a|_{\mathcal{O}}$-bounded-mind-changes, with precision ±1.

Proposition 102. *For any Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ and any notation $a \in \mathcal{O}$, there are a well-founded tree $T_a \subseteq \mathbb{N}^{<\mathbb{N}}$ with $o(T_a) = |a|_{\mathcal{O}}$ and a notation $a' \in \mathcal{O}$ with $|a'|_{\mathcal{O}} = |a|_{\mathcal{O}} + 1$ such that $P^{(a')} \leq^1_1 \bigtriangledown_{v\in(T_a,E(T_a))} P_v \leq^1_1 P^{(a)}$ holds, where $P_v = P$ for each $v \in T_a$.*

Proof. As in the usual argument, each notation $a \in \mathcal{O}$ generates a computable well-founded tree T_a, by defining the tree-homomorphism $t_a : \mathcal{O}(\leq_{\mathcal{O}} a) \to \mathbb{N}^{<\mathbb{N}}$ as follows, where $\mathcal{O}(\leq_{\mathcal{O}} a) = \{b \in \mathcal{O} : b \leq_{\mathcal{O}} a\}$. Put $t_a(a) = \langle\rangle$. Fix $b \leq_{\mathcal{O}} a$, and assume that $t_a(b)$ has been already defined. In the case that b is successor, say $b = 2^c$ for $c \in \mathcal{O}$, we put $t_a(c) = t_a(b)^\frown 0$. If b is limit, say $b = 3 \cdot 5^e$ for $e \in \mathbb{N}$ such that $\Phi_e(n) \downarrow\in \mathcal{O}$, and $\Phi_e(n) <_{\mathcal{O}} \Phi_e(n+1)$ for each $n \in \mathbb{N}$, then we put $t_a(\Phi_e(n)) = t_a(b)^\frown n$ for each $n \in \mathbb{N}$. Then we define $T_a = \{t_a(b) : b \leq_{\mathcal{O}} a\}$. It is easy to see that T_a is a computable well-founded tree, and $o(T_a) = |a|_{\mathcal{O}}$. Note that we have $|t(b)|_{T_a} = |b|_{\mathcal{O}}$ for each $b \leq_{\mathcal{O}} a$.

We also define a computable function $R : \mathbb{N} \to \mathbb{N}$ by recursion theorem. Put $R(0) = 0$, $R(2^c) = 2^{R(c)}$, and $R(3 \cdot 5^e) = 2^{3\cdot 5^{r(e)}}$ for each $c, e \in \mathbb{N}$, where $\Phi_{r(e)}(n) = R(\Phi_e(n))$. Then, put $a' = R(a)$. We claim that $|a'|_{\mathcal{O}} \leq |a|_{\mathcal{O}} + 1$. By transfinite induction, we show that $|R(b)|_{\mathcal{O}} \leq |b|_{\mathcal{O}} + 1$ for any $b \leq_{\mathcal{O}} a$. By definition, $|R(0)|_{\mathcal{O}} = |0|_{\mathcal{O}}$, Fix $b \leq_{\mathcal{O}} a$, and assume that $|R(d)|_{\mathcal{O}} \leq |d|_{\mathcal{O}} + 1$ is satisfied for any $d <_{\mathcal{O}} b$. For successor step $b = 2^c$, we have $|R(2^c)|_{\mathcal{O}} = |2^{R(c)}|_{\mathcal{O}} = |R(c)|_{\mathcal{O}} + 1 \leq |c|_{\mathcal{O}} + 2 = |2^c|_{\mathcal{O}} + 1$. For limit step $b = 3 \cdot 5^e$, $|R(3\cdot 5^e)|_{\mathcal{O}} = |2^{3\cdot 5^{r(e)}}|_{\mathcal{O}} + 1 = |3\cdot 5^{r(e)}|_{\mathcal{O}} + 1 = \sup_{n\in\mathbb{N}}(|R(\Phi_e(n))|_{\mathcal{O}}) + 1 \leq \sup_{n\in\mathbb{N}}(|\Phi_e(n)|_{\mathcal{O}} + 1) + 1 = |3\cdot 5^e|_{\mathcal{O}} + 1$, since $\{|\Phi_e(n)|_{\mathcal{O}}\}_{n\in\mathbb{N}}$ is strictly increasing. Thus, $|a'|_{\mathcal{O}} \leq |a|_{\mathcal{O}} + 1$.

For each $\zeta \in T_a$, the *ζ-root of* $\bigtriangledown_{v\in(T_a,E(T_a))} P_v$ is a unique ρ_ζ which is extendible in $\bigtriangledown_{v\in(T_a,E(T_a))} P_v$ with $\mathtt{pr}_\zeta(\sigma) = \langle\rangle$. Note that $\rho_\zeta \subseteq \rho_\xi$ whenever $\zeta \subseteq \xi$. Then, $C_\zeta = \rho_\zeta{}^\frown\mathtt{write}(\zeta, T_P)$ is called *the component of* ζ, where $\mathtt{write}(\zeta, T_P) = \{\mathtt{write}(\zeta,\sigma) : \sigma \in T_P\}$ (recall that $\mathtt{write}(\zeta,\sigma) = \zeta^{|\sigma|} \oplus \sigma = \langle(\zeta,\sigma(0)), (\zeta,\sigma(1)), \dots, (\zeta,\sigma(|\sigma|-1))\rangle$). Then, it is easy to see that $\bigcup_{\zeta\in T_a} C_\zeta$ is just the corresponding tree of $\bigtriangledown_{v\in(T_a,E(T_a))} P_v$ whenever $\zeta \neq \xi$.

We next assign some $b \in \mathcal{O}(\leq_{\mathcal{O}} a) \cup \{-1\}$ to each $\sigma \in T^{(a)}$, where we use $T^{(a)}$ to denote the corresponding tree of $P^{(a)}$. Every $\sigma \in 2^{<\mathbb{N}}$ can be represented to be of the form $\pi_{k(0)}{}^\frown\pi_{k(1)}{}^\frown\dots{}^\frown\pi_{k(n)}{}^\frown\tau$ for some $k(0), k(1), \dots, k(n) \in \mathbb{N}$, and some $\tau \in T_P$. Here, π_n is the n-th leaf in L_P. Put $\mathtt{cut}(\sigma) =$

$\langle k(0), k(1), \dots, k(n)\rangle$. We define *the change-recording* $\mathtt{record}_a(\sigma) \in (\mathcal{O}(\leq_{\mathcal{O}} a))^{<\mathbb{N}}$ *along* σ *of the mind-change-counter* as follows. Put $\mathtt{record}_a(\sigma; 0) = a$, and fix $i < |\mathtt{cut}(\sigma)|$. Assume that $\mathtt{record}_a(\sigma; 0)$ is already defined.

- If $\mathtt{record}_a(\sigma; i) \leq 0$, then set $\mathtt{record}_a(\sigma; i+1) = -1$
- If $\mathtt{record}_a(\sigma; i)$ is of the form 2^b, then set $\mathtt{record}_a(\sigma; i+1) = b$.
- If $\mathtt{record}_a(\sigma; i)$ is of the form $3 \cdot 5^e$, then set $\mathtt{record}_a(\sigma; i+1) = \Phi_e(\mathtt{cut}(\sigma; i))$.

Note that $\mathtt{record}$ is computable. Let us define $\mathtt{ord}(\sigma) \in \mathcal{O}(\leq_{\mathcal{O}} a)$ to be $\mathtt{record}_a(\sigma; |\mathtt{record}_a(\sigma)| - 1)$. Then the relation $g \in P^{(a)}$ is characterized by the condition that $\mathtt{ord}(\sigma) \geq 0$ for any $\sigma \subseteq g$. Define $T^{(a)}_b = \{\sigma \in T^{(a)} : \mathtt{ord}(\sigma) = b\}$ for each $b \leq_{\mathcal{O}} a$. Then $T^{(a)}_b$ has a root α_b, and indeed, $T^{(a)}_b = \alpha_b{}^\frown T_P$. Note that t_a is the computable tree-isomorphism between $\mathcal{O}(\leq_{\mathcal{O}} a)$ and T_a, and $\{T^{(a)}_b\}_{b \leq_{\mathcal{O}} a}$ is also uniformly computable. Then, the set of all infinite paths of the computable tree $\bigcup_{b \leq_{\mathcal{O}} a} T^{(a)}_b$ coincides with $P^{(a)}$.

We first show that $\bigtriangledown_{v \in (T_a, E(T_a))} P_v \leq^1_1 P^{(a)}$. We consider a computable mapping $\Gamma : \alpha_b \mapsto \rho_{t_a(b)}$. For each $\alpha_b{}^\frown\sigma \in T^{(a)}_b$, we define $\Gamma(\alpha_b{}^\frown\sigma) = \Phi(\alpha_b)^\frown\mathtt{write}(t_a(b), \sigma)$. Clearly, $\Gamma(\sigma) \in C_{t_a(b)}$ for each $\sigma \in T^{(a)}_b$. Thus, $\bigtriangledown_{v \in (T_a, E(T_a))} P_v \leq^1_1 P^{(a)}$ via Γ. We construct a computable function Δ witnessing $P^{(a')} \leq^1_1 \bigtriangledown_{v \in (T_a, E(T_a))} P_v$. Set $\Delta(\rho_{\langle\rangle}) = \Delta(\langle\rangle) = \langle\rangle$. Fix $\zeta \in T_a$. Inductively assume that $\Delta(\rho^-_\zeta)$ is already defined. Note that ρ^-_ζ is a leaf of C_{ζ^-}. Then $\Delta(\rho^-_\zeta)$ is of the form $\sigma^\frown\tau$, where τ is the tail of $\Delta(\rho^-_\zeta)$. By uniformly partial computable way, we can extend the tail τ to $\mathtt{leaf}(\tau) \in L_P$. If ζ^- has just one successor in T_a, i.e., $\zeta^{-\frown}1 \notin T_a$, then we set $\Delta(\rho_\zeta) = \sigma^\frown\mathtt{leaf}(\tau) \supseteq \Delta(\rho^-_\zeta)$. If ζ^- has infinitely many successors in T_a, i.e., $\zeta^{-\frown}1 \in T_a$, then we set $\Delta(\rho_\zeta) = \sigma^\frown\mathtt{leaf}(\tau)^\frown\pi_{\zeta(|\zeta|-1)} \supseteq \Delta(\rho^-_\zeta)$. For each $\rho_\zeta{}^\frown\gamma \in C_\zeta$, we set $\Delta(\rho_\zeta{}^\frown\gamma) = \Delta(\rho_\zeta)^\frown\gamma$. Put $R(\zeta) = t_a(R(b))$ for each $\zeta \in T$, where $b \in \mathcal{O}$ is the unique notation satisfying $t_a(b) = \zeta$. Then, $\Delta(\sigma) \in T^{(a')}_{R(\zeta)}$ for each $\sigma \in C_\zeta$. Consequently, $P^{(a')} \leq^1_1 \bigtriangledown_{v \in (T, E(T))} P_v$ via Δ. □

Proposition 103. *For any special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$, *if* $a, b \in \mathcal{O}$ *and* $a <_{\mathcal{O}} b$, *then* $P^{(b)}$ *does not* $(1,1)$*-cup to* $P^{(a)}$, *i.e., for any set* $R \subseteq \mathbb{N}^{\mathbb{N}}$, *if* $P^{(a)} \leq^1_1 P^{(b)} \otimes R$ *then* $P^{(a)} \leq^1_1 R$.

Proof. The assumption $a <_{\mathcal{O}} b$ implies $2^a \leq_{\mathcal{O}} b$. Therefore, we have $P^{(b)} \leq^1_1 P^{(2^a)}$. By Theorem 83, $P^{(2^a)}$ does not $(1,1)$-cup to $P^{(a)}$. Thus, $P^{(b)}$ does not $(1,1)$-cup to $P^{(a)}$. □

Fix any notation $\mathtt{omega} \in \mathcal{O}$ such that $|\Phi_{\mathtt{omega}}(n)|_{\mathcal{O}} = n$ for each $n \in \mathbb{N}$. Note that $|\mathtt{omega}|_{\mathcal{O}} = \omega$.

Proposition 104. *Let* P *be a special* Π^0_1 *subset of* $2^{\mathbb{N}}$, *and* $a \in \mathcal{O}$ *be any notation. For any set* $R \subseteq \mathbb{N}^{\mathbb{N}}$, *if* $P^{(a)} \leq^1_{tt,<\omega} P^{(a \cdot_{\mathcal{O}} \mathtt{omega})} \otimes R$, *then* $P^{(a)} \leq^1_{<\omega} R$.

Proof. By the definition of $\leq^1_{tt,<\omega}$, if $P^{(a)} \leq^1_{tt,<\omega} P^{(a \cdot_{\mathcal{O}} \mathtt{omega})} \otimes R$, then there is a $(1,n)$-truth-table function $\Gamma : P^{(a \cdot_{\mathcal{O}} \mathtt{omega})} \otimes R \to P^{(a)}$ for some $n \in \mathbb{N}$. In particular, $\Gamma : (\rho_{n+1}{}^\frown P^{(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n+1))}) \otimes R \to P^{(a)}$, where ρ_{n+1} is the $(n+1)$-th leaf of T_P. By modifying Γ, we can easily construct a $(1,n)$-truth-table function $\Theta : P^{(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n+1))} \otimes R \to P^{(a)}$. By Corollary 98, we get a $(1,n)$-computable function $\Theta^* : R \to P^{(a)}$. Thus, $P^{(a)} \leq^1_{<\omega} R$. □

Let Ψ be a learner. We say that $c : \mathbb{N}^{<\mathbb{N}} \to \mathcal{O}$ is a *mind-change counter* for Ψ if, for any $\sigma \in \mathbb{N}^{<\mathbb{N}}$, $c(\sigma) <_{\mathcal{O}} c(\sigma^-)$ whenever $\Psi(\sigma) \neq \Psi(\sigma^-)$. A learner Ψ is α*-bounded* if there is a computable mind-change counter $c : \mathbb{N}^{<\mathbb{N}} \to \mathcal{O}$ for Ψ such that $|c(\langle\rangle)|_{\mathcal{O}} \leq \alpha$.

Proposition 105. *Let* P *be a* Π^0_1 *subset of* $2^{\mathbb{N}}$, *and let* $a \in \mathcal{O}$ *be any notation. Then* $P \leq^1_\omega P^{(a)}$ *via some* $|a|_{\mathcal{O}}$*-bounded learner.*

Proof. Recall the definition of $\mathtt{record}_a : 2^{<\mathbb{N}} \to (\mathcal{O} \cup \{-1\})^{<\mathbb{N}}$ in the proof of Proposition 100. For $g \in P^{(a)}$, by counting $\mathtt{record}_a(g)$, we can easily see that $g \mapsto \mathtt{tail}^{\mathtt{cut}}(g)$ is $(1,\omega)$-computable via an $|a|_{\mathcal{O}}$-bounded learner. Here the partial function $\mathtt{tail}^{\mathtt{cut}} : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is defined as in Proposition 96. □

Corollary 106. *For every $a \in \mathcal{O}$ there exists a computable function g such that, for any Π^0_1 index e, if P_e is special then the following properties hold.*

1. *$P_{g(e,b)} <^1_1 P_{g(e,c)}$ holds for every $c <_{\mathcal{O}} b <_{\mathcal{O}} a$, indeed, $P_{g(e,b)}$ does not $(1,1)$-cup to $P_{g(e,c)}$.*
2. *$P_{g(e,b)} \equiv^1_\omega P_{g(e,c)}$ for every $b, c <_{\mathcal{O}} a$.*

Proof. Let $g(e,b)$ be an index of $P_e^{(b)}$. Then, the desired conditions follow from Proposition 103. □

For any reducibility notion r, and any ordered set $(I, \leq_I)$, an I-indexed sequence $\{\mathbf{a}_i\}_{i \in I}$ of r-degrees is *r-noncupping* if, for any $i <_I j$, the condition $\mathbf{a}_i \leq_r \mathbf{b}$ must be satisfied whenever $\mathbf{a}_i \leq_r \mathbf{a}_j \vee \mathbf{b}$, for any r-degree $\mathbf{b}$. In particular, any r-noncupping sequence is strictly decreasing, in the sense of r-degrees.

Corollary 107. *For any nonzero $(1,\omega)$-degree $\mathbf{a} \in \mathcal{P}^1_\omega$, there is a $(1,1)$-noncupping sequence of $(1,1)$-degrees inside $\mathbf{a}$ of length ω_1^{CK}. (Of course, such a sequence is not computable).*

Proof. Fix a Π^0_1 set $P \in \mathbf{a}$. Pick a path $p \subseteq \mathcal{O}$ through $\mathcal{O}$ of order type ω_1^{CK}. Then, $\{P^{(a)}\}_{a \in p}$ is a desired sequence by Proposition 103. □

4.5. **Infinitary Disjunctions along the Straight Line.** We introduce the LCM disjunctions of $\{P_i\}_{i \in \mathbb{N}}$ as $\bigtriangledown_{n \in \mathbb{N}} P_n = \bigcup_{n \in \mathbb{N}} (P_0 \triangledown \dots \triangledown P_n)$ and $\overrightarrow{\bigtriangledown}_{n \in \mathbb{N}} P_n = \bigcup_{n \in \mathbb{N}} (P_0{}^\frown \dots {}^\frown P_n)$. This is a straightforward infinitary iteration of the commutative and noncommutative concatenations. If $P_n = P$ for all $n \in \mathbb{N}$, we write $\bigtriangledown P$ for $\bigtriangledown_n P_n = \overrightarrow{\bigtriangledown}_n P_n$. In this case, to simplify our argument, we again think of $\triangledown$ as the non-commutative disjunction ${}^\frown$.

Remark. In the context of Computability Logic [39], Japaridze [40] have introduced the *sequential existential quantifier* $\bigtriangledown nA(n)$ and the *sequential corecurrence* $\bigtriangledown A$. We fix some interpretation $[\![\cdot]\!] : \mathsf{Snt} \to \mathcal{P}(\mathbb{N}^\mathbb{N})$, where Snt is the set of all sentences of a fixed language in Computability Logic. Then, our disjunction operations $\overrightarrow{\bigtriangledown}_n [\![A(n)]\!]$ and $\bigtriangledown [\![A]\!]$ coincide with the existential quantifier $\bigtriangledown nA(n)$ and the corecurrence $\bigtriangledown A$, respectively.

Proposition 108. *Let $\{P_n\}_{n \in \mathbb{N}}$ be a computable collection of Π^0_1 subsets of Baire space $\mathbb{N}^\mathbb{N}$.*

1. *$\bigoplus_n P_n \equiv^1_\omega \bigtriangledown_n P_n$.*
2. *$\overrightarrow{\bigtriangledown}_n P_n \equiv^1_1 \bigtriangledown_{v \in (\mathbb{N},S)} P_v$, where S is the successor relation.*
3. *$\left[\bigtriangledown_\omega\right]_{\langle n,m\rangle} P_{n,m} \leq^1_1 \bigtriangledown_{\langle n,m\rangle} P_{n,m} \leq^1_1 \left[\bigtriangledown_\omega\right]_n P_n \leq^1_1 \bigtriangledown_n P_n$, where $P_{n,m} = P_n$ for each $n, m \in \mathbb{N}$.*

Proof. (1) Clearly $\left[\bigtriangledown_\omega\right]_n P_n \leq^1_1 \bigtriangledown_n P_n$, and by Proposition 66. (2) For each $\sigma \in (\mathbb{N} \times \mathbb{N})^{<\mathbb{N}}$, we calculate $\mathtt{count}(\sigma) = \max\{m : \mathtt{pr}_m(\sigma) \neq \langle\rangle\}$. Then $\overrightarrow{\bigtriangledown}_n P_n \leq^1_1 \bigtriangledown_{v \in (\mathbb{N},S)} P_v$ is witnessed by $\sigma \mapsto \bigsqcap_{i \leq \mathtt{count}(\sigma)} \mathtt{pr}_i(\sigma)$, where recall that the concatenation is denoted by $\bigsqcap$. Each $\sigma \in \mathbb{N}^\mathbb{N}$ can be represented as $\sigma = \rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_n{}^\frown\tau$, where $\rho_m \in L_{P_m}$ for any $m \leq n$, and $\tau \in T_{P_{n+1}}$. Then, we define $\mathtt{cut}(\sigma) = \langle \rho_0, \rho_1, \dots, \rho_n, \tau\rangle$. Then $\Phi(\sigma) = \bigsqcap_{i < |\mathtt{cut}(\sigma)|} \mathtt{write}(i, \mathtt{cut}(\sigma; i))$ witnesses $\bigtriangledown^\frown_n P_n \geq^1_1 \bigtriangledown_{v \in (\mathbb{N},S)} P_v$.

(3) We only show $\bigtriangledown_{\langle n,m\rangle} P_{n,m} \leq^1_1 \left[\bigtriangledown_\omega\right]_n P_n$, since other relations are trivial. Each $\sigma \in \mathbb{N}^\mathbb{N}$ can be represented as $\sigma = \rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_u{}^\frown\tau$ with $\rho_v \in L_{P_{(v)_0}}$ for any $v \leq u$, and $\tau \in T_{P_{(u+1)_0}}$, where we think of each $u \in \mathbb{N}$ as a pair $\langle (u)_0, (u)_1\rangle$. Then we write $m(\sigma) = u$, and put $L(\sigma) = \rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_u{}^\frown\mathrm{leaf}(\tau)$, where $\mathrm{leaf}(\tau)$ is the least leaf of $T_{P_{(u+1)_0}}$ extending τ. Fix $\sigma = \tau^\frown\langle (i,m),(j,n)\rangle \in (\mathbb{N} \times \mathbb{N})^{<\mathbb{N}}$. Assume that $\Phi(\tau^\frown\langle (i,m)\rangle)$ is already defined. If $i = j$, then we set $\Phi(\sigma) = \Phi(\tau^\frown\langle (i,m)\rangle)^\frown\langle n\rangle$. If $i \neq j$, then we calculate the least $k \in \mathbb{N}$ such that $\langle j,k\rangle > m(\Phi(\tau^\frown\langle (i,m)\rangle))$. For such k, put $K = \langle j,k\rangle - m(\Phi(\tau^\frown\langle (i,m)\rangle)) - 1$. Then we define $\Phi(\sigma) = L(\Phi(\tau^\frown\langle (i,m)\rangle))^\frown L(\langle\rangle)^{K\frown}\mathtt{pr}_j(\sigma)$. Here $L(\langle\rangle)^K$ denotes the concatenation of K many $L(\langle\rangle)$'s. It is not difficult to see that $\bigtriangledown_{\langle n,m\rangle} P_{n,m} \leq^1_1 \left[\bigtriangledown_\omega\right]_n P_n$ via Φ. □

Proposition 109. *Let $\{P_i\}_{i \in \mathbb{N}}$ be a computable collection of nonempty Π^0_1 subsets of $2^\mathbb{N}$. Then $\bigtriangledown_n P_n$ is $(1,1)$-equivalent to a dense Σ^0_2 set in Cantor space $2^\mathbb{N}$.*

Proof. Let S denote the set $\{g \in (\mathbb{N} \cup \{\sharp\})^{\mathbb{N}} : (\exists n \in \mathbb{N})\ (\mathtt{count}(g) = n \ \&\ \mathtt{tail}(g) \in P_n)\}$, where $\mathtt{count}(g) = \#\{n \in \mathbb{N} : g(n) = \sharp\}$. Then, S is clearly a Σ^0_2 subset of $\{0, 1, \sharp\}^{\mathbb{N}}$. For any $\sigma \in \{0, 1, \sharp\}^{<\mathbb{N}}$, we have $\sigma^\frown\langle\sharp\rangle^\frown h \in S$ for any $h \in P_{\mathtt{count}(\sigma)+1}$ Thus, S intersects with any clopen set. □

Example 110. Let MLR denote the set of all Martin-Löf random reals. Then $\mathsf{MLR} \equiv^1_1 \bigtriangledown P$ for any nonempty Π^0_1 set $P \subseteq \mathsf{MLR}$, by Kučera-Gács Theorem, while $\mathsf{MLR} <^1_1 P$ for any Π^0_1 set $P \subseteq \mathsf{MLR}$ as follows.

Proposition 111 (Lewis-Shore-Sorbi [48])**.** *No somewhere dense set in Baire space* $(1, 1)$*-cup to a closed set in Baire space. In other words, for any somewhere dense set* $D \subseteq \mathbb{N}^{\mathbb{N}}$*, any closed set* $C \subseteq \mathbb{N}^{\mathbb{N}}$*, and any set* $R \subseteq \mathbb{N}^{\mathbb{N}}$*, if* $C \leq^1_1 D \otimes R$ *then* $C \leq^1_1 R$*.* □

Proposition 112. *For any somewhere dense set* $D \subseteq \mathbb{N}^{\mathbb{N}}$ *and any special closed set* $C \subseteq \mathbb{N}^{\mathbb{N}}$*, we have* $C \not\leq^{<\omega}_1 D$*.*

Proof. If $\{D_i\}_{i<b}$ is a finite partition of D, then $\bigcup_{i<b} \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D_i) = \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D)$, where the topological closure of D, in the standard Baire topology on $\mathbb{N}^{\mathbb{N}}$, is denoted by $\mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D)$. To show this claim, for every $x \in \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D)$ we have a sequence $\{x_k\}_{k\in\mathbb{N}} \subseteq D$ converging to x. By pigeonhole principle, there is $i < b$ such that there are infinitely many k such that $x_k \in D_i$. For such i, clearly $x \in \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D_i)$. However, since the somewhere density of D implies that $\mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D)$ contains some clopen set, and hence $\mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D_i)$ contains a computable element r for some i. Additionally, $\mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(C) = C$ since C is closed. If $C \leq^{<\omega}_1 D \otimes R$, then there is a finite partition $\{D_i\}_{i<b}$ of D such that $C \leq^1_1 D_i$ via a computable function $\Phi_{e(i)}$. Fix i such that $\mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D_i)$ contains a computable element. Therefore, $C = \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(C) \leq^1_1 \mathrm{Cl}_{\mathbb{N}^{\mathbb{N}}}(D_i) \supseteq \{r\}$ via $\Phi^f_{e(i)}$. Hence, C contains a computable element. □

Especially, if P is a special Π^0_1 set, then there is no nonzero $(< \omega, 1)$-degree of Π^0_1 subsets of $2^{\mathbb{N}}$ below the $(< \omega, 1)$-degree of $\bigtriangledown P$. We will see that the set $\bigtriangledown P$ has a stronger property.

Definition 113. A sequence $\langle t_n \rangle_{n\in\mathbb{N}}$ of finite strings is *a timekeeper* if there is a uniformly c.e. collection of finite sets, $\{V_n\}_{n\in\mathbb{N}}$, such that, for any $n \in \mathbb{N}$, $|t_n| = |V_n|$ and $t_n(i)$ is given as the stage at which the i-th element is enumerated into V_n, for each $i < |t_n|$.

Definition 114. For a finite string $\tau \in \mathbb{N}^{<\mathbb{N}}$, *the* τ*-delayed* $(|\tau| + 1)$*-weakening* $P^{(\tau)}$ is inductively defined as follows:

$$P^{(\tau\upharpoonright 0)} = P; \qquad P^{(\tau\upharpoonright i+1)} = \bigcup\{\sigma^\frown P : \sigma \in L_{P^{(\tau\upharpoonright i)}} \ \&\ |\sigma| \geq \tau(i)\} \text{ for each } i < |\tau|.$$

Proposition 115. *If* $\tau(m) = 0$ *for each* $m < |\tau|$*, then* $P^{(\tau)} = P^{(|\tau|+1)}$*.*

Proof. Straightforward from the definition. □

Lemma 116. *For any timekeeper* $\langle t_n \rangle_{n\in\mathbb{N}}$*, the following conditions hold.*

(1) $P^{(t_n)} \subseteq P^{(|t_n|+1)}$*. Hence,* $\bigoplus^{\longrightarrow}_n P^{(t_n)} \subset \bigtriangledown P$*.*
(2) $P^{(t_n)}$ *is* Π^0_1*, uniformly in n. Hence,* $\bigoplus^{\longrightarrow}_n P^{(t_n)}$ *is* Π^0_1*.*

Proof. (1) Straightforward. (2) We construct a computable tree $T^{(t_n)}$ corresponding to $P^{(t_n)}$. Each $\sigma \in 2^{\mathbb{N}}$ can be represented as $\sigma = \rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_k{}^\frown\tau$, where $\rho_m \in L_P$ for any $m \leq k$, and $\langle\rangle \neq \tau \in T_P$. Then $\sigma \in T^{(t_n)}$ if and only if $t_n(k)$ holds by stage $|\rho_0{}^\frown\rho_1{}^\frown \dots {}^\frown\rho_k|$. Then $T^{(t_n)}$ is a computable tree, and clearly $P^{(t_n)} = [T^{(t_n)}]$. □

Remark. Our *finite injury priority argument* often yields a timekeeper. Therefore, the delayed weakening construction is useful, since the recursive meet $\bigoplus^{\longrightarrow}_n P^{(|t_n|+1)}$ of the standard weakenings along a timekeeper $\{t_n\}_{n\in\mathbb{N}}$ is only assured to be $\Pi^{0,\emptyset'}_1$.

Theorem 117. *Let P be any* Π^0_1 *subset of* $2^{\mathbb{N}}$*. Then, for every special* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$*, there exists a* Π^0_1 *set* $\widehat{P} \subseteq \bigtriangledown P$ *such that* $Q \not\leq^1_{<\omega} \widehat{P}$*.*

Proof. Let Q be a special Π^0_1 set, and P be a given Π^0_1 set. By a uniformly computable procedure, from P, we will construct a timekeeper $\{t_n\}_{n\in\mathbb{N}}$. The desired class $\widehat{P}$ will be given by $\widehat{P} = \overrightarrow{\bigoplus}_n P^{(t_n)}$.

Requirements. We need to ensure, for all $n \in \mathbb{N}$, the following:

$$R_n \ : \ Q \leq^1_{<\omega} \widehat{P} \text{ via } n \ \rightarrow \ (\exists \Delta_n)\ \Delta_n \in Q.$$

Action of an R_n-strategy. Fix an effective enumeration $\{\rho_n : n \in \mathbb{N}\}$ of all leaves of T_P. An R_n-strategy uses nodes extending the n-th leaf ρ_n of T_P, and it constructs a finite sequence $t_n[s]$, a sequence $\tau_n[s]$ of strings, and a computable functional Δ_n. For any n, put $t_n[0] = \langle\rangle$, and $\tau_n[0] = \rho_n$ at stage 0. *An R_n-strategy acts at stage $s+1$ if the following condition holds:*

$$(\exists \rho \in T^s_P)(\exists e < n)\ \Phi_e(\tau_n[s]{}^\frown\rho) \in T_Q \ \&\ \Phi_e(\tau_n[s]{}^\frown\rho) \supsetneq \Phi_e(\tau_n[s]).$$

If an R_n-strategy acts at stage $s+1$ then, for a witness $\rho \in T^s_P$, we pick $\rho^* \in L_P$ extending ρ. Then let us define $\tau_n[s+1] = \tau_n[s]{}^\frown\rho^*$, $t_n[s+1] = t_n[s]{}^\frown\langle|\tau_n[s+1]|\rangle$, and $\Delta_{e,n} \upharpoonright l = \Phi_e(\tau_n[s+1])$, where l is the length of $\Phi_e(\tau_n[s+1])$. Otherwise, $t_n[s+1] = t_n[s]$, $\tau_i[s+1] = \tau_i[s]$. Note that the mapping $(n,m) \mapsto \tau_n(m)$ is partial computable. At the end of the construction, set $t_n = \bigcup_s t_n[s]$. As mentioned above, $\widehat{P}$ is defined by $\widehat{P} = \overrightarrow{\bigoplus}_n P^{(t_n)}$.

Claim. An R_n-strategy acts at most finitely often for each n.

Clearly $\tau_n = \bigcup_s \tau_n[s]$ is a computable string. If R_n acts infinitely often, then $\Delta_{e,n} = \Phi_e(\tau_n) \in Q$ for some $e < n$ by our choice of τ_n. Since $\Phi_e(\tau_n)$ is computable, Q contains a computable element. However, this contradicts our assumption that Q is special. Therefore, we concludes the claim. As a corollary, $\langle t_n\rangle_{n\in\mathbb{N}}$ is a timekeeper.

Claim. $P \not\leq^1_{<\omega} \widehat{P}$.

Let $\tau_n = \bigcup_s \tau_n[s]$. By induction we show that $\tau_n \in \rho_n{}^\frown T^{ext}_{P(t_n)}$. First we have the following observation:

$$\tau_n[0] = \rho_n \in \rho_n{}^\frown T^{ext}_P = \rho_n{}^\frown T^{ext}_{P(t_n\upharpoonright 0)} \subseteq \rho_n{}^\frown T^{ext}_{P(t_n[0])}.$$

Assume $\tau_n[s] \in \rho_n{}^\frown T^{ext}_{P(t_n[s])}$. If $\tau_n[s+1] = \tau_n[s]{}^\frown\rho^*$ for $\rho^* \in L_P$ then $t_n[s+1] = t_n[s]{}^\frown\langle|\tau_n[s+1]|\rangle$. In particular $\tau_n[s+1] \in \rho_n{}^\frown L_{P(t_n[s+1]\upharpoonright|t_n[s]|)}$ and $|\tau_n[s+1]| \geq t_n[s+1](|t_n[s]|)$. Hence, by the definition of $P^{(t_n[s+1])}$, it is easy to see that $\tau_n[s+1]{}^\frown P \subseteq \rho_n{}^\frown P^{(t_n[s+1])}$. Thus, $\tau_n[s+1] \in \rho_n{}^\frown T^{ext}_{P(t_n[s+1])}$. So we obtain $\tau_n \in \rho_n{}^\frown T^{ext}_{P(t_n)}$ and by our construction of τ_n there is no $\rho \in P$ and $e < n$ such that $\Phi_e(\tau_n{}^\frown\rho) \supsetneq \Phi_e(\tau_n)$. Since $\Phi_e(\tau_n)$ is a finite string, for any $g \in \rho_n{}^\frown T^{ext}_{P(t_n)} \subset \hat{P}$ extending τ_n, $\Phi_e(g)$ is also a finite string. Consequently, this g witnesses that $P \not\leq^1_{<\omega} \widehat{P}$. □

Corollary 118. (1) *For every special Π^0_1 set $P \subseteq 2^\mathbb{N}$, we have $\bigtriangledown P <^{<\omega}_1 P \equiv^1_\omega \bigtriangledown P$.*

(2) *For every special Π^0_1 set $P \subseteq 2^\mathbb{N}$ there exists a Π^0_1 set $Q \subseteq 2^\mathbb{N}$ with $Q <^{<\omega}_1 P \equiv^1_\omega Q$.*

Proof. By applying Theorem 117 to $Q = P$, we have $P \not\leq^1_{<\omega} \widehat{P} \geq^1_{<\omega} \bigtriangledown P$. Moreover, $P \oplus \widehat{P} <^{<\omega}_1 P \equiv^1_\omega P \oplus \widehat{P}$. □

4.6. **Infinitary Disjunctions along ill-Founded Trees.** To establish the nonexistence theorem on $(1,\omega)$-contiguous $(<\omega,\omega)$-degrees, the team-learning proof model of P is expected to be useful. We can directly introduce *the proof model of P by a team of n learners* as $[\bigtriangledown_\infty]_{i<n} \bigtriangledown P$.

Proposition 119. *$P \equiv^{<\omega}_\omega [\bigtriangledown_\infty]_{i<n} \bigtriangledown P$ for any $P \subseteq \mathbb{N}^\mathbb{N}$ and $n \in \mathbb{N}$.*

Proof. Each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ is represented as $\sigma = \rho_0{}^\frown\rho_1{}^\frown \ldots {}^\frown\rho_n{}^\frown\tau$, where $\rho_i \in L_P$ for each $i \leq n$ and $\tau \in T_P$. Then we put $\mathtt{tail}^{\mathtt{cut}}(\sigma) = \tau$. Let Ψ_i be a learner who guesses an index of $\mathtt{tail}^{\mathtt{cut}}(\mathtt{pr}_i(\sigma))$ for each $\sigma \in \mathbb{N}^{<\mathbb{N}}$, where note that $\mathtt{tail}^{\mathtt{cut}} \circ \mathtt{pr}_i$ is learnable, since $\mathtt{pr}_i$ is partially computable and $\mathtt{tail}^{\mathtt{cut}}$ is learnable. For any $g \in [\bigtriangledown_\infty]_{i<n} \bigtriangledown P$, obviously $\mathtt{tail}^{\mathtt{cut}}(\mathtt{pr}_i(g)) \in P$. Thus, $P \equiv^{<\omega}_\omega [\bigtriangledown_\infty]_{i<n} \bigtriangledown P$ via the team $\{\Psi\}_{i<n}$. □

However, this model may be far from Π^0_1 whenever P is Π^0_1. To break out of the dilemma, the following minor modification of consistent dynamic disjunction is helpful. For any tree $T_P \subseteq \mathbb{N}^{<\mathbb{N}}$ and $i \in \mathbb{N}$, we let $T_P{}^\frown\langle i\rangle$ denote the tree $T_P \cup \bigcup_{\rho\in L_P} \rho^\frown\langle i\rangle$, and $T_P{}^\frown T_Q$ denote the tree $T_P \cup \bigcup_{\rho\in L_P} \rho^\frown T_Q$. In other words, $T_P{}^\frown T_Q$ is a corresponding tree of $P^\frown Q$.

Definition 120. Let V be a subtree of $\mathbb{N}^{<\mathbb{N}}$, $\{P_\sigma\}_{\sigma\in V}$ be a computable sequence of Π^0_1 subsets of $2^{\mathbb{N}}$, and $T_\sigma \subseteq 2^{<\mathbb{N}}$ be the corresponding tree of P_σ for each $\sigma \in V$. Then *the concatenation of* $\{P_\sigma\}_{\sigma\in V}$ *along the tree* V is defined as follows.

$$\blacktriangledown_{\sigma\in V} P_\sigma = \left[\bigcup_{\tau\in V}\left(\bigsqcap_{i<|\tau|} T_{\tau\upharpoonright i}{}^\frown\langle\tau(i)\rangle\right)^\frown T_\tau\right].$$

Proposition 121. *Let V be a computable subtree of $2^{<\mathbb{N}}$, and $\{P_\sigma\}_{\sigma\in V}$ be a computable sequence of Π^0_1 subsets of $2^{\mathbb{N}}$. Then $\blacktriangledown_{\sigma\in V} P_\sigma$ is Π^0_1 subset of $2^{\mathbb{N}}$. Moreover, $\blacktriangledown_{\sigma\in V} P_\sigma$ is $(1,1)$-equivalent to $\blacktriangledown_{\sigma\in(V,E(V))} P_\sigma$ in the sense of Definition 79.*

Proof. To simplify our proof, we assume that T_σ is the full binary tree $2^{<\mathbb{N}}$ for each $\sigma \notin V$. Each $\alpha \in 2^{<\mathbb{N}}$ is represented as $\alpha = \rho_0{}^\frown\langle\tau(0)\rangle^\frown\rho_1{}^\frown\langle\tau(1)\rangle^\frown\dots^\frown\langle\tau(|\tau|-2)\rangle^\frown\rho_{|\tau|-1}{}^\frown\langle\tau(|\tau|-1)\rangle^\frown\beta \in 2^{<\mathbb{N}}$, where $\tau \in 2^{<\mathbb{N}}$, $\rho(i) \in T_{\tau\upharpoonright i}$ for each $i < |\tau|$, and $\beta \in T_\tau$. For such τ and β, we set $\mathtt{walk}(\alpha) = \tau$, and $\mathtt{cut}(\alpha) = \langle\rho_0, \rho_1, \dots, \rho_{|\tau|-1}, \beta\rangle$. We also define $\mathtt{tail}^{\mathtt{cut}}(\alpha) = \mathtt{cut}(\alpha; |\mathtt{walk}(\alpha)|) = \beta$. Note that $\mathtt{walk}$, $\mathtt{cut}$, and $\mathtt{tail}^{\mathtt{cut}}$ are total computable on $\mathbb{N}^{<\mathbb{N}}$. Then the set $\blacktriangledown_{\sigma\in V} P_\sigma$ is characterized as the set of all paths of the computable tree $\{\alpha \in 2^{<\mathbb{N}} : \mathtt{walk}(\alpha) \in V \ \&\ \mathtt{tail}^{\mathtt{cut}}(\alpha) \in T_{P_{\mathtt{walk}(\alpha)}}\}$. Therefore, it is Π^0_1. As we mentioned before, we now note that each $\alpha \in 2^{<\mathbb{N}}$ is represented by $\left(\bigsqcap_{i<|\mathtt{walk}(\alpha)|} \mathtt{cut}(\alpha; i)^\frown\langle\mathtt{walk}(\alpha; i)\rangle\right)^\frown\mathtt{cut}(\alpha; |\mathtt{walk}(\alpha)|)$. Then, by defining $\Phi(\alpha)$ for each $\alpha \in 2^{<\mathbb{N}}$ to be the set $\bigsqcap_{i\leq|\mathtt{walk}(\alpha)|} \mathtt{write}(\mathtt{walk}(\alpha) \upharpoonright i, \mathtt{cut}(\alpha; i))$, the function Φ witnesses $\blacktriangledown_{\sigma\in V} P_\sigma \geq^1_1 \blacktriangledown_{\sigma\in(V,E(V))} P_\sigma$.

Conversely, to see $\blacktriangledown_{\sigma\in V} P_\sigma \leq^1_1 \blacktriangledown_{\sigma\in(V,E(V))} P_\sigma$, we inductively define a computable function Ξ. Set $\Phi(\langle\rangle)$. Fix $\alpha = \alpha^{--\frown}\langle(\sigma, m), (\tau, n)\rangle \in (V \times 2)^{<\mathbb{N}}$, and assume that $\Phi(\alpha^-)$ has been already defined. If $\sigma = \tau$, then set $\Xi(\alpha) = \Xi(\alpha^-)^\frown\langle n\rangle$. If $\sigma \neq \tau$, say $\tau = \sigma^\frown\langle i\rangle$, then we first calculate the least leaf $\mathtt{leaf}(\Xi(\alpha^-))$ of T_{P_σ} extending $\Xi(\alpha^-)$. Then we set $\Xi(\alpha) = \mathtt{leaf}(\Xi(\alpha^-)^\frown\langle i, n\rangle$. Note that, for each $\alpha = \alpha^{-\frown}\langle(\tau, n)\rangle \in (V \times 2)^{<\mathbb{N}}$, we have $\mathtt{walk}(\Xi(\alpha)) = (\alpha(|\alpha| - 1))_0 = \tau$, and $\mathtt{tail}^{\mathtt{cut}}(\Xi(\alpha)) = \mathtt{pr}_{\mathtt{walk}(\alpha)}(\alpha)$. Thus, Ξ witnesses $\blacktriangledown_{\sigma\in V} P_\sigma \leq^1_1 \blacktriangledown_{\sigma\in(V,E(V))} P_\sigma$. □

Proposition 122. *Assume that V is a computable subtree of $\mathbb{N}^{<\mathbb{N}}$ and $\{P_\sigma\}_{\sigma\in V}$ is a computable collection of Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$. Then $\blacktriangledown_{\sigma\in V} P_\sigma \equiv^{<\omega}_\omega [V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$.*

Proof. Let $\mathtt{lf}(\sigma)$ be the least leaf of T_{P_σ}. Then $\mathtt{lf} : V \to \mathbb{N}^{<\mathbb{N}}$ is computable, by uniform computability of L_{P_σ}. Fix $f \in [V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$. If $f = 0^\frown g$, then put $\Phi(f) = \bigsqcap_{n\in\mathbb{N}}(\mathtt{lf}(g \upharpoonright n)^\frown\langle g(n)\rangle)$. If $f = \langle 1, \sigma\rangle^\frown g$, then $\Phi(f) = \left(\bigsqcap_{n<|\sigma|} \mathtt{lf}(\sigma \upharpoonright n)^\frown\langle\sigma(n)\rangle\right)^\frown g$. We again use the notions $\mathtt{walk}$ and $\mathtt{tail}^{\mathtt{cut}}$ in the proof of Proposition 121 In the case $f = 0^\frown g$, we have $g \in [V]$. Therefore, $\mathtt{walk}(\Phi(f)) \in [V]$. Hence, $\Phi(f) \in \blacktriangledown_{\sigma\in V} P_\sigma$. If $f = \langle 1, \sigma\rangle^\frown g$, then we have $g \in P_\sigma$. Therefore, $\mathtt{tail}^{\mathtt{cut}}(\Phi(f))$ exists, and $\mathtt{tail}^{\mathtt{cut}}(\Phi(f)) \in P_\sigma$, where $\mathtt{tail}^{\mathtt{cut}}(h) = \lim_{n\in\mathbb{N}} \mathrm{tail}(h \upharpoonright n)$. Thus, $\blacktriangledown_{\sigma\in V} P_\sigma \leq^1_1 [V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$ via Φ.

We consider two learners Ψ_0 who guesses an index of $0^\frown\mathtt{walk}(\alpha)$ on $\alpha \in \mathbb{N}^{<\mathbb{N}}$, and Ψ_1 who guesses an index of $\langle 1, \mathtt{walk}(\alpha)\rangle^\frown\mathtt{tail}^{\mathtt{cut}}(\alpha)$ on $\alpha \in \mathbb{N}^{<\mathbb{N}}$. Note that $\mathtt{walk} : f \mapsto \lim_n \mathtt{walk}(f \upharpoonright n)$ is partial computable, and $\mathtt{tail}^{\mathtt{cut}}$ is learnable. For each $f \in \blacktriangledown_{\sigma\in V} P_\sigma$, if $\mathtt{walk}(f)$ is partial, then $\mathtt{walk}(f) \in V$, and $\mathtt{tail}^{\mathtt{cut}}(f) \in P_{\mathtt{walk}}(f)$. Thus, the learner Ψ_1 succeeds to guess a correct index of a solution $\langle 1, \mathtt{walk}(f)\rangle^\frown\mathtt{tail}^{\mathtt{cut}}(f)$ to $[V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$ on f. If $\mathtt{walk}(f)$ is total, then $\mathtt{walk}(f) \in [V]$. Therefore, the learner Ψ_0 succeeds to guess a correct index of a solution $0^\frown\mathtt{walk}(f)$ to $[V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$ on f. Consequently, $\blacktriangledown_{\sigma\in V} P_\sigma \geq^{<\omega}_\omega [V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma$. □

Definition 123. For Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$, the *hyperconcatenation of P and Q* is defined by $Q \blacktriangledown P = \blacktriangledown_{\sigma\in T_Q} P_\sigma$, where T_Q denotes the corresponding tree for Q, and $P_\sigma = P$ for any $\sigma \in T_Q$.

Proposition 124. *$Q\triangledown_\infty \bigtriangledown P \leq^1_1 Q\blacktriangledown P \leq^1_1 Q\triangledown P \leq^1_1 Q \oplus P$ for any Π^0_1 sets $P, Q \subseteq \mathbb{N}^\mathbb{N}$.*

Proof. We first construct a computable function Φ witnessing $Q\triangledown_\infty \bigtriangledown P \leq^1_1 Q\blacktriangledown P$. Each $\alpha \in \mathbb{N}^{<\mathbb{N}}$ is represented as $\left(\bigoplus_{i<|\mathtt{walk}(\alpha)|} \mathtt{cut}(\alpha; i)^\frown\langle\mathtt{walk}(\alpha; i)\rangle\right)^\frown\mathtt{tail}^{\mathtt{cut}}(\alpha)$, where $\mathtt{cut}(\alpha; i) \in L_P$ for each $i < |\tau|$, $\mathtt{walk}(\alpha) \in \mathbb{N}^{<\mathbb{N}}$, and $\mathtt{tail}^{\mathtt{cut}}(\alpha) \in T_P$. Then we define Φ as follows.

$$\Phi(\alpha) = \left(\prod_{i<|\mathtt{walk}(\alpha)|} \mathtt{write}(1, \mathtt{cut}(\alpha; i))^\frown\langle(0, \mathtt{walk}(\alpha; i))\rangle\right)^\frown\mathtt{write}(1, \mathtt{tail}^{\mathtt{cut}}(\alpha)).$$

Note that $\mathtt{pr}_0(\Phi(\alpha)) = \mathtt{walk}(\alpha)$ and $\mathtt{pr}_1(\Phi(\alpha)) = \left(\prod_{i<|\tau|} \rho_i\right)^\frown\mathtt{tail}(\alpha)$. Therefore, for any $g \in Q\blacktriangledown P$, if $\mathtt{walk}(g)$ is total, then $\mathtt{pr}_0(\Phi(g)) = \mathtt{walk}(g) \in Q$. If $\mathtt{walk}(g)$ is partial, then $\mathtt{tail}^{\mathtt{cut}}(g)$ is total, and $\mathtt{pr}_1(\Phi(g))$ is of the form $\rho_0{}^\frown \dots {}^\frown\rho_n{}^\frown\mathtt{tail}(g)$, where $\rho_i \in L_P$ for each $i \leq n$. Thus, $\mathtt{pr}_1(\Phi(g)) \in \bigtriangledown P$, since $\mathtt{tail}^{\mathtt{cut}}(g) \in P$. Hence, $Q\triangledown_\infty \bigtriangledown P \leq^1_1 Q\blacktriangledown P$.

We next show $Q\blacktriangledown P \leq^1_1 Q\triangledown P$. Fix a leaf $\rho \in L_P$. Each $\sigma \in \mathbb{N}^\mathbb{N}$, if σ extends a leaf $\tau \in L_Q$, say $\sigma = \tau^\frown\eta$, then we set $\Phi(\sigma) = \left(\prod_{i<|\tau|} \rho^\frown\langle\tau(i)\rangle\right)^\frown\eta$. If $\sigma \in T_Q$, then set $\Phi(\sigma) = \left(\prod_{i<|\sigma|} \rho^\frown\langle\sigma(i)\rangle\right)$. Fix $f \in Q\triangledown P$. Then $f \in Q$ or $f = \tau^\frown g$ for some $\tau \in L_Q$ and $g \in P$. If $f \in Q$, then $\mathtt{walk}(\Phi(f)) = f \in Q$. If f is of the form $f = \tau^\frown g$, then $\mathtt{tail}^{\mathtt{cut}}(\Phi(f)) = g \in P$. Hence, $\Phi(f) \in Q\blacktriangledown P$. □

Remark. The relation $Q\triangledown_\infty \bigtriangledown P \leq^1_1 Q\blacktriangledown P$ in Proposition 124 indicates that the hyperconcatenation $Q\blacktriangledown P$ can be viewed as a special case of team-learning proof model of $Q \vee P$, because $\bigtriangledown Q\triangledown_\infty \bigtriangledown P$ is a proof model of $Q \vee P$ by a team of two learners. Later, we will see that $Q\blacktriangledown P$ is also related to Σ^0_2-LEM, the *law of excluded middle for Σ^0_2 formulas*. See Theorem 161 and 181.

Corollary 125. *$[\bigtriangledown_\infty]_{i<2} \bigtriangledown P \leq^1_1 P\blacktriangledown P$ for any Π^0_1 set $P \subseteq \mathbb{N}^\mathbb{N}$.*

Proof. Because $P\triangledown_\infty \bigtriangledown P \subseteq [\bigtriangledown_\infty]_{i<2} \bigtriangledown P$ and by Proposition 124. □

The next theorem concerning the hyperconcatenation $\blacktriangledown$ and the $(1, \omega)$-reducibility $\leq^1_\omega$ is a counterpart of Theorem 83 concerning the concatenation $\triangledown$ and the $(1, 1)$-reducibility $\leq^1_1$.

Theorem 126. *For every special Π^0_1 sets $P, Q \subseteq 2^\mathbb{N}$, and for any R, if $P \leq^1_\omega (Q\blacktriangledown P) \otimes R$ then $P \leq^1_\omega R$ holds.*

Proof. Let $T_\blacktriangledown$ denote the corresponding computable tree for $Q\blacktriangledown P$. *The heart of $T_\blacktriangledown$, $T^\heartsuit_\blacktriangledown$,* is the set of all strings $\gamma \in T_\blacktriangledown$ such that $\gamma \subseteq \prod_{i<n}(\sigma_i{}^\frown\langle\tau(i)\rangle)$ for some $\{\sigma_i\}_{i<n} \subseteq L_P$, and $\tau \in T^{ext}_Q$. Now we assume $P \leq^1_\omega (Q\blacktriangledown P) \otimes R$ via a learner Ψ. To show the theorem it is needed to construct a new learner Δ witnessing $P \leq^1_\omega R$. Fix $g \in R$.

Lemma 127. *There exists a string $\rho \in T^\heartsuit_\blacktriangledown$ such that, for every $\tau \in T^\heartsuit_\blacktriangledown$ extending ρ, we have $\Psi(\rho \oplus (g \upharpoonright |\rho|)) = \Psi(\gamma)$ for any γ with $\rho \oplus (g \upharpoonright |\rho|) \subseteq \gamma \subseteq \tau \oplus (g \upharpoonright |\tau|)$.*

Proof. If Lemma 127 is false, we can inductively define an increasing sequence $\{\tau_i\}_{i\in\omega}$ of strings. First let $\tau_0 = \langle\rangle$, and τ_{i+1} be the least $\tau \supsetneq \tau_i$ such that $\tau \in T^\heartsuit_\blacktriangledown$ and $\Psi(\tau \oplus (g \upharpoonright (|\tau| + i))) \neq \Psi(\rho \oplus (g \upharpoonright (|\rho| + j)))$ for some $i, j < 2$. Since $\bigcup_i \tau_i \in Q\blacktriangledown P$, clearly $(\bigcup_i \tau_i) \oplus g \in (Q\blacktriangledown P) \otimes R$. However, based on the observation $(\bigcup_i \tau_i) \oplus g$, the learner Ψ changes his mind infinitely often. This means that his prediction $\lim_n \Psi((\bigcup_i \tau_i) \oplus g)$ diverges. This contradicts our assumption that $P \leq^1_\omega (Q\blacktriangledown P) \otimes R$ via the learner Ψ. Thus, our claim is verified. □

Lemma 127 can be seen as an analogy of an observation of Blum-Blum [9] in the theory of inductive inference for total computable functions on $\mathbb{N}$. Such ρ is sometimes called *a locking sequence*.

Lemma 128. *There exist an effective procedure $\Theta : \mathbb{N}^\mathbb{N} \times 2^{<\mathbb{N}} \times 2 \to \mathbb{N}^\mathbb{N}$ and a Π^0_1 condition φ such that, for any $g \in Q$, $\varphi(g, \rho, m)$ holds for some $\rho \in 2^{<\mathbb{N}}$, and $m < 2$, and that for any $\rho \in 2^{<\mathbb{N}}$ and $m \in \mathbb{N}$, if $\varphi(g, \rho, m)$ holds, then $\Theta(g, \rho, m) \in P$.*

Proof. The desired condition $\varphi(g,\rho,m)$ is given by the conjunction of the following three conditions.

1. ρ is of the form $\bigsqcap_{i<n}(\sigma_i{}^\frown\langle\tau(i)\rangle)$ for some $\{\sigma_i\}_{i<n}\subseteq L_P$, $\tau\in T_Q^{ext}$, and $|\tau|=n$.
2. $\tau^\frown\langle m\rangle\in T_Q^{ext}$.
3. $\Psi(\rho\oplus(g\upharpoonright|\rho|))=\Psi(\gamma)$ for any $\gamma\in(\rho^\frown T_P{}^\frown\langle m\rangle^\frown T_P)\otimes\{g\}$.

The first two conditions are clearly Π^0_1, and since Ψ is total computable, the last condition is also Π^0_1. Consequently, φ is Π^0_1. We first show that $\varphi(g,\rho,m)$ holds for some $\rho\in 2^{<\mathbb{N}}$ and $m\in\mathbb{N}$. Pick a locking sequence $\rho\in T^{\heartsuit}_{\blacktriangledown}$ forcing to stop changing the mind of Ψ, as in the previous claim. Without loss of generality, we can assume that ρ satisfies the condition (1). Since $\tau\in T_Q^{ext}$, there exists $m\in\omega$ such that $\tau^\frown\langle m\rangle\in T_Q^{ext}$, and this m satisfies the condition (2). From conditions (1) and (2), we conclude that $\rho^\frown P^\frown\langle m\rangle^\frown P=(\rho^\frown P)\cup(\rho^\frown\bigcup_{\sigma\in L_P}\sigma^\frown\langle m\rangle^\frown P)\subseteq T^{\heartsuit}_{\blacktriangledown}$, and so condition (3) is satisfied. Since we assume that $P\leq^1_\omega(Q\blacktriangledown P)\otimes\{g\}$ via the learner Ψ, if $\varphi(g,\rho,m)$ is satisfied, then the following holds.

$$P\leq^1_1(\rho^\frown P^\frown\langle m\rangle^\frown P)\otimes\{g\}\text{ via }\Phi_{\Psi(g\upharpoonright|\rho|\oplus\rho)}.$$

Our proof process in Theorem 83 is effective with respect to g, m, and an index of $\Phi_{\Psi(g\upharpoonright|\rho|\oplus\rho)}$ which are calculated from g, ρ, and an index of Ψ. To see this, recall our proof in Theorem 83. Define $V_P^m=T_P\cup\{\rho^\frown\langle m\rangle:\rho\in L_P\}$.

$$E_0^{g,\rho,m}=V_P^m\otimes\{g\};\qquad D_0^{g,\rho,m}=\Phi_{\Psi(g\upharpoonright|\rho|\oplus\rho)}(E_0^{g,\rho,m}).$$
$$E_{i+1}^{g,\rho,m}=(V_P^{m\frown}D_i^{g,\rho,m})\otimes\{g\};\qquad D_{i+1}^{g,\rho,m}=\Phi_{\Psi(g\upharpoonright|\rho|\oplus\rho)}(E_{i+1}^{g,\rho,m}).$$

Then, as in the proof of Theorem 83, $D^{g,\rho,m}=\bigcup_{i\in\mathbb{N}}D_{i+1}^{g,\rho,m}$ is a subtree of V_P, and it has no dead ends. Moreover, this construction is clearly c.e. uniformly in g, ρ, and m. Therefore, we can effectively choose an element $\Theta(g,\rho,m)\in[D^{g,\rho,m}]\subseteq P$, uniformly in g, ρ, and m. □

Now, a procedure to get $P\leq^1_\omega R$ is follows. For given $g\in Q$, on *the i-th challenge* of a learner Δ, the learner Δ chooses the lexicographically i-th least pair $\langle\rho,m\rangle\in 2^{<\mathbb{N}}\times\mathbb{N}$, and Δ calculates an index $e(\rho,m)$ of the computable functional $g\mapsto\Theta(g,\rho,m)$, that is to say, $\Delta(g\upharpoonright s)=e(\rho,m)$ at the current stage s. At each stage in the i-th challenge, the learner Δ tests whether the Π^0_1 condition $\varphi(g,\rho,m)$ is refuted. When $\varphi(g,\rho,m)$ is refuted, Δ changes his mind, and goes to the $(i+1)$-th challenge. Clearly $\lim_s\Delta(g\upharpoonright s)$ converges, and $\Phi_{\lim_s\Delta(g\upharpoonright s)}(g)\in P$ holds. □

Corollary 129. *For every special Π^0_1 set $P\subseteq 2^{\mathbb{N}}$ there exists a Π^0_1 set $Q\subseteq 2^{\mathbb{N}}$ with $Q<^1_\omega P\equiv^{<\omega}_\omega Q$.*

Proof. By Theorem 126, if $P\leq^1_\omega(P\blacktriangledown P)\otimes 2^{\mathbb{N}}\equiv^1_1 P\blacktriangledown P$, then $P\leq^1_\omega 2^{\mathbb{N}}$, i.e., P contains a computable element. As P is special, we must have $P\not\leq^1_\omega P\blacktriangledown P$. By Proposition 122, $P\leq^{<\omega}_\omega P\blacktriangledown P$. Therefore, for $Q=P\blacktriangledown P$, we have $Q<^1_\omega P\equiv^{<\omega}_\omega Q$. □

Corollary 130. *Every nonzero $\mathbf{a}\in\mathcal{P}^1_\omega$ has the strong anticupping property.*

Proof. Fix $P\in\mathbf{a}$. Let $\mathbf{b}$ be the $(1,\omega)$-degree of $P\blacktriangledown P$. Then, by Theorem 126, for any $(1,\omega)$-degree $\mathbf{c}$, if $\mathbf{a}\leq\mathbf{b}\vee\mathbf{c}$, then $\mathbf{a}\leq\mathbf{c}$. □

The primary motivation of the second author behind introducing the notions of learnability reduction was to attack an open problem on Π^0_1 subsets of $2^{\mathbb{N}}$. The problem (see Simpson [61]) is whether the Muchnik degrees ($(\omega,1)$-degrees) of Π^0_1 classes are dense. Cenzer-Hinman [20] showed that the Medvedev degrees ($(1,1)$-degrees) of Π^0_1 classes are dense. One can easily apply their priority construction to prove densities of $(1,<\omega)$-degrees and $(<\omega,1)$-degrees. The reason is that the arithmetical complexity of $A^\alpha_\beta=\{(i,j)\in\mathbb{N}^2:P_i\leq^\alpha_\beta P_j\}$ is Σ^0_3 for $(\alpha,\beta)\in\{(1,1),(1,<\omega),(<\omega,1)\}$, where $\{P_e\}_{e\in\mathbb{N}}$ is an effective enumeration of all Π^0_1 subsets of $2^{\mathbb{N}}$. It enables us to use priority argument directly. However, for other reductions (α,β), the complexity of A^α_β seems to be Π^1_1. This observation hinders us from using priority arguments. Hence it seems to be a hard task to prove densities of such (α,β)-degrees. Nevertheless, our disjunctive notions turn out to be useful to obtain some partial results.

Theorem 131 (Weak Density)**.** *For nonempty Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$, if $P <^1_\omega Q$ and $P <^{<\omega}_\omega Q$ then there exists a Π^0_1 set $R \subseteq 2^{\mathbb{N}}$ such that $P <^1_\omega R <^1_\omega Q$.*

Proof. Assume $P <^1_\omega Q$ and $P <^{<\omega}_\omega Q$. Let $R = (Q \blacktriangledown Q) \otimes P$. Then $P \leq^1_\omega R \leq^1_\omega Q$. Moreover $Q \not\leq^1_\omega P$ implies $Q \not\leq^1_\omega R = (Q \blacktriangledown Q) \otimes P$, by non-cupping property of $\blacktriangledown$. On the other hand, $R = (Q \blacktriangledown Q) \otimes P \not\leq^1_\omega P$ since $Q \blacktriangledown Q \equiv^{<\omega}_\omega Q \not\leq^{<\omega}_\omega P$. Consequently, $P <^1_\omega R = (Q \blacktriangledown Q) \otimes P <^1_\omega Q$. □

4.7. **Infinitary Disjunctions along Nested ill-Founded Trees.** By using the previous method, we can obtain a finer observation. We introduce two totality restrictions for learners.

Definition 132. Let Ψ be a learner.

(1) Ψ is *confident* if $\lim_s \Psi(f \upharpoonright s)$ converges for every $f \in \mathbb{N}^{\mathbb{N}}$.
(2) Ψ is *eventually-Popperian* if, for every $f \in \mathbb{N}^{\mathbb{N}}$, $\Phi_{\lim_s \Psi(f \upharpoonright s)}(f)$ is total whenever $\lim_s \Psi(f \upharpoonright s)$ converges.
(3) Ψ is *eventually-Lipschitz* if there is a constant $c \in \mathbb{N}$ such that, for every $f \in \mathbb{N}^{\mathbb{N}}$, $|\Phi_{\lim_s \Psi(f \upharpoonright s)}(f \upharpoonright l + c)| \geq l$ for any $l \in \mathbb{N}$, whenever $\lim_s \Psi(f \upharpoonright s)$ converges.

Proposition 133.

(1) *For any set $X, Y \subseteq \mathbb{N}^{\mathbb{N}}$, if $X \leq^{<\omega}_{tt,\omega} Y$, then $X \leq^{<\omega}_\omega Y$ via a team of eventually-Popperian learners.*
(2) *For any Σ^0_2 set $S \subseteq 2^{\mathbb{N}}$ and any set $R \subseteq \mathbb{N}^{\mathbb{N}}$, if $R \leq^1_\omega S$, then it can be witnessed by an eventually-Popperian learner. Moreover, if S is Π^0_1, then it can be witnessed by a confident eventually-Popperian learner.*
(3) *For any Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ and any set $Q \subseteq \mathbb{N}^{\mathbb{N}}$, if $P \leq^{<\omega}_1 Q$ then $P \leq^{<\omega}_\omega Q$ by a team of confident learners.*

Proof. (1) Straightforward from the definition.

(2) Fix a computable increasing sequence $\{T_i\}_{i\in\omega}$ of infinite computable trees such that $S = \bigcup_i [T_i]$. By padding, there is a computable function $p : \mathbb{N}^2 \to \mathbb{N}$ such that $\Phi_{p(e,n)}$ corresponds exactly to Φ_e, and $p(e, n+1) > p(e, n)$ for any index e and n. Assume that $R \leq^1_\omega S$ via a learner Ψ. We need to construct a eventually-Popperian learner Δ witnessing $R \leq^1_\omega S$. At each stage s, we define a value of $\Delta(\sigma)$ for each $\sigma \in 2^s$. For a given $\sigma \in 2^s$, we compute $q(\sigma) = \min(\{i < s : (\forall \tau \in 2^s)\ \tau \supseteq \sigma \ \rightarrow\ \tau \in T_i\} \cup \{s\})$, and put $\Delta(\sigma) = p(\Psi(\sigma), q(\sigma))$. If $f \notin S$, then $\lim_n q(f \upharpoonright n)$ diverges. Therefore, $\lim_n \Delta(f \upharpoonright n)$ diverges. On the other hand, if $f \in S$, then $\lim_n q(f \upharpoonright n)$ converges to some q. Then $\Phi_{\lim_n \Delta(f \upharpoonright n)}(f) = \Phi_{p(\lim_n \Psi(f \upharpoonright n), q)}(f) = \Phi_{\lim_n \Psi(f \upharpoonright n)}(f) \in R$. Consequently, Δ is eventually-Popperian, and witnesses $R \leq^1_\omega S$. If S is Π^0_1, then we modify Δ by setting $\Delta(\sigma)$ to be a fixed index of a total computable function $g \mapsto 0^\omega$, whenever σ extends a leaf of T_S. Then, Δ is also confident.

(3) If $P \leq^{<\omega}_1 Q$ via n many computable functions $\{\Phi_i\}_{i<n}$, then each learner Ψ_i for each $i < n$ guesses an index of Φ_i. Note that Ψ_i does not change his mind. In particular, Ψ_i is confident. □

Proposition 134. *Let V be a computable subtree of $\mathbb{N}^{<\mathbb{N}}$, and $\{P_\sigma\}_{\sigma\in V}$ be a computable collection of Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$. Then $[V] \oplus \bigoplus_{\sigma\in\mathbb{N}^{<\mathbb{N}}} P_\sigma \leq^{<\omega}_\omega \blacktriangledown_{\sigma\in V} P_\sigma$ by a team of a confident learner and an eventually-Popperian learner. Moreover, if P and Q are Π^0_1 subsets of $2^{\mathbb{N}}$, then $Q \oplus P \leq^{<\omega}_\omega Q \triangledown_\infty \bigtriangledown P$ via a team of a confident learner and an eventually-Popperian learner.*

Proof. As in the proof of Proposition 122, we consider two learners: a learner Ψ_0 who guesses an index of $\alpha \mapsto 0^\frown\mathtt{walk}(\alpha)$, and a learner Ψ_1 who guesses an index of $\alpha \mapsto \langle 1, \mathtt{walk}(\alpha)\rangle^\frown\mathtt{tail}^{\mathtt{cut}}(\alpha)$. As $f \mapsto 0^\frown\mathtt{walk}(f)$ is partial computable, Ψ_0 does not change his mind. In particular, Ψ_0 is confident. On $f \in \mathbb{N}^{\mathbb{N}}$, the learner Ψ_1 changes his mind whenever $\mathtt{walk}(f \upharpoonright n+1)$ properly extends $\mathtt{walk}(f \upharpoonright n)$. If $\lim_{n\in\mathbb{N}} \Psi_1(f \upharpoonright n)$ converges, then $\mathtt{walk}(f)$ must be partial. Thus, $\mathtt{tail}^{\mathtt{cut}}(f)$ must be total. Then, $\langle 1, \mathtt{walk}(f)\rangle^\frown\mathtt{tail}^{\mathtt{cut}}(f)$ is total. Therefore, Ψ_1 is eventually-Popperian. Moreover, $Q \oplus P \leq^{<\omega}_\omega Q \triangledown_\infty \bigtriangledown P$ via a team of two learners, $\{\Psi_i\}_{i<2}$, where Ψ_0 guesses $\sigma \mapsto \mathtt{pr}_0(\sigma)$, and Ψ_1 guesses $\sigma \mapsto \mathtt{tail}^{\mathtt{cut}}(\mathtt{pr}_1(\sigma))$. As in the previous argument, we can easily see that Ψ_0 is confident and Ψ_1 is eventually-Popperian. □

Proposition 135. *Let* P_0, P_1, Q_0, Q_1 *be* Π^0_1 *subsets of* $2^{\mathbb{N}}$ *such that* $Q_0 \leq^1_\omega Q_1$ *via an eventually Lipschitz learner and that* $P_0 \leq^1_1 P_1$*. Then,* $Q_0 \blacktriangledown P_0 \leq^1_\omega Q_1 \blacktriangledown P_1$*.*

Proof. For any partial computable function Φ, without loss of generality, we may assume $|\Phi(\sigma)| \leq |\Phi(\sigma^-)| + 1$ for any string $\sigma \in \mathbb{N}^{<\mathbb{N}}$. For given indices i and j, we effectively construct a computable function $\Phi_{\mathtt{hyp}(i,j)}$ as follows. Put $\Phi_{\mathtt{hyp}(i,j)}(\langle\rangle) = \langle\rangle$, and assume that $\Phi_{\mathtt{hyp}(i,j)}(\sigma^-)$ has been already defined. Note that, either $|\mathtt{walk}(\sigma)| = |\mathtt{walk}(\sigma^-)| + 1$ or $|\mathtt{tail}^{\mathtt{cut}}(\sigma)| = |\mathtt{tail}^{\mathtt{cut}}(\sigma^-)| + 1$ is satisfied. Here, the notation $\mathtt{tail}^{\mathtt{cut}}$ is used in referring to decomposing $Q_1 \blacktriangledown P_1$. If the former is the case (i.e., $|\mathtt{walk}(\sigma)| = |\mathtt{walk}(\sigma^-)| + 1$), then we extend $\mathtt{tail}^{\mathtt{cut}}(\Phi_{\mathtt{hyp}(i,j)}(\sigma^-))$ to $\mathtt{leaf} \circ \mathtt{tail}^{\mathtt{cut}}(\Phi_{\mathtt{hyp}(i,j)}(\sigma^-))$, the least leaf of T_{P_0} extending it, and then, concatenate the bit $\Phi_i(\mathtt{walk}(\sigma); |\mathtt{walk}(\sigma)| - c)$ to it. Formally, for a string $\tau \in \mathbb{N}^{<\mathbb{N}}$ with $\Phi_{\mathtt{hyp}(i,j)}(\sigma^-) = \tau^\frown \mathtt{tail}^{\mathtt{cut}}(\Phi_{\mathtt{hyp}(i,j)}(\sigma^-))$, we define

$$\Phi_{\mathtt{hyp}(i,j)}(\sigma) = \tau^\frown \mathtt{leaf} \circ \mathtt{tail}^{\mathtt{cut}}(\Phi_{\mathtt{hyp}(i,j)}(\sigma^-))^\frown \langle \Phi_i(\mathtt{walk}(\sigma); |\mathtt{walk}(\sigma)| - c)\rangle.$$

Here, we fix some string $\rho \in T_{Q_0}$ of length c, and we set $\Phi_i(\sigma; k - c) = \sigma(k)$ for each $k < c$. If $\Phi_i(\mathtt{walk}(\sigma); |\mathtt{walk}(\sigma)| - c)$ is undefined, then $\Phi_{\mathtt{hyp}(i,j)}(\tau)$ is undefined for any $\tau \supseteq \sigma$. If the former is not the case (then, $|\mathtt{tail}^{\mathtt{cut}}(\sigma)| = |\mathtt{tail}^{\mathtt{cut}}(\sigma^-)| + 1$), then we concatenate the new values of $\Phi_j(\mathtt{tail}^{\mathtt{cut}}(\sigma))$ to $\Phi_{\mathtt{hyp}(i,j)}(\sigma^-)$ if it belongs to T_{P_0}. Formally, if $\Phi_j(\mathtt{tail}^{\mathtt{cut}}(\sigma^-)) \subsetneq \Phi_j(\mathtt{tail}^{\mathtt{cut}}(\sigma)) \in T_{P_0}$, say $\Phi_j(\mathtt{tail}^{\mathtt{cut}}(\sigma)) = \Phi_j(\mathtt{tail}^{\mathtt{cut}}(\sigma^-))^\frown \rho$, then we define $\Phi_{\mathtt{hyp}(i,j)}(\sigma) = \Phi_{\mathtt{hyp}(i,j)}(\sigma^-)^\frown \rho$. Otherwise, we set $\Phi_{\mathtt{hyp}(i,j)}(\sigma) = \Phi_{\mathtt{hyp}(i,j)}(\sigma^-)$.

Now assume that $P_0 \leq^1_\omega P_1$ via a computable function Φ_e, and $Q_0 \leq^\omega Q_1$ via an eventually Lipschitz learner Ψ with a constant c. We construct a learner Δ witnessing $Q_0 \blacktriangledown P_0 \leq^1_\omega Q_1 \blacktriangledown P_1$. At first the learner Δ guesses the index $\Delta(\langle\rangle) = \mathtt{hyp}(\Psi(\langle\rangle), e)$. Fix $\sigma \in \mathbb{N}^{<\mathbb{N}}$, and assume that $\Delta(\sigma^-)$ has been already defined. If $\Psi(\mathtt{walk}(\sigma)) \neq \Psi(\mathtt{walk}(\sigma^-))$, then Δ also changes his mind as $\Delta(\langle\rangle\rangle = \mathtt{hyp}(\Psi(\mathtt{walk}(\sigma)), e)$. Assume not. In the case $|\mathtt{walk}(\sigma)| > |\mathtt{walk}(\sigma^-)|$, if either $|\mathtt{walk}(\sigma)| < c$ or $\mathtt{walk}(\sigma) \notin T^{ext}_{Q_1}$ is witnessed, the learner Δ changes his mind (this situation occurs only finitely often). Otherwise, the learner Δ keeps his previous guess, i.e., $\Delta(\sigma) = \Delta(\sigma^-)$. In this way, it is not hard to see that we may construct a learner Δ witnessing $Q_0 \blacktriangledown P_0 \leq^1_\omega Q_1 \blacktriangledown P_1$. □

We may introduce the iteration of the hyperconcatenation $\blacktriangledown$ along any directed graph (V, E). However, the iteration of $\blacktriangledown$ does not represented by our dynamic proof model. We may introduce another new model called *the nested disjunction model*.

The nested tape model: As an example, first we consider the nested disjunction $T^* = \blacktriangledown_{\sigma \in T^0} \blacktriangledown_{\tau \in T^1_\sigma} [T^2_{\sigma,\tau}]$ along the graph $G = (\{0,1,2\}, \{(0,1),(1,2)\})$ with the initial vertex $\varepsilon = 0$, where $T = \{T^0\} \cup \{T^1_\sigma\}_{\sigma \in \mathbb{N}^{<\mathbb{N}}} \cup \{T^2_{\sigma,\tau}\}_{\langle\sigma,\tau\rangle \in (\mathbb{N}^{<\mathbb{N}})^2}$ is a given collection of subtrees of $\mathbb{N}^{<\mathbb{N}}$. The nested tape model for T^* consists of a collection $\{\Lambda_\square\} \cup \{\Lambda^0\} \cup \{\Lambda^1_\sigma\}_{\sigma \in \mathbb{N}^{<\mathbb{N}}} \cup \{\Lambda^2_{\sigma,\tau}\}_{\langle\sigma,\tau\rangle \in (\mathbb{N}^{<\mathbb{N}})^2}$ of infinite tapes.

Generally, a *nested system* (G, T, Λ) consists of a graph $G = (V, E)$ with the initial vertex ε, a collection $T = \{T^v_\sigma\}_{v \in V, \sigma \in (\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}}$ of (ill-founded) trees, and a collection $\Lambda = \{\Lambda_\square\} \cup \{\Lambda^v_\sigma\}_{v \in V, \sigma \in (\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}}$ of infinite tapes. A verifier Ψ is only allowed to write a letter on tapes which are *available*. Assume that a word $\mathtt{pr}[v, \sigma]$ is written on Λ^v_σ for each $v \in V$ and $\sigma \in (\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$. Then, the availability conditions are given as follows.

- $\Lambda_\square$ and $\Lambda^\varepsilon_{\langle\rangle}$ are available at each stage.
- If a finite word $v = \langle v[0], v[1], \dots, v[l]\rangle$ is written on the tape $\Lambda_\square$, then the following tapes are available.

$$\Lambda^{v[1]}_{\mathtt{pr}[v[0],\langle\rangle]}, \Lambda^{v[2]}_{\mathtt{pr}[v[1],\mathtt{pr}[v[0],\langle\rangle]]}, \dots, \Lambda^{v[i]}_{\mathtt{pr}[v[i-1],\mathtt{pr}[v[i-2],\dots,\mathtt{pr}[v[1],\mathtt{pr}[v[0],\langle\rangle]]]]}.$$

Here, on the tape $\Lambda_\square$, the verifier Ψ is only allowed to write a path starting from the initial vertex ε within the graph $G = (V, E)$.

Example 136. On the nested tape model for T^*, let $\alpha \in ((I \cup \{\square\}) \times \mathbb{N})^{<\mathbb{N}}$ be the record of a proof process of Ψ by some stage, i.e., $\mathtt{pr}_\square(\alpha)$ and $\mathtt{pr}_{(v,\sigma)}(\alpha)$, for each $(v, \sigma) \in I^{<\mathbb{N}}$, represent the words written on $\Lambda_\square$


$G:$
0
1
2
$\blacktriangledown_{\sigma\in T^0}T^1_\sigma$
$\blacktriangledown_{\tau\in T^1_\sigma}T^2_{\sigma,\tau}$
T^0
T^1_{101}
$T^2_{101,1001}$
101
1001

FIGURE 3. An example nested tape model when G is a linear order of length 3: $\langle 012\rangle$ is written on $\Lambda_\Box$; $\langle 101\rangle$ is written on Λ^0; $\langle 1001\rangle$ is written on Λ^1_{101}; then $\Lambda_\Box$, Λ^0, Λ^1_{101}, and $\Lambda^2_{101,1001}$ are available.

and Λ^v_σ, respectively. Here, I denotes $V\times(\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$. If the letter 1 representing the vertex $1\in V$ has been written on $\Lambda_\Box$ (i.e., $\mathtt{pr}_\Box(\alpha)\supseteq\langle 01\rangle$), then the three tapes $\Lambda_\Box$, Λ^0, and Λ^1_p are available, where $p=\mathtt{pr}_0(\alpha)$.

The verifier Ψ *succeeds* if he eventually writes a correct solution on some tape from Λ (i.e., some solution $f\in[T^v_\sigma]$ is eventually written on Λ^v_σ for some $(v,\sigma)\in V\times(\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$, or otherwise, some infinite path though G is written on $\Lambda_\Box$). For each $u,v\in V$ and $(v,\sigma)\in V\times(\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$, the tuple $\langle T^v_\sigma,\Lambda^v_\sigma,T^u_{\sigma,\tau},\Lambda^u_{\sigma,\tau}\rangle_{\tau\in T^v_\sigma}$ is called *the* (σ,v,u)*-component of* (G,T,Λ). The (σ,v,u)-component of our nested system consists of an infinite disjunction along an ill-founded tree, $\blacktriangledown_{\tau\in T^v_\sigma}[T^u_{\sigma,\tau}]$. In other words, on the (σ,v,u)-component of the system (I,Λ,T,G), the set Λ^v_σ plays the role of the declaration $\Box$, and $\Lambda^u_{\sigma,\tau}$ plays the role of the working tape for each $\tau\in T^v_\sigma$, as in the dynamic tape model.

Definition 137. Fix a directed graph $G=(V,E)$, and we denotes $V\times(\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$ by I. Assume that a collection $\{T_{(v,\sigma)}\}_{(v,\sigma)\in I}$ of subtrees of $\mathbb{N}^{<\mathbb{N}}$ are given. For $\alpha\in((I\cup\{\Box\})\times\mathbb{N})^{<\mathbb{N}}$, we inductively define *the n-th available index along* α, $p(\alpha,n)\in I$, for each $n\leq|\mathtt{pr}_\Box(\alpha)|$, as follows.

$$p(\alpha,0)=(\varepsilon,\langle\rangle),\qquad p(\alpha,i+1)=(\mathtt{pr}_\Box(\alpha)(i),(p(\alpha,i))_1{}^\frown\langle\mathtt{pr}_{p(\alpha,i)}(\alpha)\rangle).$$

Then we define the set of all indices of *available tapes along* α by $A(\alpha)=\{p(\alpha,n):n\leq|\mathtt{pr}_\Box(\alpha)|\}$. The set $S(\alpha)$ of *successors of* α is defined as follows:

$$\begin{aligned}S(\alpha)=\{(p,n)\in(I\cup\{\Box\})\times\mathbb{N}:p\in A(\alpha)\ \&\ \mathtt{pr}_p(\alpha)^\frown n\in T_p\}\\ \cup\{(\Box,v):(\mathtt{pr}_\Box(\alpha)(|\mathtt{pr}_\Box(\alpha)|-1),v)\in E\}.\end{aligned}$$

Then *the nested infinitary disjunction* $\mathbf{W}_{\sigma\in I}[T_\sigma]\subseteq((I\cup\{\Box\})\times\mathbb{N})^{\mathbb{N}}$ *of* $\{T^v_\sigma\}_{(v,\sigma)\in I}$ is defined by

$$\mathbf{W}_{\sigma\in I}[T_\sigma]=\{f\in((I\cup\{\Box\})\times\mathbb{N})^{\mathbb{N}}:(\forall n\in\mathbb{N})\ f(n)\in S(f\upharpoonright n)\}.$$

We can also define $\mathbb{W}_{\sigma\in I}[T_\sigma]=\{f\in\mathbf{W}_{\sigma\in I}[T_\sigma]:|\mathtt{pr}_\Box(f)|<\infty\}$.

Proposition 138. *Assume that G is a computable directed graph, and $\{T_\sigma\}_{\sigma\in I}$ is a computable collection of computable subtrees of $\mathbb{N}^{\mathbb{N}}$, where $I=V\times(\mathbb{N}^{<\mathbb{N}})^{<\mathbb{N}}$. Then, $\mathbf{W}_{\sigma\in I}[T_\sigma]$ is Π^0_1. Moreover, if G and T_σ are subtrees of $2^{<\mathbb{N}}$ for each $\sigma\in I$, then $\mathbf{W}_{\sigma\in I}[T_\sigma]$ is $(1,1)$-equivalent to a Π^0_1 subset of $2^{\mathbb{N}}$.*

Proof. Note that $\alpha\mapsto A(\alpha)$ is computable. Therefore, $\alpha\mapsto S(\alpha)$ is also computable. Thus, $\mathbf{W}_{\sigma\in I}[T_\sigma]$ is Π^0_1.

Assume that $G=(V,E(V))$ and T_σ are subtrees of $2^{<\mathbb{N}}$ for each $\sigma\in I$. Fix new symbols $+,-$ which does not belong to $\mathbb{N}$. To construct a Π^0_1 subset of $(\{+,-\}\cup 2)^{\mathbb{N}}$ which is $(1,1)$-equivalent to $\mathbf{W}_{\sigma\in I}[T_\sigma]$, we inductively define a computable function head : $(\{+,-\}\cup 2)^{<\mathbb{N}}\to\mathbb{Z}$. Fix $\alpha=\alpha^{-\frown}\langle w\rangle\in(\{+,-\}\cup 2)^{<\mathbb{N}}$. Put head$(\langle\rangle)=0$, Put head$(\alpha)=$ head$(\alpha^-)+1$ if $w=+$; put head$(\alpha)=$ head(α^-) if $w\notin\{+.-\}$; and put

$\mathrm{head}(\alpha) = \mathrm{head}(\alpha^-) - 1$ if $w = -$. If $\alpha = \alpha^{--\frown}\langle +,+\rangle$ and $\mathrm{head}(\alpha) = \max\{\mathrm{head}(\beta) : \beta \subsetneq \alpha\} + 2$, or if $\mathrm{head}(\alpha) = -1$, then we say that α is *overflowing*. If α has an overflowing initial segment $\beta \subseteq \alpha$, then we also say that α is overflowing. Let Rule denote the set of all non-overflowing strings $\alpha \in (\{+,-\} \cup 2)^{<\mathbb{N}}$ which has neither $\langle +,-\rangle$ nor $\langle -,+\rangle$ as substrings. Note that Rule is computable.

We now inductively define $\tilde{\mathtt{pr}}_\square$, $\tilde{p}$, and $\tilde{\mathtt{pr}}_\sigma$ for each $\sigma \in V$. Put $\tilde{\mathtt{pr}}_\square(\langle\rangle)$, and $\tilde{p} = \langle\langle\rangle\rangle$. Fix $\alpha = \alpha^{-\frown}w \in$ Rule. Assume that $\tilde{\mathtt{pr}}_\square(\alpha^-)$, and $\tilde{p}(\alpha^-)$ have been already defined. If $w \in \{+,-\}$, then $\tilde{\mathtt{pr}}_\square(\alpha) = \tilde{\mathtt{pr}}_\square(\alpha^-)$ and $\tilde{p}(\alpha) = \tilde{p}(\alpha^-)$. Assume $w \notin \{+,-\}$. Then, if $\mathrm{head}(\alpha) > \max\{\mathrm{head}(\beta) : \beta \subsetneq \alpha\}$, then we define $\tilde{\mathtt{pr}}_\square(\alpha) = \tilde{\mathtt{pr}}_\square(\alpha^-)^\frown w$. Otherwise, set $\tilde{\mathtt{pr}}_\square(\alpha) = \tilde{\mathtt{pr}}_\square(\alpha^-)$. If $\tilde{\mathtt{pr}}_\square(\alpha) \neq \tilde{\mathtt{pr}}_\square(\alpha^-)$, then $\tilde{p}(\alpha) = \tilde{p}(\alpha^-)^\frown\langle\langle\rangle\rangle$. Otherwise, define $\tilde{p}(\alpha) \in (2^{<\mathbb{N}})^{|V(\alpha)|}$ as follows.

$$(\tilde{p}(\alpha))(n) = \begin{cases} (\tilde{p}(\alpha^-))(n), & \text{if } n < \mathrm{head}(\alpha); \\ (\tilde{p}(\alpha^-))(n)^\frown w, & \text{if } n = \mathrm{head}(\alpha); \\ \langle\rangle, & \text{if } h(\alpha) < n \leq |\tilde{\mathtt{pr}}_\square(\alpha)|. \end{cases}$$

Then, for each $\sigma \in V$, we define $\tilde{\mathtt{pr}}_\sigma(\alpha) = (\tilde{p}(\beta))(|\sigma|)$ for the greatest $\beta \subseteq \alpha$ such that $\sigma \subseteq \tilde{p}(\beta)$. Set $\mathrm{Rule}_\forall = \{f \in (\{+,-\} \cup 2)^{\mathbb{N}} : (\forall n \in \mathbb{N})\ f \upharpoonright n \in \mathrm{Rule}\}$. Note that any $g \in \mathrm{Rule}_\forall$ has no infinite $\{+,-\}$-sequence; otherwise $g \upharpoonright s$ for some $s \in \mathbb{N}$ is overflowing or has a substring $\langle +,-\rangle$ or $\langle -,+\rangle$, and hence $g \upharpoonright s$ must go against Rule. Then P is defined as follows.

$$P = \{f \in \mathrm{Rule}_\forall : (\forall n \in \mathbb{N})\ (\tilde{\mathtt{pr}}_\square(f \upharpoonright n) \in V \ \&\ (\forall \sigma \in I)\ \tilde{\mathtt{pr}}_\sigma(f \upharpoonright n) \in T_\sigma)\}.$$

Clearly, P is computably bounded, and Π^0_1. It remains to show that $P \equiv^1_1 \mathbf{W}_{\sigma \in I}[T_\sigma]$. We first inductively define a computable function Φ witnessing $P \geq^1_1 \mathbf{W}_{\sigma \in I}[T_\sigma]$. Set $\Phi(\langle\rangle) = \langle\rangle$, fix $\alpha = \alpha^{-\frown}w \in$ Rule, and assume that $\Phi(\alpha^-)$ has been already defined. If $w \in \{+.-\}$, then set $\Phi(\alpha) = \Phi(\alpha^-)$. Assume $w \notin \{+,-\}$. If $\mathrm{head}(\alpha) > \max\{\mathrm{head}(\beta) : \beta \subsetneq \alpha\}$, then we set $\Phi(\alpha) = \Phi(\alpha^-)^\frown\langle(\square, w)\rangle$. Otherwise, we set $\Phi(\alpha) = \Phi(\alpha^-)^\frown\langle((\tilde{\mathtt{pr}}_\square(\alpha), \tilde{p}(\alpha) \upharpoonright h(\alpha)), w)\rangle$. It is not hard to check $P \geq^1_1 \mathbf{W}_{\sigma \in I}[T_\sigma]$ via Φ.

To prove $P \geq^1_1 \mathbf{W}_{\sigma \in I}[T_\sigma]$, we first define a computable function head^*. Firstly put $\mathrm{head}^*(\langle\rangle) = 0$. Fix $\alpha = \alpha^{-\frown}\langle(\sigma, w)\rangle \in ((I \cup \{\square\}) \cup \mathbb{N})^{<\mathbb{N}}$. If $\sigma = \square$, then we set $\mathrm{head}^*(\alpha) = |\mathtt{pr}_\square(\alpha)|$. If $\sigma \in I$, then we set $\mathrm{head}^*(\alpha) = |(\sigma)_1|$. Set $\Phi(\langle\rangle) = \langle\rangle$, and assume that $\Phi(\alpha^-)$ has already been defined. Put $d = \mathrm{head}^*(\alpha) - \mathrm{head}^*(\alpha^-)$. If $d \geq 0$, then $\Phi(\alpha) = \Phi(\alpha^-)^\frown +^{d\ \frown} w$. If $d < 0$, then $\Phi(\alpha) = \Phi(\alpha^-)^\frown -^{-d\ \frown} w$. It is not hard to check $P \leq^1_1 \mathbf{W}_{\sigma \in I}[T_\sigma]$ via Φ. □

If T^v_σ only depends on $v \in V$, i.e., $T^v_\sigma = T_v$, then the nested system (I, Λ, T, G) can be viewed as the iteration of the hyperconcatenation $\blacktriangledown$ along the graph G. In this case, we write $\mathbf{W}_{v \in (V,E)} P_v$ for this notion.

Proposition 139. *Let (V, E) be a computable directed graph, and $\{P_v\}_{v \in V}$ be a computable collection of Π^0_1 subsets of $\mathbb{N}^{\mathbb{N}}$. Then, $\mathbf{W}_{v \in (V,E)} P_v \leq^1_1 \blacktriangledown_{v \in (V,E)} P_v$.*

Proof. We inductively define a computable function Φ which witnesses the condition $\mathbf{W}_{v \in (V,E)} P_v \leq^1_1 \blacktriangledown_{v \in (V,E)} P_v$. Set $\Phi(\langle\rangle) = \langle\rangle$. Fix $\alpha = \alpha^{--\frown}\langle(u, i), (v, j)\rangle \in (V \times \mathbb{N})^{<\mathbb{N}}$. Assume that $\Phi(\alpha^-)$ has already been defined, and $\Phi(\alpha^-)$ is of the form $\Phi(\alpha^-) = \beta^\frown\langle(\sigma, k)\rangle$ for some $\beta \in ((I \cup \{\square\}) \times \mathbb{N})^{<\mathbb{N}}$, $\sigma \in I \cup \{\square\}$, and $k \in \mathbb{N}$. If $v = u$, then we set $\Phi(\alpha) = \Phi(\alpha^-)^\frown\langle(\sigma, j)\rangle$. If $v \neq u$, then we set $\Phi(\alpha) = \Phi(\alpha^-)^\frown\langle(\square, v), ((v, (\sigma)_1{}^\frown\mathtt{pr}_u(\alpha)), j)\rangle$. Fix $g \in \blacktriangledown_{v \in (V,E)} P_v$. By induction, we can show $\mathtt{pr}_{v[n]}(g \upharpoonright n + 1) = \mathtt{pr}_{\sigma[n]}(\Phi(g \upharpoonright n + 1))$, where $g(n) = (v[n], j)$ and $\Phi(g \upharpoonright n + 1) = \beta^\frown\langle(\sigma[n], j)\rangle$. Then, $(\sigma[n])_1 = (\sigma[n]^-)_1{}^\frown\mathtt{pr}_{\sigma[n]^-}(\Phi(g \upharpoonright n + 1))$, by our definition of Φ. Therefore, $\sigma[n]$ is available whenever $\sigma[n]^-$ is available. By induction, $\sigma[n]$ is available at $g \upharpoonright n$, for any $n \in \mathbb{N}$. Moreover, $\mathtt{pr}_{\sigma[n]}(\Phi(g)) = \mathtt{pr}_{v[n]}(g) \in T_{v[n]} = T_{\sigma[n]}$, and $\mathtt{pr}_\square(\Phi(g)) = \mathrm{walk}(g)$. Here $\mathrm{walk}(g)$ is inductively defined as follows. Set $\mathrm{walk}(g \upharpoonright 1) = (g(0))_0$. If $(g(n+1))_0 = (g(n))_0$, then $\mathrm{walk}(g \upharpoonright n+1) = \mathrm{walk}(g \upharpoonright n)$. If $(g(n+1))_0 \neq (g(n))_0$, then $\mathrm{walk}(g \upharpoonright n + 1) = \mathrm{walk}(g \upharpoonright n)^\frown(g(n + 1))_0$. Note that $\langle \mathrm{walk}(g; n), \mathrm{walk(g; n + 1)}\rangle \in E$ for each $n < |\mathrm{walk}(g)| - 1$. Thus, $\Phi(g; s) \in S(\Phi(g) \upharpoonright s)$ for any $s \in \mathbb{N}$. Consequently, $\Phi(g) \in \mathbf{W}_{v \in (V,E)} P_v$. □

If $G = (V, E)$ is linearly ordered, then we have no choice of the next vertex at each stage. In this case, to simplify our argument, we assume that only $\{\Lambda_\sigma\}_{\langle v,\sigma\rangle \in I}$ is given, i.e., the (v, σ)-th tape Λ^v_σ does not depend on the vertex $v \in V$, and. Moreover, if $T_\sigma = T_\tau$ for any $\sigma, \tau \in I$, then we only require $\{\Lambda_{|\sigma|}\}_{\langle v,\sigma\rangle \in I}$. We will use the simplest depth n nested system. The system (G, T, Λ) is *an* $\mathbb{N}^{<n}$*-nested system* if $G = (n, S)$ and $T_\sigma = T_\tau$ for any $\sigma, \tau \in I$. This system is equivalent to the n-th iteration of $\blacktriangledown$. Let $P^{\blacktriangledown(0)} = P$, and $P^{\blacktriangledown(n+1)} = P \blacktriangledown P^{\blacktriangledown(n)}$. We also write $\blacktriangledown P$ for $\bigcup_{n \in \mathbb{N}} P^{\blacktriangledown(n)}$.

Proposition 140. *Let* $G = (n+2, S)$*, where* $n + 2 = \{m \in \mathbb{N} : m < n+2\}$ *and* $S = \{(m, m+1) : m \leq n\}$*, and* $\{P^v_\sigma\}_{\langle v,\sigma\rangle \in I}$ *be a computable collection of* Π^0_1 *subsets of* $\mathbb{N}^\mathbb{N}$*. Let* T^v_σ *denote the corresponding tree of* P^v_σ *for each* $\langle v, \sigma\rangle \in I$*. Then* $\mathbf{W}_{\langle v,\sigma\rangle \in I} P^v_\sigma$ *is* $(1, 1)$*-equivalent to the following set:*

$$Q = \blacktriangledown_{\sigma(0) \in T^0_{\langle\rangle}} \left(\blacktriangledown_{\sigma(1) \in T^1_{\sigma(0)}} \left(\dots \left(\blacktriangledown_{\sigma(n) \in T^n_{\sigma(0),\dots,\sigma(n-1)}} P^{n+1}_{\sigma(0),\dots,\sigma(n)} \right) \dots \right) \right).$$

In particular, $\mathbf{W}_{v \in (n,S)} P = P^{\blacktriangledown(n)}$ *for any* Π^0_1 *subset of* $\mathbb{N}^\mathbb{N}$*.*

Proof. Straightforward. □

Proposition 141. *For any* Π^0_1 *set* $P \subseteq 2^\mathbb{N}$*, the following conditions hold.*

1. $[\bigtriangledown_\infty]_{i<n+1} \bigtriangledown P \leq^1_1 P^{\blacktriangledown(n)}$ *for each* $n \in \mathbb{N}$*.*
2. *There is an* $(n+1, \omega)$*-computable function* $P^{\blacktriangledown(n+1)} \to P$ *via a team* $\{\Psi_i\}_{i \leq n}$ *of learners, where* Ψ_0 *is confident, and* Ψ_n *is eventually-Popperian. In particular,* $P \equiv^{<\omega}_\omega \blacktriangledown P$*.*

Proof. The item (1) follows from Corollary 125. To prove the item (2), we use the Π^0_1 subset of $2^\mathbb{N}$ corresponding to $P^{\blacktriangledown(n+1)}$ constructed as in Proposition 138. Then, on each $\alpha \in \mathbb{N}^{<\mathbb{N}}$, the i-th learner Ψ_i guesses an index of $\mathtt{pr}_{\tilde{p}(\alpha)\upharpoonright i}$. Note that $\mathtt{pr}_{\tilde{p}(\alpha)\upharpoonright i} = \mathtt{pr}_{\langle\rangle}$ is partial computable. In particular, Ψ_0 is confident. Fix $g \in P^{\blacktriangledown(n+1)}$. Then, $\lim_s \Psi_i(g \upharpoonright s)$ converges if and only if $\lim_s(\tilde{p}(g \upharpoonright s) \upharpoonright i)$ converges. For the greatest i such that $\lim_s(\tilde{p}(g \upharpoonright s) \upharpoonright i)$ converges to some $\sigma \in (2^{<\mathbb{N}})^{<i}$, we must have $\mathtt{pr}_\sigma(g) \in P$. Thus, if $\lim_s \Psi_n(g \upharpoonright s)$ converges, then $\lim_s \tilde{p}(g \upharpoonright s)$ converges to some $\sigma \in (2^{<\mathbb{N}})^{<n}$, and hence $\Phi_{\lim_s \Psi_n(g \upharpoonright s)}(g) = \mathtt{pr}_\sigma(g) \in P$. In particular, Ψ_n is eventually-Popperian. □

Proposition 142. *Let* P *be a* Π^0_1 *subset of* $\mathbb{N}^\mathbb{N}$*, and* Q *be a subset of* $\mathbb{N}^\mathbb{N}$*. If* $P \leq^{<\omega}_\omega Q$ *by a team of* n *eventually-Popperian learners, then* $P^{\blacktriangledown(n)} \leq^1_\omega Q$*.*

Proof. Assume that $P \leq^{<\omega}_\omega P^{\blacktriangledown(n+1)} \otimes R$ by a team $\{\Psi_i\}_{i<n}$ of n eventually-Popperian learners. For $f \in Q$, assume that the $\mathbb{N}^{<n}$-nested tape $\{\Lambda_\sigma\}_{\sigma \in \mathbb{N}^{<n}}$ is given. Assume that the q-th challenge of Γ is started at stage s_q. At the beginning of the stage s_q, all words on $\{\Lambda_\sigma\}_{\sigma \in \mathbb{N}^{<n}}$ are removed. The learner Γ uses $\{\Lambda_\sigma\}_{\sigma \in \mathbb{N}^{<n}}$ to get an element of $P^{\blacktriangledown(n)}$, at the q-th challenge. The $\Gamma(f \upharpoonright s_q)$-th algorithm $\Phi_{\Gamma(f \upharpoonright s_q)}$ which is guessed by Γ at stage s_q is denoted by Γ^*_q.

- We say that Γ^*_q *writes on* σ *at stage* t if the learner Γ simulates the calculation of $\Phi_{\Psi_{|\sigma|}(f \upharpoonright t)}(f \upharpoonright t)$ on the tape Λ_σ at stage t, i.e., the learner Γ tries to make $\mathtt{pr}_\sigma \circ \Phi_{\Gamma(f \upharpoonright s_q)}(f \upharpoonright t) = \Phi_{\Psi_{|\sigma|}(f \upharpoonright t)}(f \upharpoonright t)$.
- We say that Γ^*_q *is currently-refuted on* σ *at stage* $t+1$ if either $\Psi_{|\sigma|}(f \upharpoonright t+1) \neq \Psi_{|\sigma|}(f \upharpoonright t)$ or $\Phi_{\Psi_{|\sigma|}(f \upharpoonright t+1)}(f \upharpoonright t+1) \notin T_P$.

Now we describe the strategy of Γ^*_q. *The active leaf at the beginning stage* s_q is the leftmost leaf $0^{n-1} \in \mathbb{N}^{<n}$. At each stage $t \geq s_q$ of the q-th trial of the learner Γ, his algorithm Γ^*_q proceeds as follows.

- Γ^*_q writes on $\rho \upharpoonright n-m-1$ for the active leaf ρ at substage m of stage t, and goes to next stage $t+1$, if it is not currently-refuted.
- If Γ^*_q is currently-refuted on $\rho \upharpoonright n-m-1$ for the active leaf ρ at substage m of stage t, then Γ^*_q acts as follows:
  - The word written on the tape $\Lambda_{\rho \upharpoonright n-m-1}$ before stage s_q is extended to a leaf of T_P.
  - The active leaf is switched to $(\rho \upharpoonright n-m-2)^\frown\langle \rho(n-m-2)+1\rangle^\frown\langle 00\dots0\rangle \in \mathbb{N}^{<n}$, i.e., the leftmost leaf of $\mathbb{N}^{<n}$ extending $(\rho \upharpoonright n-m-2)^\frown\langle \rho(n-m-2)+1\rangle$.
  - Go to next substage $m+1$.
- If Γ^*_q is currently-refuted on $\langle\rangle$ at some stage u, then the $(q+1)$-th challenge of Γ begins at stage u.

By the eventually-Popperian assumption, the word written on any tape must be eventually currently-refuted or belong to P, since P is Π^0_1. Therefore, the learner Γ changes his mind at most finitely often. Since we have $P \leq^{<\omega}_{\omega} P^{\blacktriangledown(n+1)} \otimes R$ via $\{\Psi_i\}_{i<n}$, an element of P is written on some tape. Therefore, $P^{\blacktriangledown(n)} \leq^1_\omega P^{\blacktriangledown(n+1)} \otimes R$ is satisfied via the learner Γ, as desired. □

Definition 143. Let P be a Π^0_1 subset of $2^{\mathbb{N}}$. For each $a \in \mathcal{O}$, we define *the a-hyperweakening $P^{\blacktriangledown(a)}$* as follows.

1. $P^{\blacktriangledown(0)} = P$.
2. If $a = 2^b$, then $P^{\blacktriangledown(a)} = P \blacktriangledown P^{\blacktriangledown(b)}$.
3. If $a = 3 \cdot 5^e$, then $P^{\blacktriangledown(a)} = P^\frown \{P^{\blacktriangledown(\Phi_e(n))}\}_{n\in\mathbb{N}}$.

Remark. We can give another definition for $P^{\blacktriangledown(a)}$, in the case that $a = 3 \cdot 5^e$. Let $P \blacktriangledown \{P_n\}_{n\in\mathbb{N}}$ denote $\blacktriangledown_{\sigma \in T_P} P_\sigma$, where put $P_\sigma = P_n$ if σ is the n-th node of T_P. Then $P^{\blacktriangledown(a)}$ can be defined as $P \blacktriangledown \{P^{\blacktriangledown(\Phi_e(n))}\}_{n\in\mathbb{N}}$. However, we will use the previous definition for $P^{\blacktriangledown(a)}$ to simplify our argument.

Proposition 144. *Let $P \subseteq 2^{\mathbb{N}}$ be a Π^0_1 set, and $a \in \mathcal{O}$ be a notation. Then, the a-hyperweakening $P^{\blacktriangledown(a)}$ is a Π^0_1 subset of $2^{\mathbb{N}}$.*

Proof. The proof is straightforward, as in the proof of Proposition 100. □

Proposition 145. *For any special Π^0_1 set $P \subseteq 2^{\mathbb{N}}$, if $a, b \in \mathcal{O}$ and $a <_{\mathcal{O}} b$, then $P^{\blacktriangledown(b)}$ does not $(1,\omega)$-cup to $P^{\blacktriangledown(a)}$, i.e., for any set $R \subseteq \mathbb{N}^{\mathbb{N}}$, if $P^{\blacktriangledown(a)} \leq^1_\omega P^{\blacktriangledown(b)} \otimes R$ then $P^{\blacktriangledown(a)} \leq^1_\omega R$.*

Proof. The assumption $a <_{\mathcal{O}} b$ implies $2^a \leq_{\mathcal{O}} b$. Therefore, we have $P^{\blacktriangledown(b)} \leq^1_\omega P^{\blacktriangledown(2^a)}$. By Theorem 126, $P^{\blacktriangledown(2^a)}$ does not $(1,\omega)$-cup to $P^{\blacktriangledown(a)}$. Thus, $P^{\blacktriangledown(b)}$ does not $(1,\omega)$-cup to $P^{\blacktriangledown(a)}$. □

Fix again any notation $\mathtt{omega} \in \mathcal{O}$ such that $|\Phi_{\mathtt{omega}}(n)|_{\mathcal{O}} = n$ for each $n \in \mathbb{N}$.

Proposition 146. *Let P be a special Π^0_1 subset of $2^{\mathbb{N}}$, and $a \in \mathcal{O}$ be any notation. For any set $R \subseteq \mathbb{N}^{\mathbb{N}}$, if $P^{\blacktriangledown(a)} \leq^{<\omega}_\omega P^{\blacktriangledown(a \cdot_{\mathcal{O}} \mathtt{omega})} \otimes R$ by a team of eventually-Popperian learners, then $P^{\blacktriangledown(a)} \leq^{<\omega}_\omega R$.*

Proof. If $P^{\blacktriangledown(a)} \leq^{<\omega}_\omega P^{(a +_{\mathcal{O}} \mathtt{omega})} \otimes R$ via a team of eventually-Popperian learners, then this reduction is also witnessed by a team of n eventually-Popperian learners, for some $n \in \mathbb{N}$. In particular, by modifying this reduction, we can easily construct a team of n eventually-Popperian learners witnessing $P^{\blacktriangledown(a)} \leq^{<\omega}_\omega P^{\blacktriangledown(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n+1))} \otimes R$. Then, by Proposition 142, we have $P^{\blacktriangledown(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n))} \leq^1_\omega P^{\blacktriangledown(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n+1))} \otimes R = (P \blacktriangledown P^{\blacktriangledown(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n))}) \otimes R$. By Theorem 126, $P^{\blacktriangledown(a \cdot_{\mathcal{O}} \Phi_{\mathtt{omega}}(n))} \leq^1_\omega R$. Hence, $P^{\blacktriangledown(a)} \leq^{<\omega}_\omega R$ is witnessed by a team of n learners, by Proposition 141. □

Corollary 147. *For every $a \in \mathcal{O}$ there exists a computable function g such that, for any Π^0_1 index e, if P_e is special then the following properties hold.*

1. *$P_{g(e,b)} <^1_\omega P_{g(e,c)}$ holds for every $c <_{\mathcal{O}} b <_{\mathcal{O}} a$, indeed, $P_{g(e,b)}$ does not $(1,\omega)$-cup to $P_{g(e,c)}$.*
2. *$P_{g(e,b)} \equiv^\omega_1 P_{g(e,c)}$ for every $b, c <_{\mathcal{O}} a$.*

Proof. Let $g(e,b)$ be an index of $P_e^{\blacktriangledown(b)}$. Then the desired conditions follow from Proposition 145. □

Corollary 148. *For any nonzero $(\omega,1)$-degree $\mathbf{a} \in \mathcal{P}^\omega_1$, there is a $(1,\omega)$-noncupping sequence of $(1,\omega)$-degrees inside $\mathbf{a}$ of length ω^{CK}_1. (Of course, such a sequence is not computable.)*

Proof. Fix a Π^0_1 set $P \in \mathbf{a}$. Pick a path $p \subseteq \mathcal{O}$ through $\mathcal{O}$ of order type ω^{CK}_1. Then, $\{P^{\blacktriangledown(a)}\}_{a\in p}$ is a desired sequence, by Proposition 145. □

Remark. We may introduce the "*nested nested*" model, the "*nested nested nested*" model, and so on. Let $Q\mathbf{w}P$ be $\mathbf{W}_{v \in (T_Q, E(T_Q))} P_v$, where $P_v = P$ for each $v \in T_Q$. Then, for example, the nested nested model can be introduced as the iteration of $\mathbf{w}$ along any directed graph (V, E). Therefore, inside the $(\omega,1)$-degree of any Π^0_1 set $P \subseteq 2^{\mathbb{N}}$, one may iterate this procedure as "*nested nested nested ... nested nested*

. . . ” Actually one may iterate “*nested nested nested . . . nested nested . . .* ” along any directed graph, for example, along the corresponding tree of P. If we call it a “*hypernested*” model, then, of course, we may introduce models which are “*hypernested hypernested*”, and “*hypernested hypernested hypernested*”, and so on. By iterating this notion along the corresponding tree of P, we obtain a “*hyperhypernested*” model. Iterating this procedure, of course, we have the iteration of “*hyper*” along the correspoding tree of P. This observation reveals to us that there are a fine structure, a deep hierarchy, and a morass inside each $(\omega,1)$-degree (or equivalently, each Turing upward closure) of a Π^0_1 subset of $2^{\mathbb{N}}$.

4.8. **Infinitary Disjunctions along Infinite Complete Graphs.**

Theorem 149. *For every special Π^0_1 set $P,Q\subseteq 2^{\mathbb{N}}$ there exists a Π^0_1 set $\widehat{P}\subseteq 2^{\mathbb{N}}$ such that $Q\not\leq^{<\omega}_{\omega}\widehat{P}$ and $\widehat{P}$ is $(\omega,1)$-equivalent to P.*

Proof. We construct a Π^0_1 set $P\subseteq 2^{\mathbb{N}}$ by priority argument with infinitely many requirements $\{\mathcal{P}_e,\mathcal{G}_e\}_{e\in\mathbb{N}}$. Each preservation ($\mathcal{P}_e$-)strategy will injure our coding ($\mathcal{G}$-)strategy of P into $\widehat{P}$ infinitely often. In other words, for each $\mathcal{P}_e$-requirement, $\widehat{P}$ contains an element g^f_e which is a counterpart of each $f\in P$, but each g^f_e has infinitely many noises. Indeed, to satisfy the $\mathcal{P}$-requirements, we need to ensure that there is no uniformly team-learnable way to extract the information of $f\in P$ from its code $g^f_e\in\widehat{P}$. Nevertheless, the global ($\mathcal{G}$-)requirement must guarantee that $f\in P$ is computable in $g^f_e\in\widehat{P}$ via a non-uniform way. Let $\{\Psi^e_i\}_{i<b(e)}$ be the e-th team of learners, where $b=b(e)$ is the number of members of the e-th team.

Requirements. It suffices to construct a Π^0_1 set $\widehat{P}\subseteq 2^{\mathbb{N}}$ satisfying the following requirements.

$$\mathcal{P}_e : (\exists g_e\in\widehat{P})(\forall i<b)\ \left(\lim_s \Psi^e_i(g_e\upharpoonright s)\downarrow\rightarrow\ \Phi_{\lim_s\Psi^e_i(g_e\upharpoonright s)}(g_e)\notin Q\right).$$
$$\mathcal{G}_e : (\forall f\in P)\ f\leq_T g^f_e.$$

Here, the desired Π^0_1 set $\widehat{P}\subseteq 2^{\mathbb{N}}$ will be of form $P\cup\{g^f_e : e\in\mathbb{N}\ \&\ f\in P\}$.

Construction. We will construct a computable sequence of computable trees $\{T_s\}_{s\in\mathbb{N}}$, and a computable sequence of natural numbers $\{h_s\}_{s\in\mathbb{N}}$. The desired set $\widehat{P}$ is defined as $[\bigcup_s T_s]$, and h_s is called *active height at stage s*. We will ensure that the tree T_s consists of strings of length $\leq h_s$. The strategy for the $\mathcal{P}_e$-requirement acts on some string extending the e-th leaf ρ_e of T_{CPA}.

We will inductively define a string $\gamma_e(\alpha,s)\in T_s$ extending ρ_e for each $s\in\mathbb{N}$ and $\alpha\in T_P$ of height $\leq s$. The map $\alpha\mapsto\lim_s\gamma_e(\alpha,s)$ restricted to T^{ext}_P will provide a tree-isomorphism between T^{ext}_P and $(\bigcup_s T_s)^{ext}$, i.e., $\widehat{P}\cap[\rho_e]$ will be constructed as the set of all infinite paths of the tree generated by $\{\lim_s\gamma_e(\alpha,s):\alpha\in T_P\}$. In other words, g^f_e is defined by $\bigcup_{\alpha\subset f}\lim_s\gamma_e(\alpha,s)$, and each string $\gamma_e(\alpha,s)$ is an approximation of $g_e\in\widehat{P}$ witnessing to satisfy the $\mathcal{P}_e$ requirements.

We will also define a finite set $M_e(\alpha,s)\subseteq b$ for each $s\in\mathbb{N}$ and $\alpha\in T_P$ of height $\leq s$. Intuitively, $M_e(\alpha,s)$ contains any index of the learner who have been already changed his mind $|\alpha|$ times along any string extending α of length s, and the string $\gamma_e(\alpha,s)$ also plays the role of an active node for learners in $M_e(\alpha,s)$. To satisfy the $\mathcal{P}_e$-requirement, each learner in $M_e(\alpha,s)$ can act on $\gamma_e(\alpha,s)$ at stage $s+1$, and then he extends $\gamma_e(\alpha,s)$ to some new string $\gamma_e(\alpha,s+1)$ of length h_s, and *injures* all constructions of $\gamma_e(\beta,s+1)$ for $\beta\supsetneq\alpha$. We assume that, for any $\alpha\in T_P$ of length s, $\{M_e(\beta,s)\}_{\beta\subseteq\alpha}$ is a partition of $\{i\in\mathbb{N}:i<b\}$.

Stage 0**.** At first, put $T_s=\{\langle\rangle\}$, $h_s=0$, $M_e(\langle\rangle,0)=\{i\in\mathbb{N}:i<b\}$, and $\gamma_e(\langle\rangle,0)=\rho_e$.

Stage $s+1$**.** At the beginning of each stage $s+1$, assume that T_s and h_s are given, and that $M_e(\beta,s)$ and $\gamma_e(\beta,s)$ have been already defined for each $s\in\mathbb{N}$ and $\beta\in T_P$ of height $\leq s$. For each $i,e\in\mathbb{N}$ and each $\tau\in 2^{\mathbb{N}}$, *the length-of-agreement function $l^i_e(\tau)$* is the maximal $l\in\mathbb{N}$ such that $\Phi_{\Psi^e_i(\tau)}(\tau;x)\downarrow$ for each $x<l$, and $\Phi_{\Psi^e_i(\tau)}(\tau)\in T_Q$.

Fix a string $\alpha \in T_P$ of length s, and then each i belongs to some $M_e(\beta, s)$ for $\beta \subseteq \alpha$. In this case, the learner Ψ^e_i can act on $\gamma_e(\beta, s)$. Then, we say that *the learner Ψ^e_i requires attention along α at stage* $s+1$ if there exists $\tau \in T_s$ of length h_s extending $\gamma_e(\beta, s)$ such that either of the following conditions are satisfied.

1. Ψ^e_i *changes on* $(\gamma_e(\beta, s), \tau]$, i.e., there is a string σ such that $\gamma_e(\beta, s) \subsetneq \sigma \subseteq \tau$ and $\Psi^e_i(\sigma^-) \neq \Psi^e_i(\sigma)$.
2. or, $l^i_e(\tau) > \max\{l^i_e(\sigma) : \sigma \subseteq \gamma_e(\beta, s)\}$.

Let R_s be the set of all $\alpha \in T_P$ of length s such that some learner requires attention along α at stage $s+1$. For $\alpha \in R_s$, let $m(\alpha)$ be the least m such that there is a string $\beta \subseteq \alpha$ of length m and an index $i \in M_e(\beta, s)$ such that Ψ^e_i requires attention along α at stage $s+1$. That is to say, some learner Ψ^e_i who has already changed his mind $m(\alpha)$ times requires attention.

Claim. For any $\alpha, \beta \in R_s$, we have that $\alpha \upharpoonright m(\alpha) = \beta \upharpoonright m(\beta)$ holds or $\alpha \upharpoonright m(\alpha)$ is incomparable with $\beta \upharpoonright m(\beta)$.

Put $R^*_s = \{\alpha \upharpoonright m(\alpha) : \alpha \in R_s\}$. Then, for $\beta \in R^*_s$, let $i(\beta)$ be the least $i \in M(m(\alpha), s)$ such that Ψ^e_i requires attention along some $\alpha \supseteq \beta$ of length s at stage $s+1$. For $\beta \in R^*_s$, we say that $\Psi^e_{i(\beta)}$ *acts at stage* $s+1$. Moreover, for $\beta \in R^*_s$, let $\tau(\beta)$ be the lexicographically least string $\tau \in T_s$ of length h_s extending $\gamma_e(\beta, s)$ such that τ witnesses that the learner $\Psi^e_{i(\beta)}$ requires attention along some $\alpha \supseteq \beta$ of length s at stage $s+1$. Then $R^{**}_s \subseteq R^*_s$ is defined as the set of all $\beta \in R^*_s$ such that $\Psi^e_{i(\beta)}$ changes on $(\gamma_e(\beta, s), \tau(\beta)]$.

For each $\beta \in R^{**}_s$, put $M_e(\beta, s+1) = M_e(\beta, s) \setminus \{i(\beta)\}$, and put $M_e(\beta^\frown i, s+1) = M_e(\beta, s) \cup \{i(\beta)\}$ for $\beta^\frown i \in T_P$. For any $\beta \notin R^{**}_s$, put $M_e(\beta, s+1) = M_e(\beta, s)$. For each $\beta \in R^*_s$, if $\beta^\frown\sigma \in T_P$ is length $\leq s$ for some $\sigma \in 2^{<\mathbb{N}}$, then put $\gamma_e(\beta^\frown\sigma, s+1) = \tau(\beta)^\frown\sigma$. If $\alpha \in T_P$ of length $\leq s$ has no substring $\beta \in R^*_s$, then put $\gamma_e(\alpha, s+1) = \gamma_e(\alpha, s)$. For each $\alpha \in T_P$ of length s, if $|\gamma_e(\alpha, s+1)| < h_s$ then pick the lexicographically least node $\gamma^*_e(\alpha, s+1) \in T_s$ such that $|\gamma^*_e(\alpha, s+1)| = h_s$ and $\gamma^*_e(\alpha, s+1) \supseteq \gamma_e(\alpha, s+1)$. Otherwise put $\gamma^*_e(\alpha, s+1) = \gamma_e(\alpha, s+1)$. Then, for each $\alpha^\frown i \in T_P$ of length s, put $\gamma_e(\alpha^\frown i, s+1) = \gamma^*_e(\alpha, s+1)^\frown i$. Put $h_{s+1} = \max\{|\gamma_e(\alpha, s+1)| : \alpha \in T_P \ \& \ |\alpha| = s+1\}$. Then we define the approximation of $\widehat{P}$ at stage $s+1$ as follows.

$$T_{s+1} = T_s \cup \{\sigma \subseteq \gamma_e(\alpha, s+1)^\frown 0^{h_{s+1} - |\gamma_e(\alpha, s+1)|} : \alpha \in T_P \ \& \ |\alpha| = s+1 \ \& \ e \in \mathbb{N}\}.$$

Finally, we set $\widehat{P} = [\bigcup_{s \in \mathbb{N}} T_s]$. Clearly, $\widehat{P}$ is a nonempty Π^0_1 subset of $2^{\mathbb{N}}$.

Lemma 150. *$\lim_s \gamma_e(\alpha, s)$ converges for any $e \in \mathbb{N}$ and $\alpha \in T_P$.*

Proof. Note that $\gamma_e(\alpha, s)$ is incomparable with $\gamma_e(\beta, s)$ whenever α is incomparable with β. Therefore, $\gamma_e(\alpha, s)$ changes only when some learner in $M_e(\beta, s)$ acts for some $\beta \subseteq \alpha$. Assume that $\gamma_e(\alpha, s)$ changes infinitely often. Then there is $\beta \subseteq \alpha$, $t \in \mathbb{N}$ and $i \in M_e(\beta, t)$ such that $i \in M_e(\beta, s)$ for any $s \geq t$, and $\Psi^e_{i(\beta)}$ acts infinitely often. However, by our construction, $g^\alpha_e = \lim_s \gamma_e(\alpha, s)$ is computable. Additionally, since $i \in M_e(\beta, s)$ for any $s \geq t$, $\lim_n \Psi^e_{i(\beta)}(g^\alpha_e \upharpoonright n)$ exists, and $\Phi_{\lim_n \Psi^e_{i(\beta)}(g^\alpha_e \upharpoonright n)}(g^\alpha_e) \in Q$. This contradicts our assumption that Q is special. □

For $f \in P$, put $g^f_e = \bigcup_{\alpha \subset f} \lim_s \gamma_e(\alpha, s)$. By this lemma, such g^f_e exists, and we observe that $\widehat{P}$ can be represented as $\widehat{P} = P \cup \{g^f_e : e \in \mathbb{N} \ \& \ f \in P\}$. For each $e \in \mathbb{N}$ and $\alpha \in T_P$, we pick $t(e, \alpha) \in \mathbb{N}$ such that $\gamma_e(\alpha, s) = \gamma_e(\alpha, t)$ for any $s, t \geq t(e, \alpha)$.

Lemma 151. *The $\mathcal{P}$-requirements are satisfied.*

Proof. Assume that $P \leq^{<\omega}_{\omega} \widehat{P}$ via the e-th team $\{\Psi_i\}_{i<b}$ of learners. Then, for any $f \in P$, there is $i < b$ such that $\lim_n \Psi_i(g^f_e \upharpoonright n)$ exists and $\Phi_{\lim_n \Psi_i(g^f_e \upharpoonright n)}(g^f_e) \in Q$. Since $\lim_n \Psi_i(g^f_e \upharpoonright n)$ exists, there exists $\alpha \subset f$ such that $i \in M_e(\alpha, t(e, \alpha))$. However, by the previous claim, no learner in $\bigcup_{\beta \subseteq \alpha} M_e(\beta, t(e, \alpha))$ requires attention after stage $t(e, \alpha)$. This implies $\lim_n l^i_e(g^f_e \upharpoonright n) < \infty$. In other words, $\Phi_{\lim_n \Psi_i(g^f_e \upharpoonright n)}(g^f_e) \notin Q$. This contradicts our assumption. □

$$\begin{tikzcd}
P \arrow[dr, shift left, "\mathrm{id}"] & & & & \\
& P^{\triangledown} \arrow[ul, shift left, "(1{,}<\omega)\text{-computable}"] \arrow[ul, bend right=60, "(1{,}1)\text{-computable}"', "\times" marking] \arrow[dr, shift left, "\mathrm{id}"] & & & \\
& & \bigtriangledown P \arrow[ul, shift left, "(1{,}\omega)\text{-computable}"] \arrow[ul, bend right=60, "(<\omega{,}1)\text{-computable}"', "\times" marking] \arrow[dr, shift left, "\mathrm{id}"] & & \\
& & & P^{\blacktriangledown} \arrow[ul, shift left, "(<\omega{,}\omega)\text{-computable}"] \arrow[ul, bend right=60, "(1{,}\omega)\text{-computable}"', "\times" marking] \arrow[dr, shift left, "\mathrm{id}"] & \\
& & & & \bigtriangledown_{\infty} P \arrow[ul, shift left, "(\omega{,}1)\text{-computable}"] \arrow[ul, bend right=60, "(<\omega{,}\omega)\text{-computable}"', "\times" marking]
\end{tikzcd}$$

FIGURE 4. The dynamic proof model for a special Π^0_1 set $P \subseteq 2^{\mathbb{N}}$.

Lemma 152. *The $\mathcal{G}$-requirements are satisfied.*

Proof. It suffices to show that $f \leq_T g^f_e$ for any $e \in \mathbb{N}$ and $f \in P$. Assume that $\{\Psi_i\}_{i<b}$ is the e-th team of learners. Let H^f_e denote the set of all $i < b$ such that $\lim_n \Psi_i(g^f_e \upharpoonright n)$ converges. By our construction and the first claim, if $i \in H^f_e$ then $i \in M_e(\alpha_i, t(e, \alpha_i))$ for some $\alpha_i \subset f$. If $i \notin H^f_e$ then for any $\alpha \subset f$ there exists s such that $i \in M_e(\alpha, s)$. Set $l = \max_{i \in H^f_e} |\alpha_i|$, and $u = \max_{i \in H^f_e} t(e, \alpha_i)$. For $n > l$, to compute $f(n)$, we wait for stage $v(n) > u$ such that, for every $i \notin H^f_e$, $i \in M_e(f \upharpoonright m, v(n))$ for some $m \geq n + 1$. By our construction, it is easy to see that we can extract $f(n)$ from $\gamma_e(f \upharpoonright n+1, v(n))$, by a uniformly computable procedure in n. □

Thus, we have $Q \not\leq^{<\omega}_{\omega} \widehat{P}$ by Lemma 151, and $P \subseteq \widehat{P} \subseteq \widehat{\mathrm{Deg}}(P)$ by Lemma 152. Thus, $\widehat{P}$ is a Π^0_1 set satisfying $Q \not\leq^{<\omega}_{\omega} \widehat{P} \equiv^{\omega}_1 P$. This concludes the proof. □

Corollary 153. *For any nonempty Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$, if $Q \leq^{<\omega}_{\omega} \widehat{\mathrm{Deg}}(P)$ then Q contains a computable element.*

Proof. Assume that $Q \leq^{<\omega}_{\omega} \widehat{\mathrm{Deg}}(P)$ is satisfied. Suppose that P has no computable element. Then, for $P, Q \subseteq 2^{\mathbb{N}}$, we obtain $Q \not\leq^{<\omega}_{\omega} \widehat{P} \equiv^{\omega}_1 P$ by Theorem 149. Note that the condition $\widehat{P} \equiv^{\omega}_1 P$ implies $\widehat{P} \subseteq \widehat{\mathrm{Deg}}(P)$. Then, $Q \leq^{<\omega}_{\omega} \widehat{\mathrm{Deg}}(P) \leq^1_1 \widehat{P}$. It involves a contradiction. □

5. CHARACTERIZATIONS, APPLICATIONS, AND QUESTIONS

5.1. **Characterization by Piecewise Computability.** We can introduce the iteration of $\triangledown$ and $\triangledown_\infty$ for *any* subsets of Baire space $\mathbb{N}^{\mathbb{N}}$. We also introduce the concatenation $P^\frown Q$ for *any* sets $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, as the noncommutative version of $[\![P \vee Q]\!]^3_{\mathsf{LCM}[2]}$.

Definition 154. Let P, Q be subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.

(1) The *concatenation of P and Q* is defined as follows.

$$P^\frown Q = P \cup \{\sigma^\frown \sharp^\frown g : \sigma \in \mathbb{N}^{<\mathbb{N}} \ \& \ g \in Q\}.$$

(2) $\bigtriangledown^1_1 P = P$; and $\bigtriangledown^1_{m+1} P = P^\frown \bigtriangledown^1_m P$.

(3) $[\bigtriangledown_\infty]^1_n P = \{f \in (\mathbb{N} \times \mathbb{N})^{\mathbb{N}} : (\exists i < n)\ \mathtt{pr}_i(f) \in P\}$.

(4) $\left[\bigtriangledown_\omega\right]^1_n P = [\bigtriangledown_\infty]^1_n P \cap \{f \in (\mathbb{N} \times \mathbb{N}^{\mathbb{N}}) : \mathtt{mc}(f) < \infty\}$.

(5) $\bigtriangledown P = \bigcup_m \bigtriangledown^1_m P$, and $\bigtriangledown_\infty P = \bigcup_m [\bigtriangledown_\infty]^1_m P$.

Remark. Recall that $[\![P \vee Q]\!]^3_{\mathsf{LCM}[2]}$ is $(1, 1)$-equivalent to $[\![P \vee Q]\!]^2_{\mathsf{LCM}[2]}$. They are also $(1, 1)$-equivalent to $P \triangledown Q$ for Π^0_1 sets, and we also have $P \triangledown P \equiv^1_1 P^\frown P$. We may also consider $\bigtriangledown_{v \in (V,E)} P_v$ for a collection $\{P_v\}_{v \in V}$ of *any* subsets of $\mathbb{N}^{\mathbb{N}}$, by setting $\mathrm{Con}(T_{P_v})_{v \in V} = (V \times \mathbb{N})^{\mathbb{N}}$, for a fixed directed graph (V, E). Then one can also introduce $\bigtriangledown^1_m P$ as the dynamic disjunction of P along a linear order of cardinality m.

Definition 155. For a class Γ of subsets of Baire space $\mathbb{N}^{\mathbb{N}}$, we say that a collection $\{Q_i\}_{i\in\mathbb{N}}$ is Γ if the set $\{(i,f)\in\mathbb{N}\times\mathbb{N}^{\mathbb{N}} : f\in Q_i\}$ belongs to Γ. A partition $\{Q_i\}_{i\in\mathbb{N}}$ of Q is Γ if there is a Γ collection $\{Q_i^*\}_{i\in\mathbb{N}}$ such that $Q_i = Q\cap Q_i^*$ for any $i\in\mathbb{N}$. We say that $\{Q_i\}_{i\in\mathbb{N}}$ is *a Γ layer of Q* if there is a Γ collection $\{Q_i^*\}_{i\in\mathbb{N}}$ such that $Q_i^*\subseteq Q_{i+1}^*$ for each $i\in\mathbb{N}$, $\{Q_i^*\}_{i\in\mathbb{N}}$ covers Q, and $Q_i = Q\cap Q_i^*$. We also say that $\{Q_i\}_{i\in\mathbb{N}}$ is *a Γ d-layer of Q* if there is a Γ layer $\{Q_i^*\}_{i\in\mathbb{N}}$ of Q such that $Q_i = Q_i^*\setminus Q_{i-1}^*$ for any $i\in\mathbb{N}$, where $Q_{-1}^* = \emptyset$. Note that every Γ d-layer of Q is a $(\Gamma)_2$ partition of Q. Here a set $Q\subseteq\mathbb{N}^{\mathbb{N}}$ is $(\Gamma)_2$ if it is the difference of two Γ sets.

Theorem 156.

(1) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^1_{<\omega} Q$.*
  - (b) *There is $m\in\mathbb{N}$ such that $\bigtriangledown^1_m P\leq^1_1 Q$.*
  - (c) *There is a finite Π^0_1 d-layer $\{Q_i\}_{i<m}$ of Q such that $P\leq^1_1 Q_i$ for any $i<m$.*

(2) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^1_{\omega|<\omega} Q$.*
  - (b) *There is $m\in\mathbb{N}$ such that $\left[\bigtriangledown_\omega\right]^1_m P\leq^1_1 Q$.*
  - (c) *There is a finite Δ^0_2 partition $\{Q_i\}_{i<m}$ of Q such that $P\leq^1_1 Q_i$ for any $i<m$.*
  - (d) *There is a finite Σ^0_2 cover $\{Q_i\}_{i<m}$ of Q such that $P\leq^1_1 Q_i$ for any $i<m$.*

(3) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^1_\omega Q$.*
  - (b) *$\bigtriangledown P\leq^1_1 Q$.*
  - (c) *$\bigtriangledown P\leq^1_1 \bigtriangledown Q$.*
  - (d) *There is a Π^0_1 partition $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P\leq^1_1 Q_i$ uniformly in $i\in\mathbb{N}$.*
  - (e) *There is a Σ^0_2 cover $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P\leq^1_1 Q_i$ uniformly in $i\in\mathbb{N}$.*

  Here, we say that $P\leq^1_1 R_i$ uniformly in $i\in\mathbb{N}$ *if there is a computable sequence $\{e(i)\}_{i\in\mathbb{N}}$ of indices such that $P\leq^1_1 R_i$ via $\Phi_{e(i)}$ for any $i\in\mathbb{N}$.*

(4) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^{<\omega}_1 Q$.*
  - (b) *There is $m\in\mathbb{N}$ such that $[\bigtriangledown_\infty]^1_m P\leq^1_1 Q$.*
  - (c) *There is a finite partition $\{Q_i\}_{i<m}$ of Q such that $P\leq^1_1 Q_i$ for any $i<m$.*

(5) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^{<\omega}_\omega Q$.*
  - (b) *There is $m\in\mathbb{N}$ such that $[\bigtriangledown_\infty]^1_m \bigtriangledown P\leq^1_1 Q$.*
  - (c) *There is a finite partition $\{Q_i^0\}_{i<m}$ of Q and Π^0_1 partitions $\{Q^1_{i,j}\}_{j\in\mathbb{N}}$ of Q_i^0 such that $P\leq^1_1 Q^1_{i,j}$ uniformly in $(i,j)\in m\times\mathbb{N}$.*
  - (d) *There is a finite partition $\{Q_i^0\}_{i<m}$ of Q and Σ^0_2 covers $\{Q^1_{i,j}\}_{j\in\mathbb{N}}$ of Q_i^0 such that $P\leq^1_1 Q^1_{i,j}$ uniformly in $(i,j)\in m\times\mathbb{N}$.*

(6) *The following are equivalent for any $P,Q\subseteq\mathbb{N}^{\mathbb{N}}$:*
  - (a) *$P\leq^\omega_1 Q$.*
  - (b) *$\bigtriangledown_\infty P\leq^1_1 Q$.*
  - (c) *$\bigtriangledown_\infty P\leq^1_1 \bigtriangledown_\infty Q$.*
  - (d) *$\widehat{\mathrm{Deg}}(P)\supseteq Q$, where $\widehat{\mathrm{Deg}}(P)$ denote the Turing upward closure of P.*
  - (e) *There is an infinite partition $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P\leq^1_1 Q_i$ for any $i\in\mathbb{N}$.*

Proof. Item **1** (a)→(b): Assume that $P\leq^1_{<\omega} Q$ via a learner Ψ with mind-change-bound n. We need to construct a computable function Δ witnessing $\bigtriangledown^1_n P\leq^1_1 Q$. For any $g\in Q$, by uniformly computable procedure, we enumerate *the mind-change location along g of the learner Ψ*, a finite sequence

$\{\mathtt{mcl}_\Psi(g;i)\}_{i<k}$ of length $k<n$, as follows. To simplify out argument, put $\mathtt{mcl}_\Psi(g;-1)=\langle\rangle$. For a given $\mathtt{mcl}_\Psi(g;i-1)\in 2^{<\mathbb{N}}$, we wait for $\mathtt{mcl}_\Psi(g;i)=\min\{\tau:\mathtt{mcl}_\Psi(g;i)\subset\tau\subset g\ \&\ \Psi(\mathtt{mcl}_\Psi(g;i))\neq\Psi(\tau)\}$. If we find $\mathtt{mcl}_\Psi(g;i)$, then we next wait for $\mathtt{mcl}_\Psi(g;i+1)$. Finally, we define $\Delta(g)$ as follows.

$$\begin{aligned}\Delta(g) &= \Phi_{\Psi(\langle\rangle)}(\mathtt{mcl}_\Psi(g;0)^-)^\frown\sharp\\ &{}^\frown\left(\prod_{i<|\mathtt{mcl}(g)|-1}\Phi_{\Psi(\mathtt{mcl}_\Psi(g;i))}(\mathtt{mcl}_\Psi(g;i+1)^-)^\frown\sharp\right)^\frown\Phi_{\Psi(\mathtt{mcl}_\Psi(g;|\mathtt{mcl}(g)|-1))}(g).\end{aligned}$$

It is easy to see that Δ is computable. Note that, $\mathtt{tail}(\Delta(g))=\Phi_{\Psi(\mathtt{mcl}_\Psi(g;|\mathtt{mcl}(g)|-1))}(g)\in P$, since $P\leq^1_{<\omega}Q$ via Ψ, and $\lim_s\Psi(g\upharpoonright s)$ converges to $\Psi(\mathtt{mcl}_\Psi(g;|\mathtt{mcl}(g)|-1))$. Furthermore, $\sharp$ occurs k times in $\Delta(g)$, and $k<n$ because of mind-change-bound n. Thus, $\Delta(g)\in\nabla^1_n P$ for any $g\in Q$, as desired.

Item **1** (b)→(c): For each $\sigma\in\mathbb{N}^{<\mathbb{N}}$ let $\mathtt{count}(\sigma)$ denote the set $\#\{n<|\sigma|:\sigma(n)=\sharp\}$. Assume that $\nabla^1_m P\leq^1_1 Q$ via a computable function Φ. Then $\Phi(g\upharpoonright n)<m$ for each $n\in\mathbb{N}$ and any $g\in Q$. Define $S_i=\{\tau\in\mathbb{N}^{<\mathbb{N}}:\mathtt{count}(\Phi(\tau))\leq i\}$, for each $i<m$. Clearly, $\{[S_i]\}_{i<m}$ is a uniform increasing sequence of Π^0_1 sets. Moreover, $\lim_n\mathtt{count}(\Phi(g\upharpoonright n))<m$ for every $g\in Q$, since $\nabla^1_m P\leq^1_1 Q$ via Φ. This means $g\in[S_i]$ for some $i<m$. Therefore, $Q\subseteq\bigcup_{i<m}[S_i]$. For a given $g\in S_i\setminus S_{i-1}$, we calculate *the i-th mind-change location along g with respect to $\sharp$*, $\mathtt{mcl}_\sharp(g;i)$ such that $\Phi(g;\mathtt{mcl}_\sharp(g;i))$ is the i-th $\sharp$ in $\Phi(g)$, i.e., $\mathtt{mcl}_\sharp(g;i)$ is the least m such that $\mathtt{count}(\Phi(g)\upharpoonright\mathtt{mcl}_\sharp(g;i))>i$. Then we set $\Gamma_i(g)=\Phi(g)^{\leftharpoonup\mathtt{mcl}_\sharp(g;i)+1}$, where recall that $h^{\leftharpoonup k}$ denotes the k-th shift of h, $h^{\leftharpoonup k}=\lambda n.h(n+k)$. It is easy to see that $P\leq^1_1 Q\cap([S_i]\setminus[S_{i-1}])$ via Γ_i.

Item **1** (c)→(a): Assume that $Q\subseteq\bigcup_{i<m}Q_i$ for some computable increasing sequence $\{Q_i\}_{i\in\mathbb{N}}$ of Π^0_1 sets, and $P\leq^1_1 Q\cap(Q_i\setminus Q_{i-1})$ via a computable function $\Phi_{e(i)}$. For each $\sigma\in\mathbb{N}^{<\mathbb{N}}$, we compute the least $i(\sigma)$ such that $\sigma\in T_{Q_{i(\sigma)}}$, i.e., $\sigma\in T_{Q_{i(\sigma)}}\setminus T_{Q_{i(\sigma)-1}}$. Then, on $\sigma\in\mathbb{N}^{<\mathbb{N}}$, the learner Ψ guesses $\Psi(\sigma)=e(i(\sigma))$. By our assumption, for any $g\in Q$, we have $g\in Q_i$ for some $i\in\mathbb{N}$. Then, $\lim_n\Psi(g\upharpoonright n)$ converges to the least $e(i)$ such that $g\in Q_i$. Again, by our assumption, we have $\Phi_{\lim_n\Psi(g\upharpoonright n)}(g)=\Phi_{e(i)}(g)\in P$ for any $g\in Q\cap(Q_i\setminus Q_{i-1})$. Therefore, $P\leq^1_{<\omega}Q$ via the learner Ψ.

Item **2** (a)→(b): Assume that $P\leq^1_{\omega|<\omega}Q$ via a leaner Ψ, where $\#\{\Psi(g\upharpoonright s):s\in\mathbb{N}\}<n$ for any $g\in Q$. We need to construct a computable function Δ witnessing $\left[\nabla_\omega\right]^1_m P\leq^1_1 Q$. We define $\varepsilon(\sigma)$ for each $\sigma\in\mathbb{N}^{<\mathbb{N}}$. Put $\varepsilon(\langle\rangle)=0$. Fix $\sigma\in\mathbb{N}^{<\mathbb{N}}$, and assume that $\varepsilon(\tau)$ has been already defined for each $\tau\subsetneq\sigma$. If $\Psi(\sigma)=\Psi(\tau)$ for some $\tau\subsetneq\sigma$, then we set $\varepsilon(\sigma)=\varepsilon(\tau)$ for such τ. If there is no such τ, then we set $\varepsilon(\sigma)=\max\{\varepsilon(\tau):\tau\subsetneq\sigma\}+1$. Pick the greatest substring $\tau\subsetneq\sigma$ such that $\Psi(\tau)=\Psi(\sigma)$. Then, define $\eta^\sigma=\Phi_{\Psi(\sigma)}(\sigma)^{\leftharpoonup|\Phi_{\Psi(\sigma)}(\tau)|}$. Then, $\Phi_{\Psi(\sigma)}(\sigma)=\Phi_{\Psi(\sigma)}(\tau)^\frown\eta^\sigma$. Here, if there is no such τ, then we may assume $\Phi_{\Psi(\sigma)}(\tau)=\langle\rangle$. Assume that $\Delta(\sigma^-)$ has been already defined. Then, put $\Delta(\sigma)=\Delta(\sigma^-)^\frown\mathtt{write}(\varepsilon(\sigma),\eta^\sigma)$. Here, recall that $\mathtt{write}(\varepsilon(\sigma),\eta^\sigma)=\varepsilon(\sigma)^{|\eta(\sigma)|}\oplus\eta^\sigma=\langle\varepsilon(\sigma),\eta^\sigma(0),\dots,\varepsilon(\sigma),\eta^\sigma(|\eta|-1)\rangle$ represents the instruction to write η on the $\varepsilon(\sigma)$-th tape. Fix $g\in Q$. Note that $\varepsilon(g\upharpoonright s)<n$ for each $s\in\mathbb{N}$, since $\#\{\Psi(g\upharpoonright s):s\in\mathbb{N}\}<n$. Thus, we have $\Delta(g)\in(n\times\mathbb{N})^{\mathbb{N}}$. Moreover, $\mathtt{mc}(\Delta(g))<\infty$, since Ψ is a learner converging on Q. Thus, $\lim_s\Psi(g\upharpoonright s)$ and hence $\lim_s\varepsilon(g\upharpoonright s)$ converge. Therefore, $\mathtt{pr}_{\lim_s\varepsilon(g\upharpoonright s)}(\Delta(g))=\Phi_{\lim_s\Psi(g\upharpoonright s)}(g)\in P$. Hence, $\left[\nabla_\omega\right]^1_n P\leq^1_1 Q$.

Item **2** (b)→(c): Assume that $\left[\nabla_\omega\right]^1_b P\leq^1_1 Q$ via a computable function Φ. For any $p\in\mathbb{N}$, set $Q^\Pi_p=\{g\in\mathbb{N}^{\mathbb{N}}:(\forall m)(\exists n>m)\ (\Phi(g\upharpoonright n))_0=p\}$ and $Q^\Sigma_p=\{g\in\mathbb{N}^{\mathbb{N}}:(\exists m)(\forall n>m)\ (\Phi(g\upharpoonright n))_0=p\}$. Then $Q\cap Q^\Pi_p=Q\cap Q^\Sigma_p=\{g\in Q:\lim_n(\Phi(g\upharpoonright n))_0=p\}$ for any $p\in\mathbb{N}$. Note that, for any $g\in Q$, we have $g\in Q_p$ for some $p<b$, since $(\Phi(g\upharpoonright n))_0<b$ for each $n\in\mathbb{N}$. Therefore, $\{Q\cap Q^\Pi_p\}_{p<b}$ is a finite Δ^0_2 partition of Q such that $P\leq^1_1 Q_p$ via the computable function $\mathtt{pr}_p\circ\Phi$.

Item **2** (c)→(d): Obvious.

Item **2** (d)→(a): Assume that there is a finite Σ^0_2 partition $\{Q_p\}_{p<b}$ of Q such that $P\leq^1_1 Q_p$ via $\Phi_{e(p)}$ for any $p<b$. Then, there is a computable relation $R\subseteq\mathbb{N}\times\mathbb{N}^{<\mathbb{N}}$ such that $Q_p=\{g\in Q:(\exists m)(\forall n>$

$m)\ R(p, g \upharpoonright n)\}$. We set $\Psi(\sigma) = e(\min(\{p : R(p,\sigma)\} \cup \{b-1\}))$. For any $g \in Q$, we have $g \in Q_p$ for some $p < b$, since $Q \subseteq \bigcup_{p<b} Q_p$. Therefore, $\lim_n \Psi(g \upharpoonright n)$ converges to some value $e(p)$, where $g \in Q_p$. Hence, it is easy to see that $P \leq^1_\omega Q$ via Ψ.

Item **3** (a)→(b): By similar argument used in proof of "Item **1** (a)→(b)."

Item **3** (b)→(d): For each $\sigma \in \mathbb{N}^{<\mathbb{N}}$ let $\mathtt{count}(\sigma)$ again denote the set $\#\{n \in \mathbb{N} : \sigma(n) = \sharp\}$. Assume that $\bigtriangledown P \leq^1_1 Q$ via a computable function Φ. Define S_i as follows for each $i \in \mathbb{N}$:

$$S_i = \{\tau \in \mathbb{N}^{<\mathbb{N}} : |\tau| < i \text{ or } (\forall t)\ (i < t \leq |\tau| \ \rightarrow\ \mathtt{count}(\Phi(\tau)) = \mathtt{count}(\Phi(\tau \upharpoonright t)))\}.$$

Clearly, $\{[S_i]\}_{i\in\mathbb{N}}$ is a uniform increasing sequence of Π^0_1 sets. Moreover, the limit $\lim_n \mathtt{count}(\Phi(g \upharpoonright n))$ converges for every $g \in Q$, since $\bigtriangledown P \leq^1_1 Q$ via Φ. This means $g \in [S_i]$ for some $i \in \mathbb{N}$. Therefore, $Q \subseteq \bigcup_i [S_i]$. For a given $g \in Q$, Γ_i computes *the last mind-change location along g with respect to* $\sharp$, $\mathtt{mcl}^*_\sharp(g \upharpoonright i) = \max\{m < |\Phi(g \upharpoonright i)| : \Phi(g \upharpoonright i; m) = \sharp\} + 1$, and set $\Gamma_i(g) = \Phi(g)^{\leftarrow \mathtt{mcl}^*_\sharp(g\upharpoonright i)}$. It is easy to see that $\{\Gamma_i\}_{i\in\mathbb{N}}$ is uniformly computable, and $P \leq^1_1 Q \cap [S_i]$ via Γ_i. Set $Q_i = [S_i]$ for each $i \in \mathbb{N}$. Then $Q_{i+1} \setminus Q_i = \bigcup_{n\in\mathbb{N}} (Q_{i+1} \cap [\rho^i_n])$, where ρ^i_n is the n-th leaf of the corresponding tree of Q_i. We set $Q_{i+1,n} = Q_{i+1} \cap [\rho^i_n]$. Then $\{Q \cap Q_0\} \cup \{Q \cap Q_{i+1,n}\}_{i,n\in\mathbb{N}}$ is a Π^0_1 partition of Q, and $P \leq^1_1 Q \cap Q_{i+1,n}$ via Γ_{i+1}, for each $i, n \in \mathbb{N}$.

Item **3** (d)→(e): Obvious.

Item **3** (e)→(a): Assume that there is a Σ^0_2 partition $\{Q_p\}_{p\in\mathbb{N}}$ of Q such that $P \leq^1_1 Q_p$ via $\Phi_{e(p)}$ for any $p \in \mathbb{N}$. Then, there is a computable relation $R \subseteq \mathbb{N} \times \mathbb{N}^{<\mathbb{N}}$ such that $Q_p = \{g \in Q : (\exists m)(\forall n > m)\ R(p, g \upharpoonright n)\}$. We set $\Psi(\sigma) = e(\min\{p : R(p,\sigma)\})$. For any $g \in Q$, we have $g \in Q_p$ for some $p \in \mathbb{N}$, since $Q \subseteq \bigcup_p Q_p$. Therefore, $\lim_n \Psi(g \upharpoonright n)$ converges to some value $e(p)$, where $g \in Q_p$. Hence, it is easy to see that $P \leq^1_\omega Q$ via Ψ.

Item **3** (b)↔(c): Note that $Q \equiv^1_\omega \bigtriangledown Q$. Thus, $\bigtriangledown P \leq^1_1 Q$ if and only if $P \leq^1_\omega Q$ if and only if $P \leq^1_\omega \bigtriangledown Q$ if and only if $\bigtriangledown P \leq^1_1 \bigtriangledown Q$.

Item **4** (a)→(b): Assume that $P \leq^{<\omega}_1 Q$ via a finite collection $\{\Phi_e\}_{e<b}$ of partial computable functions. We need to construct a computable function Δ witnessing $[\bigtriangledown_\infty]^1_b P \leq^1_1 Q$. For $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $e < b$, we define $\eta^\sigma_e = \Phi_e(\sigma)^{\leftarrow |\Phi_e(\sigma^-)|}$, i.e., $\Phi_e(\sigma) = \Phi_e(\sigma^-)^\frown \eta^\sigma_e$. We recursively construct a computable function $\Delta : Q \rightarrow [\bigtriangledown_\infty]^1_b P$. Assume that $\Delta(\sigma^-)$ is already defined. Put $\Delta(\sigma) = \Delta(\sigma^-)^\frown \bigsqcap_{e<b} \mathtt{write}(e, \eta^\sigma_e)$. Note that $\mathtt{pr}_e(\Delta(g)) \in P$ if $\Phi_e(g) \in P$. Thus, for any $g \in Q$, we have $\mathtt{pr}_e(\Delta(g)) \in P$ for some $e < b$. In other words, $[\bigtriangledown_\infty]^1_b P \leq^1_1 Q$ via Δ.

Item **4** (b)→(c): Assume that $[\bigtriangledown_\infty]^1_b P \leq^1_1 Q$ via a computable function Φ. Set $Q_e = \{g \in Q : \mathtt{pr}_e(\Phi(g)) \in P\}$. Then, it is easy to see that $\{Q_e\}_{e<b}$ covers Q, and $P \leq^1_1 Q_e$ via $\mathtt{pr}_e \circ \Phi$ for each $e < b$.

Item **4** (c)→(a): Assume that there is a partition $\{Q_i\}_{i<m}$ such that $P \leq^1_1 Q_i$ via a computable function $\Phi_{e(i)}$ for any $i < m$. Then, $P \leq^1_1 Q$ via $\{\Phi_j : j \leq \max_{i<m} e(i)\}$.

Item **5** (a)→(b): Assume that $P \leq^{<\omega}_\omega$ via a team $\{\Psi_i\}_{i<b}$ of learners. We construct a computable function Δ. We first set $\Delta(\langle\rangle) = \langle\rangle$. Fix $\sigma \in \mathbb{N}^{<\mathbb{N}}$, and assume that $\Delta(\sigma^-)$ has been already defined. We define $\eta^\sigma_i \in \mathbb{N}^{<\mathbb{N}}$ for each $i < b$ as follows. Fix $i < b$. If $\Psi_i(\sigma) = \Psi_i(\sigma^-)$, say $\Phi_{\Psi_i(\sigma)}(\sigma) = \Phi_{\Psi_i(\sigma)}(\sigma^-)^\frown\tau$ for some string $\tau \in \mathbb{N}^{<\mathbb{N}}$, then let η^σ_i be such τ. If $\Psi_i(\sigma) \neq \Psi_i(\sigma^-)$, then we put $\eta^\sigma_i = \sharp^\frown \Phi_{\Psi_i(\sigma)}(\sigma)$. Then, we define $\Delta(\sigma) = \Delta(\sigma^-)^\frown \bigsqcap_{i<b} \mathtt{write}(i, \eta^\sigma_i)$. Pick $g \in Q$. Then, by our assumption, $\Phi_{\lim_n \Psi_i(g\upharpoonright n)}(g) \in P$ for some $i < b$. Then $\mathtt{tail}(\mathtt{pr}_i(\Delta(g)))$ converges, and $\mathtt{tail}(\mathtt{pr}_i(\Delta(g)))^{\leftarrow 1} = \Phi_{\lim_n \Psi_i(g\upharpoonright n)}(g) \in P$. Thus, $\Delta(g) \in [\bigtriangledown_\infty]^1_b \bigtriangledown P$.

Item **5** (b)→(c): Assume that $[\bigtriangledown_\infty]^1_b \bigtriangledown P \leq^1_1 Q$ via a computable function Φ. Set $Q^0_i = \{g \in Q : \mathtt{pr}_i(\Phi(g)) \in \bigtriangledown P\}$. Then, it is easy to see that $\{Q_e\}_{e<b}$ is a partition of Q. Then we can use the similar argument as in proof of "Item **3** (b)→(d)."

Item **5** (c)→(d): Obvious.

Item **5** (d)→(a): Assume that there is a finite partition $\{Q^0_i\}_{i<m}$ and Σ^0_2 partitions $\{Q^1_{i,j}\}_{j\in\mathbb{N}}$ of Q^0_i such that $P \leq^1_1 Q^1_{i,j}$ via $\Phi_{e(i,j)}$ for any $(i, j) \in m \times \mathbb{N}$. Then, there are computable relations $R_i \subseteq \mathbb{N} \times \mathbb{N}^{<\mathbb{N}}$ such that $Q^1_{i,j} = \{g \in Q^0_i : (\exists m)(\forall n > m)\ R_i(j, g \upharpoonright n)\}$. We set $\Psi_i(\sigma) = e(i, \min\{j : R_i(j,\sigma)\})$ for each $i < m$. For any $g \in Q$, if $g \in Q^0_i$, then we have $g \in Q^1_{i,j}$ for some $j \in \mathbb{N}$, since $Q^0_i \subseteq \bigcup_{j\in\mathbb{N}} Q^1_{i,j}$. Therefore, for any $g \in Q^0_i$, the limit $\lim_n \Psi_i(g \upharpoonright n)$ converges to some value $e(i, j)$, where we must have $g \in Q^i_j$ for such $j \in \mathbb{N}$. Hence, by our assumption, we have $P \leq^{<\omega}_{\omega} Q$ via a team $\{\Psi_i\}_{i<m}$ of learners.

Item **6** (a)→(b): Assume that $P \leq^{\omega}_1 Q$. We need to construct a computable function Δ witnessing $\bigtriangledown_\infty P \leq^1_1 Q$. For $\sigma \in \mathbb{N}^{<\mathbb{N}}$ and $e < |\sigma|$, we again define $\eta^\sigma_e = \Phi_e(\sigma)^{\leftharpoonup|\Phi_e(\sigma^-)|}$, i.e., $\Phi_e(\sigma) = \Phi_e(\sigma^-)^\frown\eta^\sigma_e$. We recursively construct a computable function $\Delta : Q \to \bigtriangledown_\infty P$. Assume that $\Delta(\sigma^-)$ has been already defined. Put $\Delta(\sigma) = \Delta(\sigma^-)^\frown(\bigsqcap_{e<|\sigma|} \mathtt{write}(e, \eta^\sigma_e))^\frown(\mathtt{write}(e, \Phi_e(\sigma)))$. Note that $\mathtt{pr}_e(\Delta(g)) = \Phi_e(g)$. Thus, for any $g \in Q$, we have $\mathtt{pr}_e(\Delta(g)) \in P$ for some $e \in \mathbb{N}$. In other words, $\bigtriangledown_\infty P \leq^1_1 Q$ via Δ.

Item **6** (a)↔(d): Obvious.

Item **6** (b)→(e): Assume that $\bigtriangledown_\infty P \leq^1_1 Q$ via a computable function Φ. Set $Q_e = \{g \in Q : \mathtt{pr}_e(\Phi(g)) \in P\}$. Then, clearly $\{Q_e\}_{e<b}$ covers Q, and $P \leq^1_1 Q_e$ via a computable function $\mathtt{pr}_e \circ \Phi$.

Item **6** (e)→(a): If there is an infinite partition $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P \leq^1_1 Q_i$ for any $i \in \mathbb{N}$, then, in particular, $Q_i \subseteq \widehat{\mathrm{Deg}}(P)$ for any $i \in \mathbb{N}$. This implies $Q = \bigcup_{i\in\mathbb{N}} Q_i \subseteq \widehat{\mathrm{Deg}}(P)$, and hence $P \leq^{\omega}_1 Q$.

Item **6** (b)↔(c): Note that $Q \equiv^{\omega}_1 \bigtriangledown_\infty Q$. Thus, $\bigtriangledown_\infty P \leq^1_1 Q$ if and only if $P \leq^{\omega}_1 Q$ if and only if $P \leq^{\omega}_1 \bigtriangledown_\infty Q$ if and only if $\bigtriangledown_\infty P \leq^1_1 \bigtriangledown_\infty Q$. □

Proposition 157. *Let P, Q be Π^0_1 subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

1. *$P \leq^{<\omega}_{tt,1} Q$ if and only if there is a finite Π^0_1 cover $\{Q_i\}_{i<n}$ of Q such that $P \leq^1_1 Q_i$ for each $i < n$.*
2. *$P \leq^{\omega}_{tt,1} Q$ if and only if there is a Π^0_1 cover $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P \leq^1_1 Q_i$ for each $i \in \mathbb{N}$.*
3. *$P \leq^{<\omega}_1 Q$ if and only if there is a finite Π^0_2 d-layer $\{Q_i\}_{i<n}$ of Q such that $P \leq^1_1 Q_i$ for each $i < n$.*
4. *$P \leq^{\omega}_1 Q$ if and only if there is a Π^0_2 partition $\{Q_i\}_{i\in\mathbb{N}}$ of Q such that $P \leq^1_1 Q_i$ uniformly in $i \in \mathbb{N}$.*

Proof. For the items 1 and 2, we assume that $P \leq^{\omega}_{tt,1} Q$ via a collection $\{\Theta_i\}_{i\in\mathbb{N}}$ of total computable functions. Note that $\Theta^{-1}(P)$ is Π^0_1, whenever Θ is total computable, and P is Π^0_1. Thus, it is easy to show that $Q_i = Q \cap \Theta^{-1}_i(P)$ satisfies the desired condition. We next assume that $P \leq^1_1 Q_i$ via Φ_{e_i} for each $i \in \mathbb{N}$, where we do not assume that $i \mapsto e_i$ is computable. Then $\Phi_{e_i} : Q_i \to P$ can be extended to a total computable function on $\mathbb{N}^{\mathbb{N}}$, since Q_i is Π^0_1, for each $i \in \mathbb{N}$. Therefore, $P \leq^{\omega}_{tt,1} Q$ via $\{\Phi_{e_i}\}_{i\in\mathbb{N}}$.

For the items 3 and 4, we assume that $P \leq^{\omega}_1 Q$. Then, by Proposition 50, there is a computable function $\Psi : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}$ such that $\liminf_n \Psi(g \upharpoonright n) \in P$ for any $g \in Q$. We put $Q_e = \{g \in Q : \liminf_n \Psi(g \upharpoonright n) \leq e\}$ for each $e \in \mathbb{N}$. Then, the collection $\{Q_e\}_{e\in\mathbb{N}}$ is uniformly Π^0_2, and it forms a layer of Q, i.e., $\bigcup_{e\in\mathbb{N}} Q_e$ covers Q. Moreover, for each $e \in \mathbb{N}$, we have $P \leq^1_1 Q_e \setminus Q_{e-1}$ via Φ_e, where $Q_{-1} = \emptyset$. Thus, $\{Q_e \setminus Q_{e-1}\}_{e\in\mathbb{N}}$ is a Π^0_2 d-layer of Q, and $P \leq^1_1 Q_e \setminus Q_{e-1}$ uniformly in $e \in \mathbb{N}$. Furthermore, each $(\Pi^0_2)_2$ set $Q_e \setminus Q_{e-1}$ is divided into infinitely many Π^0_2 sets as follows. For each $e \in \mathbb{N}$, let $\{Q^n_{e-1}\}_{n\in\mathbb{N}}$ be a uniform collection of Π^0_1 sets such that $\bigcup_{n\in\mathbb{N}} Q^n_{e-1}$ is the compliment of Q_{e-1}, i.e., $Q_e \setminus Q_{e-1} = Q_e \cap \bigcup_{n\in\mathbb{N}} Q^n_{e-1} = \bigcup_{n\in\mathbb{N}}(Q_e \cap Q^n_{e-1})$. Then, $R^n_e = Q_e \cap Q^n_{e-1}$ is Π^0_2, for each $e, n \in \mathbb{N}$. Note that we can calculate an index of such collection $\{R^n_e\}_{n\in\mathbb{N}}$ from any Σ^0_2-indices of Q_{e-1} and Q_e. Thus, $\{R^n_e\}_{e,n\in\mathbb{N}}$ is a Π^0_2 partition of Q, and $P \leq^1_1 R^n_e$ via Φ_e, for each $e, n \in \mathbb{N}$. □

The direct sum $\oplus$ is also called the disjoint union $\sqcup$. It is also correct that, for Π^0_1 sets, if P and Q are disjoint, then $P \oplus Q$ is equivalent to $P \cup Q$ modulo the $(1, 1)$-equivalence. However, if P and Q are not Π^0_1, the above claim is false, in general.

Proposition 158.

1. *For every sets $P, Q \subseteq \mathbb{N}^{\mathbb{N}}$, the following conditions are equivalent:*

   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *partition* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There is a computable function* $\Gamma : Q \to P$.

   In particular, if $P \cap Q = \emptyset$*, then* $P \cup Q \equiv^1_1 P \oplus Q$.

(2) *For any special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there exists a* $(\Pi^0_1)_2$ *set* $Q \subseteq 2^{\mathbb{N}}$ *such that* $P \cap Q = \emptyset$ *but* $P \cup Q <^1_1 P \oplus Q$.

Proof. For the item 1, we assume that $\Gamma_i : Q_i \to P$ is computable for each $i < m$ with a finite Π^0_1 partition $\{Q_i\}_{i<m}$ of Q. Then we can extend each function Γ_i to a total computable function Γ^*_i. For a given $g \in Q$, we wait for $g \notin Q_j$ for $m-1$ many j's. By this way, we can calculate such $i < m$ such that $g \notin Q_j$ for each $j \neq i$. Then $\Gamma_i(g) \in Q$. For the item 2, put $Q = (P \triangledown P) \setminus P$. For any $g \in Q$, there is a leaf $\rho \in L_P$ such that $\rho \subset g$. So we wait for such a leaf $\rho \in L_P$. Then $g^{\leftarrow|\rho|}$ belongs to P. Hence, $P \leq^1_1 Q$. Thus, we have $P \leq^1_1 P \oplus Q$, while $P \cup Q = P \triangledown P <^1_1 P$ by Theorem 83. □

Corollary 159 (As piecewise computable functions)**.**

(1) *There exist* Π^0_1 *sets* $P, Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *partition* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *cover* $\{Q_i\}_{i<m}$ *of* Q.

(2) *For every special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there is a* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *partition* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.

(3) *There exist* Π^0_1 *sets* $P, Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Δ^0_2 *partition* $\{Q_i\}_{i<m}$ *of* Q.

(4) *There exist* Π^0_1 *sets* $P, Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *uniformly with a* Σ^0_2 *cover* $\{Q_i\}_{i\in\mathbb{N}}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_2 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.

(5) *There is a special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, the following conditions are equivalent for any* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$*:*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_1 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Σ^0_2 *cover* $\{Q_i\}_{i<m}$ *of* Q.
   (c) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_2 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.

(6) *For every special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there is a* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *with a finite* Π^0_2 *d-layer* $\{Q_i\}_{i<m}$ *of* Q.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *uniformly with a* Π^0_1 *partition* $\{Q_i\}_{i\in\mathbb{N}}$ *of* Q.

(7) *For every special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there is a* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_i : Q_i \to P$ *uniformly with a* Σ^0_2 *cover* $\{Q_i\}_{i\in\mathbb{N}}$ *of* Q.
   (b) *There are computable functions* $\Gamma_{i,j} : Q^1_{i,j} \to P$ *uniformly with a finite* Π^0_2 *d-layer* $\{Q^0_i\}_{i<m}$ *and* Π^0_1 *partitions* $\{Q^1_{i,j}\}_{j\in\mathbb{N}}$ *of* Q^0_i.

(8) *For every special* Π^0_1 *set* $P \subseteq 2^{\mathbb{N}}$*, there is a* Π^0_1 *set* $Q \subseteq 2^{\mathbb{N}}$ *which satisfies (b)&¬(a):*
   (a) *There are computable functions* $\Gamma_{i,j} : Q^1_{i,j} \to P$ *uniformly with a finite* Π^0_2 *d-layer* $\{Q^0_i\}_{i<m}$ *and* Σ^0_2 *covers* $\{Q^1_{i,j}\}_{j\in\mathbb{N}}$ *of* Q^0_i.
   (b) *There are computable functions* $\Gamma_i : Q_i \to P$ *uniformly with a* Π^0_2 *partition* $\{Q_i\}_{i\in\mathbb{N}}$ *of* Q.

Proof. Item **1**. By Corollary 54 (2), Proposition 157 (1), and Proposition 158 (1). Item **2**. By Corollary 88, Theorem 156 (1), and Proposition 158 (1). Item **3**. By Corollary 59 (2), and Theorem 156 (1,2). Item **4**. By Corollary 61 (1), and Theorem 156 (3), and Proposition 157 (3). Item **5**. By Theorem 72, Theorem 156 (1,2), and Proposition 157 (3). Item **6**. By Corollary 118 (2), Theorem 156 (3), and Proposition 157 (3). Item **7**. By Corollary 129, Theorem 156 (3,5). Item **8**. By Theorem 149, Theorem 156 (5), and Proposition 157 (4). □

Definition 160 (Hyperconcatenation)**.** For any strings $\sigma \in (\mathbb{N} \cup \{\texttt{pass}\})^{<\mathbb{N}}$ and $\tau \in (\mathbb{N} \cup \{\sharp, \texttt{pass}\})^{<\mathbb{N}}$, the *content of* σ, $\texttt{content}(\sigma)$, and the *walk of* τ, $\texttt{walk}(\tau)$, is inductively defined as follows.

$$\texttt{content}(\langle\rangle) = \langle\rangle, \qquad \texttt{content}(\sigma) = \begin{cases} \texttt{content}(\sigma^-)^\frown\sigma(|\sigma|-1) & \text{if } \sigma(|\sigma|-1) \neq \texttt{pass}, \\ \texttt{content}(\sigma^-) & \text{otherwise.} \end{cases}$$

$$\texttt{walk}(\tau \upharpoonright 1) = \langle\rangle, \qquad \texttt{walk}(\tau) = \begin{cases} \texttt{walk}(\tau^-)^\frown v & \text{if } \tau(|\tau|-2) = \sharp \;\&\; \tau(|\tau|-1) = v, \\ \texttt{walk}(\tau^-) & \text{otherwise.} \end{cases}$$

Then, the *content* of $f \in (\mathbb{N} \cup \{\texttt{pass}\})^{\mathbb{N}}$ and the *walk* of $g \in (\mathbb{N} \cup \{\sharp, \texttt{pass}\})^{\mathbb{N}}$ are defined by $\texttt{content}(f) = \bigcup_{n\in\mathbb{N}} \texttt{content}(f \upharpoonright n)$ and $\texttt{walk}(g) = \bigcup_{n\in\mathbb{N}} \texttt{walk}(g \upharpoonright n)$, respectively. Let P and Q be *any* subsets of Baire space $\mathbb{N}^{\mathbb{N}}$. The *hyperconcatenation* $Q \blacktriangledown P$ and the *non-Lipschitz hyperconcatenation* $Q \blacktriangledown^\circ P$ of Q and P are defined as follows.

$$Q \blacktriangledown P = \{g \in (\mathbb{N} \cup \{\sharp\})^{\mathbb{N}} : \texttt{walk}(g) \in Q \text{ or } \texttt{tail}(g)^{\leftharpoonup 1} \in P\},$$

$$Q \blacktriangledown^\circ P = \{g \in (\mathbb{N} \cup \{\sharp, \texttt{pass}\})^{\mathbb{N}} : \texttt{content} \circ \texttt{walk}(g) \in Q \text{ or } \texttt{tail}(g)^{\leftharpoonup 1} \in P\}.$$

Theorem 161 (As the Law of Excluded Middle)**.**

1. *The following are equivalent for any* $P, Q, R \subseteq \mathbb{N}^{\mathbb{N}}$*:*
   - (a) $Q^\frown P \leq^1_1 R$.
   - (b) *There is a* Σ^0_1 *sentence* φ *with computable functions* $\Phi_{e(0)}, \Phi_{e(1)}$ *such that*
     - *if* $g \in R$ *satisfies* $\varphi(g)$*, then* $\Phi_{e(0)}(g) \in P$*;*
     - *if* $g \in R$ *satisfies* $\neg\varphi(g)$ *then* $\Phi_{e(1)}(g) \in Q$*.*
2. *The implications* (b$^+$) $\to$ (a) $\to$ (a$^-$) $\leftrightarrow$ (b$^-$) *hold for any* $P, Q, R \subseteq \mathbb{N}^{\mathbb{N}}$*:*
   - (a) $Q \blacktriangledown P \leq^1_1 R$.
   - (a$^-$) $Q \blacktriangledown^\circ P \leq^1_1 R$.
   - (b$^+$) *There is a* Σ^0_2 *sentence* $\varphi \equiv \exists v \theta(v)$ *with a uniform sequence* $\{\Gamma_i\}_{i\in\mathbb{N}}, \Delta$ *of computable functions such that*
     - *if* $g \in R$ *satisfies* $\theta(v)$*, then* $\Gamma_v(g; u)\downarrow$ *for any* $u \in \mathbb{N}$*, and* $\Gamma_v(g) \in P$*.*
     - *if* $g \in R$ *satisfies* $\neg\theta(v)$*, then* $\Delta(g; u)\downarrow$ *for any* $u \leq v$*, and* $[\Delta(g) \upharpoonright v+1]$ *intersects with* Q*.*
   - (b$^-$) *There is a* Σ^0_2 *sentence* $\varphi \equiv \exists v \theta(v)$ *with a uniform sequence* $\{\Gamma_i\}_{i\in\mathbb{N}}, \Delta$ *of computable functions such that*
     - *if* $g \in R$ *satisfies* $\theta(v)$*, then* $\Gamma_v(g; u)\downarrow$ *for any* $u \in \mathbb{N}$*, and* $\Gamma_v(g) \in P$*.*
     - *if* $g \in R$ *satisfies* $\neg\exists v \theta(v)$*, then* $\Delta(g; u)\downarrow$ *for any* $u \in \mathbb{N}$*, and* $\Delta(g) \in Q$*.*

Proof. Item **1** (a)→(b): The desired Σ^0_1 sentence is defined as $\varphi(g) \equiv (\exists n)\ \Phi(g; n) = \sharp$.

Item **1** (b)→(a): Assume that there is a Σ^0_1 set $S \subseteq \mathbb{N}^{\mathbb{N}}$ such that $P \leq^1_1 R \cap S$ via $\Phi_{e(0)}$, and $Q \leq^1_1 R \cap S$ via $\Phi_{e(1)}$. Since $S \subseteq \mathbb{N}^{\mathbb{N}}$ is Σ^0_1, there is a c.e. set W_S such that $\sigma \in W_S$ if and only if $[\sigma] \subseteq S$. For a given $\sigma \in \mathbb{N}^{<\mathbb{N}}$, if there is no $n < |\sigma|$ such that $\sigma \upharpoonright n \in W_S$, then set $\Gamma(\sigma) = \Phi_{e(1)}(\sigma)$. If there is $n < |\sigma|$ such that $\sigma \upharpoonright n \in W_S$, then, for such n, set $\Gamma(\sigma) = \Phi_{e(1)}(\sigma \upharpoonright n)^\frown\sharp^\frown\Phi_{e(0)}(\sigma)$. Then, it is easy to see that Γ is computable, and $Q^\frown P \leq^1_1 R$ via Γ.

Item **2** (b$^+$)→(a): Assume that $S_i = \{g \in \mathbb{N}^{\mathbb{N}} : \Theta(g; i)\uparrow\}$ for some computable function Θ, and that $P \leq^1_1 R \cap S_i$ via Γ_i and $Q \leq^1_1 R \setminus \bigcup_{i\in\mathbb{N}} S_i$ via Δ. For a string $\sigma \in \mathbb{N}^{<\mathbb{N}}$, define $d(\sigma)$ and $t(\sigma; i)$ as follows:

$$d(\sigma) = \max\{d \in \mathbb{N} : (\forall i < d)\ \Theta(\sigma; i)\downarrow\};$$

$$t(\sigma; i) = \min\{t \in \mathbb{N} : \Theta(\sigma \upharpoonright t; i)\downarrow\}, \text{ for any } i < d(\sigma).$$

Then let us define $\Lambda(\sigma) = \prod_{i<d(\sigma)} (\Gamma_i(\sigma \upharpoonright t(\sigma; i))^\frown\sharp^\frown\Delta(\sigma; i))^\frown\Gamma_{d(\sigma)}(\sigma)$.

Item **2** (a$^-$)→(b$^-$): Assume that $Q \blacktriangledown^\circ P \leq^1_1 R$ via a computable function Φ. Set $S_v = \{g \in \mathbb{N}^{\mathbb{N}} : (\forall n \geq v)\ \Phi(g; n) \neq \sharp\}$. For a string $\sigma \in \mathbb{N}^{<\mathbb{N}}$, we first computes the following $\texttt{count}(\sigma)$ and $\texttt{mcl}_\sharp(\sigma, n)$ for each

$n \in \mathbb{N}$:

$$\begin{aligned} \mathtt{count}(\sigma) &= \#\{m < |\sigma| : \Phi(\sigma; m) = \sharp\}, \\ \mathtt{mcl}_\sharp(\sigma, n) &= \min\{m \leq |\sigma| : \mathtt{count}(\sigma \upharpoonright m) > n\}, \text{ if such } m \text{ exists.} \end{aligned}$$

Then set $\Gamma_v(\sigma) = \Phi(\sigma)^{\leftarrow \mathtt{mcl}_\sharp(\sigma, \mathtt{count}(\sigma \upharpoonright v))+1}$; and set $\Delta(\sigma) = \lambda n.\Phi(\sigma, \mathtt{mcl}_\sharp(\sigma, n))$. Note that if $g \in R \cap S_k$ for some $k \in \mathbb{N}$, then $\Gamma_k(g) \in P$; otherwise, $\Delta(g) \in Q$. Therefore, $P \leq^1_1 R \cap S_v$ via Γ_v and $Q \leq^1_1 R \setminus S$ via Δ.

Item (b^-)→(a^-): For each $\sigma \in \mathbb{N}^{<\mathbb{N}}$, let $v(\sigma)$ be the least v such that $R(u, v, \sigma)$ holds for all $u < |\sigma|$, where $\varphi(g) \equiv (\exists v)(\forall u)R(u, v, g \upharpoonright u)$. We inductively define a computable function Φ as follows. We first set $\Phi(\langle\rangle) = \langle\rangle$. Assume that $\Phi(\sigma^-)$ has been already defined.

$$\Phi(\sigma) = \begin{cases} \Phi(\sigma^-)^\frown\gamma, & \text{if } v(\sigma) = v(\sigma^-) \;\&\; \Gamma_{v(\sigma)}(\sigma) = \mathtt{tail}^+(\Phi(\sigma^-))^\frown\gamma, \\ \Phi(\sigma^-)^\frown\langle\sharp, \delta(0)\rangle, & \text{if } v(\sigma) \neq v(\sigma^-) \;\&\; \Delta(\sigma) = \mathtt{content} \circ \mathtt{walk}(\Phi(\sigma^-))^\frown\delta, \\ \Phi(\sigma^-)^\frown\langle\sharp, \mathtt{pass}\rangle, & \text{if } v(\sigma) \neq v(\sigma^-) \;\&\; \Delta(\sigma) = \mathtt{content} \circ \mathtt{walk}(\Phi(\sigma^-)). \end{cases}$$

For any $g \in \mathbb{N}^\mathbb{N}$, if $\varphi(g) \equiv (\exists v)(\forall u)R(u, v, g \upharpoonright u)$, then for the least such $v \in \mathbb{N}$, we have $\mathtt{tail}^+(\Phi(g)) = \Gamma_v(g)$. Otherwise, we have $\mathtt{content} \circ \mathtt{walk}(\Phi(g)) = \Delta(g)$. Hence, $\Phi(g) \in Q \blacktriangledown^\circ P$, for any $g \in R$. □

Remark. For degree structures $\mathcal{D}_u$ and $\mathcal{D}_r$ on $\mathcal{P}(\mathbb{N}^\mathbb{N})$, each operator $O : \mathcal{P}(\mathbb{N}^\mathbb{N}) \to \mathcal{P}(\mathbb{N}^\mathbb{N})$ induces the new operator $O_{ur} : \mathcal{D}_u \to \mathcal{D}_r$ defined by $O_{ur}(\deg_u(P)) = \deg_r(O(P))$ for any $P \subseteq \mathbb{N}^\mathbb{N}$. We identify O with O_{ur} whenever O_{ur} is well-defined. Recall that every partially ordered set can be viewed as a category. Sorbi [68] showed that $\widehat{\mathrm{Deg}} : \mathcal{D}^\omega_1 \to \mathcal{D}^1_1$ is left-adjoint to $\mathrm{id} : \mathcal{D}^1_1 \to \mathcal{D}^\omega_1$, and $\mathrm{id} \circ \widehat{\mathrm{Deg}} : \mathcal{D}^\omega_1 \to \mathcal{D}^\omega_1$ is identity. He also showed that $\widehat{\mathrm{Deg}} : \mathcal{D}^\omega_1 \to \mathcal{D}^1_1$ preserves the supremum, but does not preserve the infimum, i.e., $\widehat{\mathrm{Deg}}(P \oplus Q) <^1_1 \widehat{\mathrm{Deg}}(P) \oplus \widehat{\mathrm{Deg}}(Q)$ for some $P, Q \subseteq \mathbb{N}^\mathbb{N}$.

Corollary 162. $\bigtriangledown : \mathcal{D}^1_\omega \to \mathcal{D}^1_1$ *is left-adjoint to* $\mathrm{id} : \mathcal{D}^1_1 \to \mathcal{D}^1_\omega$*, and* $\mathrm{id} \circ \bigtriangledown : \mathcal{D}^1_\omega \to \mathcal{D}^1_\omega$ *is identity. Additionally,* $\bigtriangledown : \mathcal{D}^1_\omega \to \mathcal{D}^1_1$ *preserves the supremum, but does not preserve the infimum, i.e.,* $\bigtriangledown(P \oplus Q) <^1_1 \bigtriangledown P \oplus \bigtriangledown Q$ *for some* $P, Q \subseteq \mathbb{N}^\mathbb{N}$.

Proof. By Theorem 156 and 83. For the infimum, see the next proposition. □

To simplify our notations, in the next proposition, we use notations $P \triangledown_\omega Q$ and $P \triangledown_\infty Q$ instead of $[\![P \vee Q]\!]^2_{\mathsf{LCM}}$ and $[\![P \vee Q]\!]^2_{\mathsf{CL}}$, respectively. However, keep in mind that these notations may be different from original meanings of $\triangledown_\omega$ and $\triangledown_\infty$ when P and Q are not Π^0_1.

Proposition 163. *Let P and Q be subsets of Baire space* $\mathbb{N}^\mathbb{N}$.

(1) $\bigtriangledown(P \oplus Q) \equiv^1_1 \bigtriangledown P \triangledown_\omega \bigtriangledown Q$.
(2) $\widehat{\mathrm{Deg}}(P \oplus Q) = \widehat{\mathrm{Deg}}(P) \cup \widehat{\mathrm{Deg}}(Q) \equiv^1_1 \widehat{\mathrm{Deg}}(P) \triangledown_\infty \widehat{\mathrm{Deg}}(Q)$.

But, there are subsets P and Q of Baire space $\mathbb{N}^\mathbb{N}$ *satisfying the following conditions.*

3. $\bigtriangledown(P \oplus Q) <^1_1 \bigtriangledown P \triangledown_n \bigtriangledown Q$ *for each* $n \in \mathbb{N}$.
4. $\widehat{\mathrm{Deg}}(P \oplus Q) <^1_1 \widehat{\mathrm{Deg}}(P) \triangledown_\omega \widehat{\mathrm{Deg}}(Q)$

Proof. (1) We first construct a computable function Φ which witnesses $\bigtriangledown(P \oplus Q) \leq^1_1 \bigtriangledown P \triangledown_\omega \bigtriangledown Q$. Set $\Phi(\langle\rangle) = \langle\rangle$. Fix $\sigma = \sigma^{--}{}^\frown\langle(i, m), (j, n)\rangle \in (2 \times (\mathbb{N} \cup \{\sharp\}))^{<\mathbb{N}}$. If $i = j$, then set $\Phi(\sigma) = \Phi(\sigma^-)^\frown\langle n\rangle$. If $i \neq j$, then set $\Phi(\sigma) = \Phi(\sigma^-)^\frown\langle\sharp\rangle^\frown\langle j\rangle^\frown\mathtt{tail}(\mathtt{pr}_j(\sigma))$. Note that $\Phi(g)$ contains only finitely many $\sharp$'s for any $g \in \bigtriangledown P \triangledown_\omega \bigtriangledown Q$, by the definition of $\triangledown_\omega$. Thus, for any $g \in \bigtriangledown P \triangledown_\omega \bigtriangledown Q$, we have $\mathtt{tail}(\Phi(g)) = i^\frown\mathtt{tail}(\mathtt{pr}_i(g))$ for some $i < 2$. Hence, $\Phi(g) \in \bigtriangledown(P \oplus Q)$.

We next construct a computable function Φ witnessing $\bigtriangledown(P \oplus Q) \geq^1_1 \bigtriangledown P \triangledown_\omega \bigtriangledown Q$. Set $\Phi(\langle\rangle) = \langle\rangle$. Fix $\sigma \in (\mathbb{N} \cup \{\sharp\})^{<\mathbb{N}}$. If $\sigma = \sigma^{--}{}^\frown\langle m, n\rangle$ for some $m, n \in \mathbb{N}$ and $\mathtt{tail}(\sigma; 0)$ is defined, then $\Phi(\sigma) = \Phi(\sigma^-)^\frown\langle(\mathtt{tail}(\sigma; 0), n)\rangle$. If $\sigma = \sigma^{--}{}^\frown\langle\sharp, i, n\rangle$ for some $i < 2$ and $n \in \mathbb{N}$, then $\Phi(\sigma) =$

$\Phi(\sigma^-)^\frown\langle(0,\sharp),(1,\sharp),(i,n)\rangle$. Otherwise, put $\Phi(\sigma)=\Phi(\sigma^-)$. Note that $\texttt{tail}(g)$ converges for any $g\in\bigtriangledown(P\oplus Q)$, and hence, $\texttt{pr}_{\texttt{tail}(g;0)}(\Phi(g))$ is total. Therefore, $\texttt{tail}(g)=\texttt{tail}(g;0)^\frown\texttt{tail}(\texttt{pr}_{\texttt{tail}(g;0)}(\Phi(g)))$. Thus, $\Phi(g)\in\bigtriangledown P\triangledown_\omega\bigtriangledown Q$.

(2) Clearly $\widehat{\mathrm{Deg}}(P\oplus Q)=P\cup Q$, and recall that $P\cup Q\equiv^1_1[\![P\vee Q]\!]^1_{\mathsf{CL}}\leq^1_1[\![P\vee Q]\!]^2_{\mathsf{CL}}$. On the other hand, $P\triangledown_\infty Q\subseteq\widehat{\mathrm{Deg}}(P)\cup\widehat{\mathrm{Deg}}(Q)$.

(3) Let f_0 and f_1 be Turing incompatable elements of $\mathbb{N}^\mathbb{N}$. Fix $n\in\mathbb{N}$. Assume that $\bigtriangledown\{f_0\}\triangledown_n\bigtriangledown\{f_1\}\leq^1_1\bigtriangledown\{f_0\}\triangledown_\omega\bigtriangledown\{f_1\}$ holds via a computable function Φ. We first set $g_0=0^\mathbb{N}\oplus f_0\in\bigtriangledown\{f_0\}\triangledown_\omega\bigtriangledown\{f_1\}$. Then $\texttt{pr}_0(\Phi(g_0))$ is total, since $\texttt{pr}_1(\Phi(g_0))\in\bigtriangledown\{f_1\}$ implies $f_0\equiv_T g_0\geq_T\Phi(g_0)\geq_T f_1$ that contradicts our assumption. In particular, $(\Phi(g_0\upharpoonright s))_0=0$ for some $s\in\mathbb{N}$. Then we extend $g_0\upharpoonright s$ to $g_1=(g_0\upharpoonright s)^\frown(1^\mathbb{N}\oplus f_1)\in\bigtriangledown\{f_0\}\triangledown_\omega\bigtriangledown\{f_1\}$. Note that $g_1\equiv_T f_1$. By the same argument, $(\Phi(g_1\upharpoonright t))_0=1$ for some $t\in\mathbb{N}$. Therefore, we can ensure $\texttt{mc}(\Phi(g_1))\geq 1$. By iterating this procedure, we can easily see that $\texttt{mc}(\Phi(g_{n+1}))>n$ for some $g_{n+1}\in\bigtriangledown\{f_0\}\triangledown_\omega\bigtriangledown\{f_1\}$.

(4) Let f_0 and f_1 be Turing incompatable elements of $\mathbb{N}^\mathbb{N}$. Fix $n\in\mathbb{N}$, and assume that the condition $\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\omega\widehat{\mathrm{Deg}}(\{f_1\})\leq^1_1\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\infty\widehat{\mathrm{Deg}}(\{f_1\})$ holds via a computable function Φ. We first set $g_0=0^\mathbb{N}\oplus f_0\in\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\infty\widehat{\mathrm{Deg}}(\{f_1\})$. Then $\texttt{pr}_1(\Phi(g_0))$ is not total, since $\texttt{pr}_1(\Phi(g_0))\in\widehat{\mathrm{Deg}}(\{f_1\})$ implies $f_0\equiv_T g_0\geq_T\Phi(g_0)\geq_T f_1$ that contradicts our assumption. In particular, $(\Phi(g_0\upharpoonright s_0))_0=0$ for some $s_0\in\mathbb{N}$. Then we extend $g_0\upharpoonright s$ to $g_1=(g_0\upharpoonright s)^\frown(1^\mathbb{N}\oplus f_1)\in\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\infty\widehat{\mathrm{Deg}}(\{f_1\})$. By the same argument, $(\Phi(g_1\upharpoonright s_1))_0=1$ for some $s_1>s_0$. Therefore, we can ensure $\texttt{mc}(\Phi(g_1))\geq 1$. Assume that g_{2n+1} and s_{2n+1} have been already constructed, $\texttt{mc}(\Phi(g_{2n+1}))\geq 2n+1$, $\texttt{pr}_0(g_{2n+1})$ is an initial segment of f_0, $\texttt{pr}_1(g_{2n+1})=f_1$, and $(\Phi(g_{2n+1}\upharpoonright s_{2n+1}))_0=1$. Then we extend $g_{2n+1}\upharpoonright s_{2n+1}$ to $g_{2n+2}=(g_0\upharpoonright s_{2n+1})^\frown(0^\mathbb{N}\oplus h)$, where $\texttt{pr}_0(g_{2n+2})^\frown h=f_0$. Again we have $(\Phi(g_{2n+2}\upharpoonright s_{2n+2}))_0=0$ for some $s_{2n+2}>s_{2n+1}$, and $\texttt{mc}(\Phi(g_{2n+2}))\geq 2n+2$. We extend $g_{2n+2}\upharpoonright s_{2n+2}$ to $g_{2n+3}=(g_0\upharpoonright s_{2n+2})^\frown(1^\mathbb{N}\oplus h^*)$, where $\texttt{pr}_1(g_{2n+3})^\frown h^*=f_1$. We have $(\Phi(g_{2n+3}\upharpoonright s_{2n+3}))_0=0$ for some $s_{2n+3}>s_{2n+2}$, and $\texttt{mc}(\Phi(g_{2n+3}))\geq 2n+3$. Finally, we set $g=\bigcup_{n\in\mathbb{N}}g_n$. We can easily see that $\texttt{pr}_i(g)=f_i$ for each $i<2$, and $\texttt{mc}(\Phi(g))=\infty$. Hence, $g\in\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\infty\widehat{\mathrm{Deg}}(\{f_1\})$, but $\Phi(g)\notin\widehat{\mathrm{Deg}}(\{f_0\})\triangledown_\omega\widehat{\mathrm{Deg}}(\{f_1\})$. □

5.2. **Diagonally Noncomputable Functions.** A total function $f:\mathbb{N}\to\mathbb{N}$ is a *k-valued diagonally noncomputable function* if $f(n)<k$ for any $n\in\mathbb{N}$ and $f(e)\neq\Phi_e(e)$ whenever $\Phi_e(e)$ converges. Let DNR_k denote the set of all k-valued diagonally noncomputable functions. Jockusch [43] showed that every DNR_k function computes a DNR_2 function. However, he also noted that there is no uniformly computable algorithm finding a DNR_2 function from any DNR_k function.

Theorem 164 (Jockusch [43])**.**

(1) $\mathsf{DNR}_k>^1_1\mathsf{DNR}_{k+1}$ *for any* $k\in\mathbb{N}$.
(2) $\mathsf{DNR}_2\equiv^\omega_1\mathsf{DNR}_k$ *for any* $k\in\mathbb{N}$.

Proposition 165.

(1) *If a* $(1,\omega)$*-degree* $\mathbf{d}^1_\omega$ *of subsets of* $\mathbb{N}^\mathbb{N}$ *contains a* $(1,1)$*-degree* $\mathbf{h}^1_1$ *of homogeneous sets, then* $\mathbf{h}^1_1$ *is the greatest* $(1,1)$*-degree inside* $\mathbf{d}^1_\omega$.
(2) *If an* $(<\omega,1)$*-degree* $\mathbf{d}^{<\omega}_1$ *of* Π^0_1 *subsets of* $2^\mathbb{N}$ *contains a* $(1,<\omega)$*-degree* $\mathbf{h}^1_{<\omega}$ *of homogeneous* Π^0_1 *sets, then* $\mathbf{h}^1_{<\omega}$ *is the least* $(1,<\omega)$*-degree inside* $\mathbf{d}^{<\omega}_1$.
(3) *Every* $(<\omega,1)$*-degree of* Π^0_1 *subsets of* $2^\mathbb{N}$ *contains at most one* $(1,1)$*-degree of homogeneous* Π^0_1 *sets.*

Proof. For the item 1, we can see that, for any $P\subseteq\mathbb{N}^\mathbb{N}$ and any closed set $Q\subseteq\mathbb{N}^\mathbb{N}$, if $P\leq^1_\omega Q$ then there is a node σ such that $Q\cap[\sigma]$ is nonempty and $P\leq^1_1 Q\cap[\sigma]$. That is, σ is a locking sequence. If Q is homogeneous, then $P\leq^1_1 Q\equiv^1_1 Q\cap[\sigma]$. The item 2 follows from Theorem 72. By combining the item 1 and 2, we see that every $(<\omega,1)$-degree of Π^0_1 subsets of $2^\mathbb{N}$ contains at most one $(1,<\omega)$-degree of homogeneous Π^0_1 sets which contains at most $(1,1)$-degree of homogeneous Π^0_1 sets. □

Corollary 166. $\mathsf{DNR}_3 <^{<\omega}_1 \mathsf{DNR}_2$, *and* $\mathsf{DNR}_3 <^1_\omega \mathsf{DNR}_2$.

Proof. By Jockusch [43], we have $\mathsf{DNR}_3 <^1_1 \mathsf{DNR}_2$. Thus, Proposition 165 implies the desired condition. □

By analyzing Jockusch's proof [43] of the Muchnik equivalence of DNR_2 and DNR_k for any $k \geq 2$, we can directly establish the $(<\omega,\omega)$-equivalence of DNR_2 and DNR_k for any $k \geq 2$. However, one may find that Jockusch's proof [43] is essentially based on the Σ^0_2 law of excluded middle. Therefore, the fine analysis of this proof structure establishes the following theorem.

Theorem 167. $\mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k <^1_1 \mathsf{DNR}_{k^2}$ *for any* k.

Proof. As Jockusch [43], fix a computable function $z : \mathbb{N}^2 \to \mathbb{N}$ such that $\Phi_{z(v,u)}(z(v,u)) = \langle \Phi_v(v), \Phi_u(u)\rangle$ for any $v, u \in \mathbb{N}$. Note that every $g \in (k^2)^{\mathbb{N}}$ determines two functions $g_0 \in k^{\mathbb{N}}$ and $g_1 \in k^{\mathbb{N}}$ such that $g(n) = \langle g_0(n), g_1(n)\rangle$ for any $n \in \mathbb{N}$. We define a uniform sequence $\{\Gamma_v\}_{v\in\mathbb{N}}, \Delta$ of computable functions as $\Gamma_v(g;u) = g_1(z(v,u))$, and $\Delta(g;v) = g_0(z(v,u_v))$, where $u_v = \min\{u \in \mathbb{N} : g_1(z(v,u)) = \Phi_u(u)\downarrow\}$. Fix $g \in \mathsf{DNR}_{k^2}$. Since $\langle g_0(z(v,u)), g_1(z(v,u))\rangle = g(z(v,u)) \neq \langle \Phi_v(v), \Phi_u(u)\rangle$, either $g_0(z(v,u)) \neq \Phi_v(v)$ or $g_1(z(v,u)) \neq \Phi_u(u)$ holds for any $v, u \in \mathbb{N}$. We consider the following Σ^0_2 sentence:

$$(\exists v)(\forall u)\ (\Phi_u(u)\downarrow \to\ g_1(z(v,u)) \neq \Phi_u(u)).$$

Let $\theta(g,v)$ denote the Π^0_1 sentence $(\forall u)\ (\Phi_u(u)\downarrow \to\ g_1(z(v,u)) \neq \Phi_u(u))$. If $\theta(g,v)$ holds, then $\Gamma_v(g;u) = g_1(z(v,u)) \neq \Phi_u(u)$ for any $u \in \mathbb{N}$. Hence, $\Gamma_v(g) \in \mathsf{DNR}_k$. If $\neg\theta(g,v)$ holds, then u_v is defined. Therefore, $\Delta(g;v) = g_0(z(v,u_v))\downarrow\neq \Phi_v(v)$, since $g_1(z(v,u_v)) = \Phi_{u_v}(u_v))\downarrow$. Thus, $\Delta(g;v)$ is extendible to a function in DNR_k. Consequently, $\mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k \leq^1_1 \mathsf{DNR}_{k^2}$ by Theorem 161 (2).

To see $\mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k \not\geq^1_1 \mathsf{DNR}_{k^2}$, we note that $\mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k$ is not tree-immune. By Cenzer-Kihara-Weber-Wu [19], $\mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k$ does not cup to the generalized separating class DNR_{k^2}. □

Corollary 168. $\mathsf{DNR}_k \equiv^{<\omega}_\omega \mathsf{DNR}_2$ *for any* $k \geq 2$. *Indeed, for any* $k \in \mathbb{N}$, *the direction* $\mathsf{DNR}_k \leq^{<\omega}_\omega \mathsf{DNR}_{k^2}$ *is witnessed by a team of a confident learner and a eventually-Popperian learner. In particular,* $\mathsf{DNR}_k \equiv^\omega_1 \mathsf{DNR}_2$ *for any* $k \geq 2$.

Proof. By Proposition 134, $\mathsf{DNR}_k \leq^{<\omega}_\omega \mathsf{DNR}_k \blacktriangledown \mathsf{DNR}_k$ is witnessed by a team of a confident learner and a eventually-Popperian learner. Thus, Theorem 167 implies the desired condition. □

Corollary 169. *There is an* $(<\omega,\omega)$*-degree which contains infinitely many* $(1,1)$*-degrees of homogeneous* Π^0_1 *sets.*

Proof. By Corollary 168, the $(<\omega,\omega)$-degree of DNR_2 contains DNR_k for any $k \in \mathbb{N}$, while $\mathsf{DNR}_k \not\equiv^1_1 \mathsf{DNR}_l$ for $k \neq l$. □

For a pointclass Γ in a space X, we say that $\mathrm{SEL}(\Gamma, X)$ *holds for* (α,β)*-degrees* if, for every Γ set $S \subseteq X$ and for every nonempty Π^0_1 set $Q \subseteq 2^{\mathbb{N}}$, there exists a Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ such that $P \equiv^\alpha_\beta S \cup Q$. Jockusch-Soare [44] indicates that $\mathrm{SEL}(\Pi^0_2, \mathbb{N}^{\mathbb{N}})$ holds for $(\omega,1)$-degrees, and points out that $\mathrm{SEL}(\Pi^0_3, 2^{\mathbb{N}})$ does not hold for $(\omega,1)$-degrees, since the set of all noncomputable elements in $2^{\mathbb{N}}$ is Π^0_3. Simpson's Embedding Lemma [62] determines the limit of $\mathrm{SEL}(\Gamma, X)$ for $(\omega,1)$-degrees.

Theorem 170 (Simpson [62])**.** $\mathrm{SEL}(\Sigma^0_3, \mathbb{N}^{\mathbb{N}})$ *holds for* $(\omega,1)$*-degrees.* □

Theorem 171 (Simpson's Embedding Lemma for other degree structures)**.**

1. $\mathrm{SEL}(\Sigma^0_2, 2^{\mathbb{N}})$ *does not hold for* $(<\omega,1)$*-degrees.*
2. $\mathrm{SEL}(\Sigma^0_2, 2^{\mathbb{N}})$ *holds for* $(1,\omega)$*-degrees.*
3. $\mathrm{SEL}(\Pi^0_2, 2^{\mathbb{N}})$ *does not hold for* $(1,\omega)$*-degrees.*
4. $\mathrm{SEL}(\Pi^0_2, \mathbb{N}^{\mathbb{N}})$ *holds for* $(<\omega,\omega)$*-degrees.*
5. $\mathrm{SEL}(\Sigma^0_3, 2^{\mathbb{N}})$ *does not hold for* $(<\omega,\omega)$*-degrees.*

Proof. (1) For any Π^0_1 set $P \subseteq 2^{\mathbb{N}}$, we note that $\bigtriangledown P \subseteq 2^{\mathbb{N}}$ is Σ^0_2. By Theorem 117, there is no Π^0_1 set $2^{\mathbb{N}}$ which is $(<\omega, 1)$-below $\bigtriangledown P$. In particular, there is no Π^0_1 set $2^{\mathbb{N}}$ which is $(<\omega, 1)$-equivalent to $P \cup \bigtriangledown P = \bigtriangledown P$.

(2) For a given Σ^0_2 set $S \subseteq 2^{\mathbb{N}}$, there is a computable increasing sequence $\{P_i\}_{i\in\mathbb{N}}$ of Π^0_1 classes such that $S = \bigcup_{i\in\mathbb{N}} P_i$. We need to show $\bigcup_{i\in\mathbb{N}} P_i \equiv^1_\omega \bigoplus_{i\in\mathbb{N}} P_i$, since $\bigoplus_{i\in\mathbb{N}} P_i$ is $(1, <\omega)$-equivalent to the Π^0_1 class $\bigoplus^{\longrightarrow}_i P_i$. Then, it is easy to see $\bigcup_i P_i \leq^1_1 \bigoplus_i P_i$. For given $f \in \bigcup_i P_i$, from each initial segment $f \upharpoonright n$, a learner Ψ guesses an index of a computable function $\Phi_{\Psi(f\upharpoonright n)}(g) = i^\frown g$ for the least number i such that $f \upharpoonright n \in T_{P_i}$ but $f \upharpoonright n \notin T_{P_{i-1}}$. For any $f \in \bigcup_i P_i$, for the least i such that $f \in P_i \setminus P_{i-1}$, $\lim_n \Psi(f \upharpoonright n)$ converges to an index of $\Phi_{\lim_n \Psi(f\upharpoonright n)}(g) = i^\frown g$. Thus, $\Phi_{\lim_n \Psi(f\upharpoonright n)}(g) \in i^\frown P_i$. Consequently, $S = \bigcup_i P_i \leq^1_\omega \bigoplus^{\longrightarrow}_i P_i$.

(3) Fix any special Π^0_1 set $P \subseteq 2^{\mathbb{N}}$. By Jockusch-Soare [44], there is a noncomputable Σ^0_1 set $A \subseteq \mathbb{N}$ such that P has no A-computable element. Then $\{A\} \subseteq 2^{\mathbb{N}}$ is a Π^0_2 singleton, since A is Σ^0_1. Therefore, $P \oplus \{A\}$ is Π^0_2. It suffices to show that there is no Π^0_1 set $Q \subseteq 2^{\mathbb{N}}$ such that $Q \equiv^1_\omega P \oplus \{A\}$. Assume that $Q \equiv^1_\omega P \oplus \{A\}$ is satisfied for some Π^0_1 set $Q \subseteq 2^{\mathbb{N}}$. Then Q must have an A-computable element $\alpha \in Q$. Fix a learner Ψ witnessing $P \oplus \{A\} \leq^1_\omega Q$. Then, we have $\Phi_{\lim_n \Psi(\alpha\upharpoonright n)}(\alpha) = 1^\frown A$, since P has no element computable in $\alpha \leq_T A$. We wait for $s \in \mathbb{N}$ such that $\Psi(\alpha \upharpoonright t) = \Psi(\alpha \upharpoonright s)$ for any $t \geq s$. Then, fix $u \geq s$ with $\Phi_{\Psi(\alpha\upharpoonright u)}(\alpha \upharpoonright u; 0) \downarrow = 1$. Consider the Π^0_1 set $Q^* = \{f \in Q \cap [\alpha \upharpoonright u] : (\forall v \geq u)\ \Psi(f \upharpoonright v) = \Psi(f \upharpoonright u)\}$. Then, for any $f \in Q^*$, $\Phi_{\lim_s \Psi(f\upharpoonright s)}(f) = \Phi_{\Psi(\alpha\upharpoonright u)}(f)$ must extends $\langle 1 \rangle$. Thus, $\{1^\frown A\} \leq^1_1 Q^*$ via the computable function $\Phi_{\Psi(\alpha\upharpoonright u)}$. Since Q^* is special Π^0_1 subset of $2^{\mathbb{N}}$, this implies the computability of $1^\frown A$ which contradicts our choice of A.

(4) Fix a Π^0_2 set $S \subseteq \mathbb{N}^{\mathbb{N}}$. As Simpson's proof, there is a Π^0_1 set $\widehat{S} \subseteq \mathbb{N}^{\mathbb{N}}$ such that $S \equiv^1_1 \widehat{S}$. We can find a Π^0_1 set $\widehat{P} \subseteq \widehat{S} \blacktriangledown Q$ such that $\widehat{P} \leq^1_1 S \cup Q$, and $\widehat{P}$ is computably homeomorphic to a Π^0_1 set $P \subseteq 2^{\mathbb{N}}$. Since $S \cup Q \leq^{<\omega}_\omega \widehat{S} \blacktriangledown Q$, we have $S \cup Q \equiv^{<\omega}_\omega P$.

(5) For every Π^0_1 set $P \subseteq 2^{\mathbb{N}}$, the Turing upward closure $\widehat{\mathrm{Deg}}(P) = \{g \in 2^{\mathbb{N}} : (\exists f \in P)\ f \leq_T g\}$ of P is Σ^0_3, and $\widehat{\mathrm{Deg}}(P)$ has the least $(<\omega, \omega)$-degree inside $\deg^\omega_1(P)$. By Theorem 149, there is no Π^0_1 subset of $2^{\mathbb{N}}$ which is $(<\omega, \omega)$-equivalent to $\widehat{\mathrm{Deg}}(P)$. □

5.3. **Weihrauch Degrees and Constructive Principles.** We can also give a characterization of our nonuniformly computable functions in the context of the Weihrauch degrees which is a generalization of the Medvedev degrees. Then, our results could be translated into the results on the Weihrauch degrees. A partial function $P :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathcal{P}(\mathbb{N}^{\mathbb{N}})$ is called *a multi-valued function*. Then P is also written as $P : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$. One can think of each multi-valued function P as a collection $\{P(x)\}_{x\in\mathrm{dom}(P)}$ of mass problems $P(x) \subseteq \mathbb{N}^{\mathbb{N}}$, or a Π_2-*theorem* $(\forall x \in \mathrm{dom}(P))(\exists y)\ y \in P(x)$.

Definition 172 ([11, 13, 12, 14])**.** Let $P :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ and $Q :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be multi-valued partial functions.

1. A single-valued function $q :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is said to be *a realizer of* Q if $q(x) \in Q(x)$ for any $x \in \mathrm{dom}(Q)$.
2. We say that P *is Weihrauch reducible to* Q (written $P \leq_W Q$) if there are partial computable functions H, K such that $K(x, q \circ H(x))$ for any $x \in \mathrm{dom}(P)$ and any realizer q of Q.

Remark. If we think of the values $P(x)$ and $Q(x)$ as *relativized mass problems* P^x and Q^x, then $P \leq_W Q$ can be represented as the existence of partial computable functions $\Phi, \Delta :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ satisfying $\Phi^x : Q^{\Delta(x)} \to P^x$ for any $x \in \mathrm{dom}(Q)$, where Φ^x is the x-computable function mapping $y \in \mathbb{N}^{\mathbb{N}}$ to $\Phi(x \oplus y)$. For any subset P of Baire space $\mathbb{N}^{\mathbb{N}}$, we define $\iota(P) : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ by $\iota(P)(x) = P$ for any $x \in \mathbb{N}^{\mathbb{N}}$. Then, the map ι provides an embedding of the Medvedev degrees into the Weihrauch degrees, i.e., $P \leq^1_1 Q$ if and only if $\iota(P) \leq_W \iota(Q)$.

Definition 173 ([11, 13, 12, 14])**.** Let $P, Q :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be partial multi-valued functions.

(1) (Pairing) $\langle P, Q\rangle(x) = P(x) \times Q(x)$.
(2) (Product) $(P \times Q)(\langle x, y\rangle) = P(x) \times Q(y)$.
(3) (Coproduct) $(P \coprod Q)(0, x) = \{0\} \times P(x)$; and $(P \coprod Q)(1, x) = \{1\} \times Q(x)$.
(4) (Composition) $(P \circ Q)(x) = \bigcup\{P(y) : y \in Q(x)\}$.
(5) (Parallelization) $\widehat{P}(\langle x_i : i \in \mathbb{N}\rangle) = \prod_{i\in\mathbb{N}} P(x_i)$.

Let X be a computable metric space (for definition, see Weihrauch [75]). Then, $\mathcal{A}_-(X)$ denotes the hyperspace of closed subsets of X with the upper Fell representation ψ_- (see [11]). For example, P is a computable point in the hyperspace $\mathcal{A}_-(\mathbb{N}^\mathbb{N})$ (resp. $\mathcal{A}_-(2^\mathbb{N})$) if and only if P is a Π^0_1 subset of Baire space $\mathbb{N}^\mathbb{N}$ (resp. of Cantor space $2^\mathbb{N}$).

Definition 174 (Closed Choice [11, 13, 12, 14])**.** Let X be a computable metric space. Then, the *closed choice* operation of X is defined as the following partial function.

$$\mathsf{C}_X :\subseteq \mathcal{A}_-(X) \rightrightarrows X, \quad A \mapsto A$$

Here, $\mathrm{dom}(\mathsf{C}_X) = \{A \in \mathcal{A}_-(X) : A \neq \emptyset\}$.

Remark. One may think of the closed choice $\mathsf{C}_{2^\mathbb{N}}$ of Cantor space as the following (relativizable) mass problem: Let $\{P^x_e\}_{e\in\mathbb{N}}$ be a computable enumeration of all nonempty $\Pi^0_1(x)$ subsets of Cantor space $2^\mathbb{N}$, uniformly in $x \in \mathbb{N}^\mathbb{N}$. The $\forall$-player chooses some $(e, x) \in \mathbb{N}\times\mathbb{N}^\mathbb{N}$ which codes an instance of the problem $\mathsf{C}_{2^\mathbb{N}}$. Then the $\exists$-player receives (e, x), and then she is required to solve the relativized mass problem P^x_e. The $\exists$-player wins if she succeeds to choose an element of the e-th $\Pi^0_1(x)$ set $P^x_e \subseteq 2^\mathbb{N}$, where she can use the information of the choice (e, x) of the $\forall$-player. Otherwise, the $\forall$-player wins. Intuitively, the Weihrauch degree of $\mathsf{C}_{2^\mathbb{N}}$ is the degree of difficulty to computing a winning strategy for the $\exists$-player. It is not hard to see that $\mathsf{C}_{2^\mathbb{N}} \equiv_W \mathsf{C}_{[0,1]} \equiv_W \mathsf{WKL}$, where WKL denotes the multi-valued function representing Weak König's Lemma.

Definition 175. We introduce non-intuitinistic principles as the following partial multi-valued functions on Baire space.

(1) (The *law of excluded middle for Σ^0_1 formulas*)

$$\Sigma^0_1\text{-}\mathsf{LEM} : \mathbb{N}^\mathbb{N} \to 2, \qquad \Sigma^0_1\text{-}\mathsf{LEM}(p) = \begin{cases} 0, & \text{if } (\exists n \in \mathbb{N})\ p(n) = 0, \\ 1, & \text{otherwise.} \end{cases}$$

(2) (The *law of excluded middle for Δ^0_2 formulas*)

$$\Delta^0_2\text{-}\mathsf{LEM} :\subseteq \mathbb{N}^2 \times \mathbb{N}^\mathbb{N} \to 2, \qquad \Delta^0_2\text{-}\mathsf{LEM}(i, j, p) = \begin{cases} 0, & \text{if } p \in \mathrm{Tot}_i, \\ 1, & \text{otherwise.} \end{cases}$$

Here, $\mathrm{dom}(\Delta^0_2\text{-}\mathsf{LEM}) = \{(i, j, p) \in \mathbb{N}^2 \times \mathbb{N}^\mathbb{N} : \mathrm{Tot}_i = \mathbb{N}^\mathbb{N} \setminus \mathrm{Tot}_j\}$, where Tot_e denotes the set of all oracles $\alpha \in \mathbb{N}^\mathbb{N}$ such that $\Phi_e(\alpha; n)$ converges for all inputs $n \in \mathbb{N}$.

(3) (The *lessor limited principle of omniscience for Σ^0_2 formulas*)

$$\Sigma^0_2\text{-}\mathsf{LLPO}_\alpha :\subseteq (\mathbb{N}^\mathbb{N})^\alpha \rightrightarrows \alpha, \quad (x_i)_{i<\alpha} \mapsto \{i < \alpha : (\exists^\infty n \in \mathbb{N})\ x_i(n) = 0\}.$$

Here, $\mathrm{dom}(\Sigma^0_2\text{-}\mathsf{LLPO}_\alpha) = \{(x_i)_{i<\alpha} \in (\mathbb{N}^\mathbb{N})^\alpha : (\exists i < \alpha)(\exists^\infty n \in \mathbb{N})\ x_i(n) = 0\}$. Define $\Sigma^0_2\text{-}\mathsf{LLPO} = \Sigma^0_2\text{-}\mathsf{LLPO}_2$.

(4) (The *double negation elimination for Σ^0_2 formulas*)

$$\Sigma^0_2\text{-}\mathsf{DNE} :\subseteq \mathbb{N}^\mathbb{N} \rightrightarrows \mathbb{N}, \quad x \mapsto \{m \in \mathbb{N} : (\forall n > m)\ x(n) \neq 0\}.$$

Here, $\mathrm{dom}(\Sigma^0_2\text{-}\mathsf{DNE}) = \{x \in \mathbb{N}^\mathbb{N} : (\exists m \in \mathbb{N})(\forall n > m)\ x(n) \neq 0\}$.

(5) (The *law of excluded middle for Σ^0_2 formulas*)

$$\Sigma^0_2\text{-}\mathsf{LEM} : \mathbb{N}^{\mathbb{N}} \rightrightarrows 2 \times \mathbb{N}, \qquad \Sigma^0_2\text{-}\mathsf{LEM}(p) \ni \begin{cases} (0, s), & \text{if } (\forall m \in \mathbb{N})(\exists n > m)\ p(n) = 0, \\ (1, s), & \text{if } (\forall n > s)\ p(n) \neq 0. \end{cases}$$

(6) (The *lessor limited principle of omniscience for* $(\Sigma^0_2 \wedge \Pi^0_2)$ *formulas*)

$$(\Sigma^0_2)_2\text{-}\mathsf{LLPO}_\alpha : (x_i)_{i<\alpha} \mapsto \{(i, m) : (\exists^\infty n \in \mathbb{N})\ x_i(n) = 0\ \&\ (\forall n > m)\ x_i(n) \neq 1\}.$$

Here, $\mathrm{dom}((\Sigma^0_2)_2\text{-}\mathsf{LLPO}_\alpha) = \{(x_i)_{i<\alpha} \in (\mathbb{N}^{\mathbb{N}})^\alpha : (\exists i < \alpha)[(\exists^\infty n \in \mathbb{N})\ x_i(n) = 0\ \&\ (\neg\exists^\infty n \in \mathbb{N})\ x_i(n) = 1]\}$. Define $(\Sigma^0_2)_2\text{-}\mathsf{LLPO} = (\Sigma^0_2)_2\text{-}\mathsf{LLPO}_2$.

(7) (The *double negation elimination for Σ^0_3 formulas*)

$$\Sigma^0_3\text{-}\mathsf{DNE} : x \mapsto \{k : (\forall m \in \mathbb{N})(\exists n \geq m)\ x(\langle k, n\rangle) = 0\}.$$

Here, $\mathrm{dom}(\Sigma^0_3\text{-}\mathsf{DNE}) = \{x \in \mathbb{N}^{\mathbb{N}} : (\exists k \in \mathbb{N})(\forall m \in \mathbb{N})(\exists n \geq m)\ x(\langle k, n\rangle) = 0\}$.

Remark. (1) The single-valued function Σ^0_1-LEM is usually called *the limited principle of omniscience* (LPO). Brattka-de Brecht-Pauly [11] showed that a single-valued partial function $f :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is $(1, \omega)$-computable if and only if f is Weihrauch reducible to the closed choice $\mathsf{C}_{\mathbb{N}}$ for the discrete space $\mathbb{N}$. Here, in their term, the $(1, \omega)$-computability is called *the computability with finitely many mind changes*. It is not hard to see that the closed choice $\mathsf{C}_{\mathbb{N}}$ is Weihrauch equivalent to $\bigcup_{m\in\mathbb{N}} \Sigma^0_1\text{-}\mathsf{LEM}^m$, where $f^m : X^m \rightrightarrows Y^m$ is the m-fold product of $f : X \rightrightarrows Y$. We also note that $\Sigma^0_1\text{-}\mathsf{LEM}^m$ is Weihrauch equivalent to the Σ^0_1 *least number principle with the upper bound* $m + 1$, $\Sigma^0_1\text{-}\mathsf{LNP}_{m+1} : \mathbb{N}^{\mathbb{N}} \to m + 1$, which maps p to $\min(\{m\} \cup \{p(n) : n \in \mathbb{N}\})$. To see $\Sigma^0_1\text{-}\mathsf{LEM}^m \leq_W \Sigma^0_1\text{-}\mathsf{LNP}_{m+1}$, we consider two computable functions $H : (\mathbb{N}^{\mathbb{N}})^m \to (m + 1)^{\mathbb{N}}$ and $K : (\mathbb{N}^{\mathbb{N}})^m \times (m + 1) \to 2^m$ defined as $H(\langle x_i\rangle_{i<m})(n) = \#\{i < m : (\forall m \leq n)\ x_i(m) \neq 0\}$ and $K(\langle x_i\rangle_{i<m}, k)(j) = \min(\{1\} \cup \{x_j(m) : m \leq s_k\})$, where $s_k = \min\{s \in \mathbb{N} : H(\langle x_i\rangle_{i<m})(s) = k\}$, and another direction is trivial.

(2) $\Sigma^0_2\text{-}\mathsf{LLPO}_2$ is Weihrauch equivalent to the *jump* LLPO' *of* LLPO in the sense of Brattka-Gherardi-Marcone [14]. They also showed that LLPO' is Weihrauch equivalent to the Borzano-Weierstrass Theorem BWT_2 for the discrete space $\{0, 1\}$. We also note that $\Sigma^0_2\text{-}\mathsf{LLPO}_k$ is equivalent to the Infinite Pigeonhole Principle for k-partitions. Brattka-Gherardi-Marcone [14] also pointed out that the n-th jump of LLPO and LPO correspond to Σ^0_{n+1}-LLPO (that is, the lessor limited principle of omniscience for Σ^0_{n+1}-formulas) and Σ^0_{n+1}-LEM (the law of excluded middle for Σ^0_{n+1}-formulas), respectively.

(3) The *Dirichlet function* $\mathbf{1}_{\mathbb{Q}} : \mathbb{R} \to \{0, 1\}$, the characteristic function of the rational numbers $\mathbb{Q}$, is Weihrauch reducible to Σ^0_2-LEM. Indeed, it is easy to see that the Dirichlet function $\mathbf{1}_{\mathbb{Q}}$ is Weihrauch equivalent to the single-valued function $\Pi^0_2\text{-}\mathsf{LEM} : \mathbb{N}^{\mathbb{N}} \to 2$, where $\Pi^0_2\text{-}\mathsf{LEM}(p) = 0$, if $p(n) = 0$ for infinitely many $n \in \mathbb{N}$; and $\Pi^0_2\text{-}\mathsf{LEM}(p) = 1$, otherwise. It is well-known that the Dirichlet function $\mathbf{1}_{\mathbb{Q}}$ is of Baire class 2, but not of Baire class 1 (i.e., $\mathbf{1}_{\mathbb{Q}}$ is $\mathbf{\Sigma}^0_3$-measurable but not $\mathbf{\Sigma}^0_2$-measurable).

(4) The study of arithmetical hierarchy of semiclassical principles such as Σ^0_n-LEM, Σ^0_n-LLPO, and Σ^0_n-DNE was initiated by Akama-Berardi-Hayashi-Kohlenbach [1]. In particular, on the study of the second level of arithmetical hierarhcy for semiclassical principles, see also Berardi [4] and Toftdal [72]. The relationship between the learnability and Σ^0_2-DNE has been also studied by Nakata-Hayashi [54] in the context of a realizability interpretation of limit computable mathematics.

Definition 176 (Unique variant [14])**.** Let $P : X \rightrightarrows Y$ be a multi-valued function. Then $\mathsf{Unique}P : X \rightrightarrows Y$ is defined as the restriction of P up to $\mathrm{dom}(\mathsf{Unique}P) = \{x \in \mathrm{dom}(P) : \#P(x) = 1\}$.

Proposition 177. *$\Delta^0_2\text{-}\mathsf{LEM}$ is Weihrauch reducible to $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}$.*

Proof. To see $\Delta^0_2\text{-}\mathsf{LEM} \leq_W \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}$, given $(e_0, e_1, p) \in \mathbb{N}^2 \times \mathbb{N}^\mathbb{N}$, define $H(e_0, e_1, p)$ to be a pair (x_0, x_1), where $x_i(s) = 0$ if and only if the computation $\Phi_{e_i,s+1}(p)$ at stage $s + 1$ properly extends $\Phi_{e_i,s}(p)$ at the previous stage. Then x_i contains infinitely many 0's if and only if p is contained in Tot_{e_i}. Note that, whenever (e_0, e_1, p) is contained in the domain of $\Delta^0_2\text{-}\mathsf{LEM}$, $H(e_0, e_1, p)$ is also contained in the domain of $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}$, since $\mathrm{Tot}_{e_0} = \mathbb{N}^\mathbb{N} \setminus \mathrm{Tot}_{e_1}$. Therefore, $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO} \circ H(e_0, e_1, p) = \Delta^0_2\text{-}\mathsf{LEM}(e_0, e_1, p)$. □

Theorem 178. *Let $f :\subseteq \mathbb{N}^\mathbb{N} \to \mathbb{N}^\mathbb{N}$ be a single-valued partial function.*

1. *f is $(1, m + 1)$-computable if and only if $f \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^m$.*
2. *f is $(1, \omega|m)$-computable if and only if $f \leq_W \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m$.*
3. *f is $(1, \omega)$-computable if and only if $f \leq_W \Sigma^0_2\text{-}\mathsf{DNE}$ (equivalently, $f \leq_W \bigcup_{m\in\mathbb{N}} \Sigma^0_1\text{-}\mathsf{LEM}^m$, or equivalently $f \leq_W \bigcup_{m\in\mathbb{N}} \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m$).*

Proof. (1) Assume that f is $(1, m)$-computable via a learner Ψ. We need to construct partial computable functions $K :\subseteq \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ and $H :\subseteq \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$ such that $f(x) = K(x, \Sigma^0_1\text{-}\mathsf{LEM}^m \circ H(x))$ for any $x \in \mathrm{dom}(f)$. Fix $x \in \mathbb{N}^\mathbb{N}$, we uniformly construct $H(x) = \langle H_i(x)\rangle_{i<m}$ as follows. We first set $H_i(x; 0) = 1$ for each $i < m$. Assume that $H_i(x; k)$ has been already defined for any $k \leq l$ and each $i < m$. If $\Psi(x \upharpoonright l + 1) = \Psi(x \upharpoonright l)$, then we set $H_i(x; l + 1) = 1$ for each $i < m$. If $\Psi(x \upharpoonright l + 1) \neq \Psi(x \upharpoonright l)$, then let $j(l + 1)$ be the maximal $j < m$ such that $H_j(x; k) = 1$ for any $k \leq l$. Then, we define $H_{j(l+1)}(x; l + 1) = 0$, and, for other $i \neq j(l + 1)$, put $H_i(x; l + 1)$. Clearly, $H : x \mapsto \langle H_i(x)\rangle_{i<m}$ is computable. Assume that $\Sigma^0_1\text{-}\mathsf{LEM}^m \circ H(x) = \sigma^x \in 2^m$. Let $c(x)$ be the least u such that $\sigma^x(u) = 1$. Then $\#\{n \in \mathbb{N} : \Psi(x \upharpoonright n + 1) \neq \Psi(x \upharpoonright n)\} = c(x)$. We define a partial computable function $K : \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ as follows. For given $\langle x, c\rangle \in \mathbb{N}^\mathbb{N} \times m$, we wait for $l(x, c) \in \mathbb{N}$ such that $\#\{n < l(x, c) : \Psi(x \upharpoonright n + 1) \neq \Psi(x \upharpoonright n)\} = c$. Then we set $K(x, c) = \Phi_{\Psi(l(x,c))}(x)$. It is easy to see that $K(x, c(x)) = \Phi_{\lim_n \Psi(x\upharpoonright n)}(x) = f(x)$ for any $x \in \mathrm{dom}(f)$.

Assume that $f \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^m$ via partial computable functions $K :\subseteq \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ and $H :\subseteq \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$. Let us consider a partial computable function $\psi : \mathbb{N}^{<\mathbb{N}} \to \mathbb{N}^{<\mathbb{N}}$ defined by $\psi(\sigma) = K(\sigma, c(\sigma))$, where $c(\sigma; i) = 0$ if $H_i(\sigma; n) = 0$ for some $n < |H_i(\sigma)|$; and $c(\sigma; i) = 1$ otherwise. Here, $H(\sigma) = \langle H_i(\sigma)\rangle_{i<m}$. Note that $c(\sigma; i) \geq c(\tau; i)$ for each $i < m$ whenever $\sigma \subseteq \tau$. Thus, $\#\{n \in \mathbb{N} : c(n + 1) \neq c(n)\} \leq m$. Hence, $\#\{n \in \mathbb{N} : \psi(x \upharpoonright n + 1) \not\supseteq \psi(x \upharpoonright n)\} < m + 1$. Clearly, we have $\psi(x) = \lim_n \psi(x \upharpoonright n) = K(x, \lim_n c(x \upharpoonright n)) = f(x)$ for any $x \in \mathrm{dom}(f)$, since $\lim_n c(x \upharpoonright n)$ converges to $\Sigma^0_1\text{-}\mathsf{LEM}^m \circ H(x)$. By modifying the proof of Proposition 42, we can see that $f = \psi$ is $(1, m + 1)$-computable.

(2) Assume that f is $(1, \omega|m)$-computable via a learner Ψ. We need to construct partial computable functions $K :\subseteq \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ and $H :\subseteq \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$ such that $f(x) = K(x, U \circ H(x))$ for any $x \in \mathrm{dom}(f)$ and any U which realizes $\Sigma^0_2\text{-}\mathsf{LLPO}_m$. We inductively construct $H(x) = \langle H_i(x)\rangle_{i<m}$ as follows. We first declare that 0 *is assigned to* $\Psi(\langle\rangle)$, and put $H_0(\langle\rangle) = \langle 0\rangle$ and $H_i(\langle\rangle) = \langle 1\rangle$ for any $i \neq 0$. Assume that $H_i(\sigma^-)$ has been already defined for each $i < m$. If $\Psi(\sigma) \notin \{\Psi(\tau) : \tau \subsetneq \sigma\}$, then we declare that k *is assigned to* $\Psi(\sigma)$, where k is the least number which has not been assigned to any index $e \in \{\Psi(\tau) : \tau \subsetneq \sigma\}$. Then, put $H_k(\sigma) = H_k(\sigma^-)^\frown\langle 0\rangle$, and $H_i(\sigma) = H_i(\sigma^-)^\frown\langle 1\rangle$ for other $i \neq k$. If $\Psi(\sigma) \notin \{\Psi(\tau) : \tau \subsetneq \sigma\}$, then some k must have been assigned to $\Phi(\sigma)$. For such k, we put $H_k(\sigma) = H_k(\sigma^-)^\frown\langle 0\rangle$, and $H_i(\sigma) = H_i(\sigma^-)^\frown\langle 1\rangle$ for other $i \neq k$. Clearly, $H : x \mapsto \langle \bigcup_{n\in\mathbb{N}} H_i(x \upharpoonright n)\rangle_{i<m}$ is computable. If $x \in \mathrm{dom}(f)$, then there is a unique $i < m$ such that $H_i(x; n) = 0$ for infinitely many $n \in \mathbb{N}$, since $\lim_n \Psi(x \upharpoonright n)$ converges. Let $U(x)$ denote the unique element of $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m(x)$, for any $x \in \mathrm{dom}(\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m)$. Note that $U \circ H(x)$ is assigned to some $\Psi(x \upharpoonright s(x))$ along the construction of $H(x)$, for any $x \in \mathrm{dom}(f)$. Fix the least such $s(x) \in \mathbb{N}$. It is easy to see that $\lim_n \Psi(x \upharpoonright n)$ converges to $\Psi(x \upharpoonright s(x))$. We set $K(x, u) = \Phi_{\Psi(x\upharpoonright t(x,u))}(x)$, where $t(x, u)$ is the least t such that u is assigned to $\Psi(x \upharpoonright t)$ along the construction of $H(x)$. Then, $K : \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ is computable. Moreover, we have $K(x, U \circ H(x)) = \Phi_{\Psi(x\upharpoonright s(x))}(x) = \Phi_{\lim_n \Psi(x\upharpoonright n)}(x) = f(x)$.

Assume that $f \leq_W \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m$ via partial computable functions $K :\subseteq \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ and $H :\subseteq \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$. Let e_i be an index of $\lambda x.K(x,i)$ for each $i < m$. We first compute $h(\sigma, i) = \#\{n < |H_i(\sigma)| : H_i(\sigma; n) = 0\}$, where $H(\sigma) = \langle H_i(\sigma)\rangle_{i<m}$. Then let $c(\sigma)$ be the least $i < m$ such that $h(\sigma, k) \leq h(\sigma, i)$ for any $k < m$. Let us consider a learner $\Psi : \mathbb{N}^{<\mathbb{N}} \to \{e_i\}_{i<m}$ defined by $\Psi(\sigma) = e_{c(\sigma)}$. For any $x \in \mathrm{dom}(f)$, we have $H(x) \in \mathrm{dom}(\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m)$, and then $\lim_n h(x \upharpoonright n, i) = \infty$ for just one $i < m$. Then, $\lim_n c(x \upharpoonright n)$ also converges to such $i < m$. Moreover, for any $x \in \mathrm{dom}(f)$, $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m(H(x)) = \{i\}$ if and only if $\lim_n h(x \upharpoonright n, i) = \infty$. We fix a realizer U of $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m$, i.e., $U(x) \in \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m(x)$ for any $x \in \mathrm{dom}(\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}^m)$. Then, $\lim_n c(x \upharpoonright n) = U \circ H(x)$ for any $x \in \mathrm{dom}(f)$ Therefore, the limit $\lim_n \Psi(x \upharpoonright n)$ converges to $e_{U\circ H(x)}$, and $\#\{\Psi(x \upharpoonright n) : n \in \mathbb{N}\} \leq \#\{e_i : i < m\} \leq m$. Thus, $\Phi_{\lim_n \Psi(x\upharpoonright n)}(x) = \Phi_{e_{U\circ H(x)}}(x) = K(x, U \circ H(x)) = f(x)$ for any $x \in \mathrm{dom}(f)$. Hence, f is $(1, \omega|m)$-computable.

(3) Clearly, $\Sigma^0_2\text{-}\mathsf{DNE}$ is Weihrauch equivalent to the closed choice $\mathsf{C}_\mathbb{N}$ for discrete space $\mathbb{N}$. Therefore, the desired condition follows from Brattka-Brecht-Pauly [11]. The equivalence with $f \leq_W \bigcup_{m\in\mathbb{N}} \Sigma^0_1\text{-}\mathsf{LEM}^m$, and $f \leq_W \bigcup_{m\in\mathbb{N}} \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m$ is easily shown by the same argument as the proofs of (1) and (2). □

Theorem 179. *Let P be any subset of $\mathbb{N}^\mathbb{N}$, and Q be a Δ^0_2 subset of $\mathbb{N}^\mathbb{N}$. Then, $f : Q \to P$ for a $(1, \omega|m)$-computable function iff $f : Q \to P$ for a single-valued function $f \leq_W \Delta^0_2\text{-}\mathsf{LEM}^m$ for some $m \in \mathbb{N}$.*

Proof. By Theorem 156 (2). □

Theorem 180. *Let P be a Π^0_2 subset of $\mathbb{N}^\mathbb{N}$, and Q be any subset of $\mathbb{N}^\mathbb{N}$.*

1. *$P \leq^{<\omega}_1 Q$ iff $f : Q \to P$ for a single-valued function $f \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}_m$ for some $m \in \mathbb{N}$.*
2. *$P \leq^{<\omega}_\omega Q$ iff $f : Q \to P$ for a single-valued function $f \leq_W (\Sigma^0_2)_2\text{-}\mathsf{LLPO}_m$ for some $m \in \mathbb{N}$.*
3. *$P \leq^\omega_1 Q$ iff $f : Q \to P$ for a single-valued function $f \leq_W \Sigma^0_3\text{-}\mathsf{DNE}$ (equivalently, $f \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}_\omega$).*

Proof. (1) Assume that $P \leq^{<\omega}_1 Q$ via a collection $\{\Phi_e\}_{e<b}$ of partial computable functions. It suffices to show that $f : Q \to P$ for some $f \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}_b$. Note that the condition $\Phi_e(x)$ is total and belongs to P is Π^0_2, uniformly in $e < b$ and $x \in \mathbb{N}^\mathbb{N}$. Thus, there is a computable function $H : \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^b$ with $H(x) = \langle H_e(x)\rangle_{e<b}$ satisfying that $H_e(x; n) = 0$ for infinitely many $n \in \mathbb{N}$ if and only if $\Phi_e(x)$ is total and belongs to P. By our assumption, there is $e < b$ such that $H_e(x; n) = 0$ for infinitely many $n \in \mathbb{N}$, for any $x \in Q$. Therefore, $H(x) \in \mathrm{dom}(\Sigma^0_2\text{-}\mathsf{LLPO}_b)$ for any $x \in Q$, and, for any realizer G of $\Sigma^0_2\text{-}\mathsf{LLPO}_b$, $G \circ H(x)$ chooses $e < b$ such that $\Phi_e(x) \in P$. Then, for a computable function $K : \mathbb{N}^\mathbb{N} \times b \to \mathbb{N}^\mathbb{N}$ mapping (x, e) to $\Phi_e(x)$, we have $K(x, G \circ H(x)) = \Phi_e(x) \in P$.

If $f : Q \to P$ for a single-valued function $f \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}_m$, for some $m \in \mathbb{N}$, then there are computable functions $H : \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$ and $K : \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ such that $K(x, G \circ H(x)) \in P$ for any realizer G of $\Sigma^0_2\text{-}\mathsf{LLPO}_m$ and any element $x \in Q$. Then $K(x, i) \in P$ for some $i < m$, since $G \circ H(x) < m$. Set $\Phi_{e(i)}(x) = K(x, i)$ for each $i < m$. Then $P \leq^{<\omega}_1 Q$ via $\{\Phi_{e(i)}\}_{i<m}$.

(2) Assume that $P \leq^{<\omega}_\omega Q$ via a team $\{\Psi_i\}_{e<b}$ of learners. It suffices to show that $f : Q \to P$ for some $f \leq_W (\Sigma^0_2)_2\text{-}\mathsf{LLPO}_b$. The condition $R(i, x)$ stating that $\lim_n \Psi_i(x \upharpoonright n)$ converges and $\Phi_{\lim_n \Psi_i(x\upharpoonright n)}(x)$ is total and belongs to P is $(\Pi^0_2)_2$, uniformly in $i < b$ and $x \in \mathbb{N}^\mathbb{N}$. Thus, there is a computable function $H : \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^b$ with $H(x) = \langle H_i(x)\rangle_{i<b}$ satisfying that $H_i(x; n) \neq 1$ for any $n > m$ if and only if $\Psi(x \upharpoonright n) = \Psi(x \upharpoonright m)$ for any $n > m$, and that $H_i(x; n) = 0$ for infinitely many $n \in \mathbb{N}$ if and only if $\Phi_{\lim_n \Psi_i(x\upharpoonright n)}(x)$ is total and belongs to P, whenever $\lim_n \Psi_i(x \upharpoonright n)$ converges. By our assumption, $R(i, x)$ holds for some $i < b$, for any $x \in Q$. Therefore, $H(x) \in \mathrm{dom}((\Sigma^0_2)_2\text{-}\mathsf{LLPO}_b)$ for any $x \in Q$, and, for any realizer G of $(\Sigma^0_2)_2\text{-}\mathsf{LLPO}_b$, $G \circ H(x)$ chooses $i < b$ and $m \in \mathbb{N}$ such that $\Phi_{\Psi_i(x\upharpoonright m+1)}(x) \in P$. Then, for a computable function $K : \mathbb{N}^\mathbb{N} \times b \times \mathbb{N} \to \mathbb{N}^\mathbb{N}$ mapping (x, i, m) to $\Phi_{\Psi_i(x\upharpoonright m+1)}(x)$, we have $K(x, G \circ H(x)) = \Phi_e(x) \in P$.

If $f : Q \to P$ for a single-valued function $f \leq_W (\Sigma^0_2)_2\text{-}\mathsf{LLPO}_m$, for some $m \in \mathbb{N}$, then there are computable functions $H : \mathbb{N}^{\mathbb{N}} \to (\mathbb{N}^{\mathbb{N}})^m$ and $K : \mathbb{N}^{\mathbb{N}} \times m \times \mathbb{N} \to \mathbb{N}^{\mathbb{N}}$ such that $K(x, G \circ H(x)) \in P$ for any realizer G of $(\Sigma^0_2)_2\text{-}\mathsf{LLPO}_m$ and any element $x \in Q$. Then there is $i < m$ such that $K(x, i, k) \in P$ for any k with $H_i(x; n) \neq 1$ for any $n > k$, where $H(x) = \langle H_i(x) \rangle_{i<m}$. Let $e(i, k)$ be an index of the computable function $\lambda x.K(x, i, k)$, for each $i < m$ and $k \in \mathbb{N}$. The learner $\Delta_i(x \upharpoonright 0)$ first guesses $e(i, 0)$. Assume that $\Delta_i(x \upharpoonright n)$ has been already defined. If $H_i(x; n) = 1$, then we set $\Delta_i(x \upharpoonright n+1) = e(i, n+1)$. If $H_i(x; n) \neq 1$, then we set $\Delta_i(x \upharpoonright n + 1) = \Delta_i(x \upharpoonright n)$. Then, there is a realizer G of $(\Sigma^0_2)_2\text{-}\mathsf{LLPO}_m$ such that, for any $x \in Q$, $\Phi_{\lim_n \Delta_i(x \upharpoonright n)}(x) = K(x, G \circ H(x)) \in P$ for some $i < m$. Hence, $P \leq^{<\omega}_{\omega} Q$ via a team $\{\Delta_i\}_{i<m}$ of learners.

(3) Assume that $P \leq^{\omega}_1 Q$. It suffices to show that $f : Q \to P$ for some $f \leq_W \Sigma^0_3\text{-}\mathsf{DNE}$. Note that the condition $\Phi_e(x)$ is total and belongs to P is Π^0_2, uniformly in $e \in \mathbb{N}$ and $x \in \mathbb{N}^{\mathbb{N}}$. Thus, there is a computable function $H : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ satisfying that $H(x; e, n) = 0$ for infinitely many $n \in \mathbb{N}$ if and only if $\Phi_e(x)$ is total and belongs to P. By our assumption, there is $e \in \mathbb{N}$ such that $H(x; e, n) = 0$ for infinitely many $n \in \mathbb{N}$, for any $x \in Q$. Therefore, $H(x) \in \mathrm{dom}(\Sigma^0_3\text{-}\mathsf{DNE})$ for any $x \in Q$, and, for any realizer G of $\Sigma^0_3\text{-}\mathsf{DNE}$, $G \circ H(x)$ chooses $e < b$ such that $\Phi_e(x) \in P$. Then, for a computable function $K : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \to \mathbb{N}^{\mathbb{N}}$ mapping (x, e) to $\Phi_e(x)$, we have $K(x, G \circ H(x)) = \Phi_e(x) \in P$.

If $f : Q \to P$ for a single-valued function $f \leq_W \Sigma^0_3\text{-}\mathsf{DNE}$, then there are computable functions $H : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ and $K : \mathbb{N}^{\mathbb{N}} \times \mathbb{N} \to \mathbb{N}^{\mathbb{N}}$ such that $K(x, G \circ H(x)) \in P$ for any realizer G of $\Sigma^0_3\text{-}\mathsf{DNE}$ and any element $x \in Q$. Then $K(x, i) \in P$ for some $i \in \mathbb{N}$, since $G \circ H(x) < m$. Set $\Phi_{e(i)}(x) = K(x, i)$ for each $i \in \mathbb{N}$. Then $P \leq^{\omega}_1 Q$ via $\{\Phi_{e(i)}\}_{i \in \mathbb{N}}$. □

Theorem 181. *Let P and Q be any subsets of $\mathbb{N}^{\mathbb{N}}$. Then, $P \blacktriangledown^{\circ} P \leq^1_1 Q$ if and only if $f : Q \to P$ for a single-valued function $f \leq_W \Sigma^0_2\text{-}\mathsf{LEM}$.*

Proof. Assume that there are two computable functions $H : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ and $K : \mathbb{N}^{\mathbb{N}} \times 2 \times \mathbb{N} \to \mathbb{N}^{\mathbb{N}}$ such that $K(x, G \circ H(x)) \in P$ for any $x \in Q$ and any realizer $G : \mathbb{N}^{\mathbb{N}} \to 2 \times \mathbb{N}$ of $\Sigma^0_2\text{-}\mathsf{LEM}$. Then the Σ^0_2 sentence $(\exists v)\theta(v, x)$ is given by $(\exists v)(\forall n > v)H(x; n) \neq 0$. We also define $\Delta(x) = K(x, \langle 0, 0 \rangle)$, and $\Gamma_v(x) = K(x, \langle 1, v \rangle)$, for any $x \in \mathbb{N}^{\mathbb{N}}$. Fix $x \in Q$. If $\theta(v, x)$ is true, then there is a realizer G of $\Sigma^0_2\text{-}\mathsf{LEM}$ mapping $H(x)$ to $(1, v)$. Therefore, $\Gamma_v(x) = K(x, \langle 1, v \rangle) = K(x, G \circ H(x)) \in P$. If $(\forall v)\neg\theta(v, x)$ is true, then there is a realizer G of $\Sigma^0_2\text{-}\mathsf{LEM}$ mapping $H(x)$ to $(0, 0)$. Therefore, $\Delta(x) = K(x, \langle 0, 0 \rangle) = K(x, G \circ H(x)) \in P$. Hence, by Theorem 161, we obtain $P \blacktriangledown^{\circ} P \leq^1_1 Q$.

Conversely, we assume that $P \blacktriangledown^{\circ} P \leq^1_1 Q$. Then, there are computable collection Δ, $\{\Gamma_v\}_{v\in}$ of computable functions, and a Σ^0_2 sentence $\exists v\theta(v, x)$, as in Theorem 161. By analyzing the proof of Theorem 161, we may assume that this Σ^0_2 sentence has an additional property that, if $\theta(v, x)$ is true and $v \leq u$, then $\theta(u, x)$ is also true. For any $x \in \mathbb{N}^{\mathbb{N}}$, put $K(x, \langle 0, n \rangle) = \Delta(x)$ for each $n \in \mathbb{N}$, and $K(x, \langle 1, v \rangle) = \Gamma_v(x)$. From the Σ^0_2 sentence $\exists v\theta(v, x)$, we can easily construct a computable function $H : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ satisfying that $\theta(v, x)$ is true if and only if $H(x; n) \neq 0$ for any $n > v$. Fix $x \in Q$ If $\exists v\theta(v, x)$ is true, then any realizer G of $\Sigma^0_2\text{-}\mathsf{LEM}$ mapping $H(x)$ to some $(1, v)$ witnessing $\theta(v, x)$. Then, $K(x, G \circ H(x)) = \Gamma_v(x) \in P$. If $\forall v\neg\theta(v, x)$ is true, then any realizer G of $\Sigma^0_2\text{-}\mathsf{LEM}$ mapping $H(x)$ to $(0, s)$ for some $s \in \mathbb{N}$. Then, $K(x, G \circ H(x)) = \Delta(x) \in P$. □

Definition 182. Let $\Theta :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ be a partial multi-valued function. A partial *single*-valued function $f :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ is Θ-*computable* if f is Weihrauch reducible to Θ. By $\mathfrak{C}_\Theta$, we denote the least class containing all Θ-computable functions and closed under composition. Then, for subsets P, Q of $\mathbb{N}^{\mathbb{N}}$, we write $P \leq_\Theta Q$ if $f : Q \to P$ for some $f \in \mathfrak{C}_\Theta$. If C_Θ is the class of all computable homeomorphisms between subsets of $\mathbb{N}^{\mathbb{N}}$, then $\leq_\Theta$ is also denoted by $\leq$. If C_Θ is the class of all partial computable functions on $\mathbb{N}^{\mathbb{N}}$, then $\leq_\Theta$ is also denoted by $\leq_{\Sigma^0_1\text{-}\mathsf{DNE}}$, i.e., $\leq_{\Sigma^0_1\text{-}\mathsf{DNE}} = \leq^1_1$.

Corollary 183. *Let X, Y be subsets of $\mathbb{N}^{\mathbb{N}}$, P be a Π^0_2 subset of $\mathbb{N}^{\mathbb{N}}$, and D be a Δ^0_2 subset of $\mathbb{N}^{\mathbb{N}}$.*



FIGURE 5. Constructive principles, and nonuniform computability.

(1) *$Y \leq^1_{<\omega} X$ if and only if $Y \leq_{\Sigma^0_1\text{-LEM}} X$.*
(2) *$Y \leq^1_{\omega|<\omega} X$ if and only if $Y \leq_{\text{Unique}\Sigma^0_2\text{-LLPO}} X$.*
(3) *$Y \leq^1_{\omega|<\omega} D$ if and only if $Y \leq_{\text{Unique}\Sigma^0_2\text{-LLPO}} D$ if and only if $Y \leq_{\Delta^0_2\text{-LEM}} D$.*
(4) *$Y \leq^1_{\omega} X$ if and only if $Y \leq_{\Sigma^0_2\text{-DNE}} X$.*
(5) *$P \leq^{<\omega}_1 X$ if and only if $P \leq_{\Sigma^0_2\text{-LLPO}} X$.*
(6) *$Y\blacktriangledown^\circ Y \leq^1_1 X$ if and only if $Y \leq_{\Sigma^0_2\text{-LEM}} X$.*
(7) *$P \leq^{<\omega}_{\omega} X$ if and only if $P \leq_{(\Sigma^0_2)_2\text{-LLPO}} X$.*
(8) *$P \leq^{\omega}_1 X$ if and only if $P \leq_{\Sigma^0_3\text{-DNE}} X$.*

Proof. By Theorem 178, 179, 180, and 181. □

Theorem 184. *The symbols P, Q, and R range over all special Π^0_1 subset of $2^{\mathbb{N}}$, and X ranges over all subsets of $\mathbb{N}^{\mathbb{N}}$.*

(1) *For every P, there exists $Q \leq P$ such that $P \leq_{\Sigma^0_1\text{-LEM}} Q$, whereas, for every X, if $P \leq_{\Sigma^0_1\text{-DNE}} Q \otimes X$ then $P \leq_{\Sigma^0_1\text{-DNE}} X$.*
(2) *There are P and $Q \leq P$ such that $P \leq_{\Delta^0_2\text{-LEM}} Q$ but $P \not\leq_{\Sigma^0_1\text{-LEM}} Q$.*
(3) *There are P and $Q \leq P$ such that $P \leq_{\Sigma^0_2\text{-LLPO}} Q$ but $P \not\leq_{\Delta^0_2\text{-LEM}} Q$.*
(4) *There is P such that, for every Q, if $P \leq_{\Sigma^0_2\text{-LLPO}} Q$, then $P \leq_{\Sigma^0_1\text{-LEM}} Q$.*
(5) *For every P and R, there exists $Q \leq P$ such that $P \leq_{\Sigma^0_2\text{-DNE}} Q$ but $R \not\leq_{\Sigma^0_2\text{-LLPO}} Q$.*
(6) *For every P, there exists $Q \leq P$ such that $P \leq_{\Sigma^0_2\text{-LEM}} Q$, whereas, for every X, if $P \leq_{\Sigma^0_2\text{-DNE}} Q \otimes X$ then $P \leq_{\Sigma^0_2\text{-DNE}} X$.*
(7) *For every P and R, there exists $Q \leq P$ such that $P \leq_{\Sigma^0_3\text{-DNE}} Q$ but $R \not\leq_{(\Sigma^0_2)_2\text{-LLPO}} Q$.*

Proof. Item **1**. By Corollary 88 and 183 (1). Item **2**. By Corollary 59 (2) and 183 (1,3). Item **3**. By Corollary 61 (1) and 183 (3,5). Item **4**. By Theorem 72, and Corollary 183 (1,5). Item **5**. By Corollary 118 (2) and 183 (4,5). Item **6**. By Corollary 129 and 183 (4,6). Item **7**. By Theorem 149, and Corollary 183 (7,8). □

5.4. Duality between Dynamic Operations and Nonconstructive Principles.

Definition 185 (Brattka-Pauly-le Roux [16])**.** For any multi-valued functions $F, G :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$, the *perfect information composition* $F \star G :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ of F and G is defined as follows.

$$F \star G : \langle e\rangle^\frown x \mapsto \{x \oplus y \oplus z : y \in G(x) \ \&\ z \in F \circ \Phi_e(y)\}.$$

Theorem 186 (Brattka-Pauly-le Roux [16])**.** *Let* $F, G :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ *be any multi-valued functions. Then,* $F \star G = \max_{\leq_W}\{F^* \circ G^* : F^* \leq_W F \ \&\ G^* \leq_W G\}$. □

If multi-valued functions $C, D :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ satisfy the condition

$$D \circ E \leq_W F \iff E \leq_W C \star F$$

for any multi-valued functions $E, F :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$, then we may think of D as the *inverse* of C. One could think of our disjunction operators as inverse operators of various constructive principles.

Definition 187. Fix $x \in \mathbb{N}^{\mathbb{N}}$.

(1) $\triangledown(x) = \{y \in (\mathbb{N} \cup \{\sharp\})^{\mathbb{N}} : \#\{n \in \mathbb{N} : y(n) = \sharp\} \leq 1 \ \&\ \mathtt{tail}(y) = x\}$.
(2) $\triangledown_\omega(x) = \{y \in (2 \times \mathbb{N})^{\mathbb{N}} : (\exists i < 2)\ \mathtt{pr}_i(y) = x \ \&\ \mathtt{mc}(y) < \infty\}$.
(3) $\triangledown_\infty(x) = \{y \in (2 \times \mathbb{N})^{\mathbb{N}} : (\exists i < 2)\ \mathtt{pr}_i(y) = x\}$.
(4) $\widehat{\deg}_T(x) = \{y \in \mathbb{N}^{\mathbb{N}} : x \leq_T y\}$.

A multi-valued function $P :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ is *Popperian* if there is a computable function $r :\subseteq \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ satisfying $\Sigma^0_1\text{-}\mathsf{LEM} \circ r(x, y) = \mathbf{1}_{P(x)}(y)$, for any $x \in \mathrm{dom}(P)$ and $y \in \mathbb{N}^{\mathbb{N}}$, where $\mathbf{1}_{P(x)}$ denotes the characteristic function of $P(x)$. In other words, P is Popperian if and only if the condition $y \in P(x)$ is Π^0_1, uniformly in $x \in \mathrm{dom}(P)$ and $y \in \mathbb{N}^{\mathbb{N}}$. Every Popperian multi-valued function is clearly Weihrauch reducible to the closed choice $\mathsf{C}_{\mathbb{N}^{\mathbb{N}}}$ of Baire space $\mathbb{N}^{\mathbb{N}}$.

Proposition 188. *Let* $E, F : \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ *be any multi-valued functions.*

(1) $\triangledown^{(n)} \circ E \leq_W F$ *if and only if* $E \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^n \star F$.
(2) $\triangledown^{(n)}_\omega \circ E \leq_W F$ *if and only if* $E \leq_W \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_n \star F$.
(3) $\triangledown \circ E \leq_W F$ *if and only if* $E \leq_W \Sigma^0_2\text{-}\mathsf{DNE} \star F$, *where* $\triangledown = \bigcup_{n \in \mathbb{N}} \triangledown^{(n)}$.

Moreover, if E is Popperian, then we also have the following conditions.

4. $\triangledown^{(n)}_\infty \circ E \leq_W F$ *if and only if* $E \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}_n \star F$.
5. $\triangledown^{(n)}_\infty \circ \triangledown \circ E \leq_W F$ *if and only if* $E \leq_W (\Sigma^0_2)_2\text{-}\mathsf{LLPO}_n \star F$.
6. $\widehat{\deg}_T \circ E \leq_W F$ *if and only if* $E \leq_W \Sigma^0_3\text{-}\mathsf{DNE} \star F$.

Proof. (1) Assume that there are partial computable functions $H :\subseteq \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ and $K :\subseteq \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ such that $K(x, f \circ H(x)) \in \triangledown^{(n)} \circ E(x)$ for any $x \in \mathrm{dom}(\triangledown^{(n)} \circ E)$ and any realizer f of F. Then, for any realizer f of F, we have the following condition for any $x \in \mathrm{dom}(E)$.

$$K \circ (\mathrm{id} \times f) \circ \langle \mathrm{id}, H \rangle (x) = K(x, f \circ H(x)) \in \triangledown^{(n)} \circ E(x) = \bigtriangledown^1_n E(x).$$

Note that $H^* = \langle \mathrm{id}, H \rangle : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}}$ is computable, $F^* = K \circ (\mathrm{id} \times F) : \mathbb{N}^{\mathbb{N}} \times \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$ is Weihrauch reducible to F. As in the proof of Theorem 156, we can construct an $(1, n)$-computable function $\gamma : \bigtriangledown^1_n E(x) \to E(x)$, uniformly in $x \in \mathrm{dom}(E)$. Therefore, by Theorem 178, we have a function $\gamma \leq_W \Sigma^0_1\text{-}\mathsf{LCM}^n$ satisfying $\gamma \circ f^* \circ H^*(x) \in E(x)$ for any $x \in \mathrm{dom}(E)$ and any realizer f^* of F^*. Consequently, $E \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^n \star F$.

Conversely, we assume that $E \leq_W S^* \circ F^*$ for some $S^* \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^n$ and $F^* \leq_W F$. Then there are computable functions H^*, K^* such that $K^*(x, H^* \circ f(x)) \in F^*(x)$ for any realizer f of F. From any single valued function $f : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$, we can effectively obtain $f^*(x) = K^*(x, H^* \circ f(x))$. Assume that $S^* \leq_W \Sigma^0_1\text{-}\mathsf{LEM}^n$ via $\tilde{H}$ and $\tilde{K}$, and $E \leq_W S^* \circ F^*$ via H and K. We consider $H_f(x) = \tilde{H} \circ f^* \circ H(x)$ and $K_f(x, i) = K(x, \tilde{K}(f^* \circ H(x), i))$ Then, we have the following condition for any $x \in \mathrm{dom}(E)$.

$$K_f(x, \Sigma^0_1\text{-}\mathsf{LCM}^n \circ H_f(x))) \in E(x).$$

By calculating $H_f(x) = \tilde{H} \circ f^* \circ H(x)$, we can approximate $i(f; x) = \Sigma^0_1\text{-}\mathsf{LEM}^n \circ H_f(x)$ uniformly in f. Therefore, we can construct F^+_f to show $\triangledown^{(n)} \circ E \leq_W F$ by the following way. Set $F^+_f(\langle\rangle) = \langle\rangle$, fix $\sigma \in \mathbb{N}^{<\mathbb{N}}$, and assume that $F^+_f(\sigma^-)$ has been already defined. If $i(f; \sigma) \neq i(f; \sigma^-)$, we put $F^+_f(\sigma) =$

$F^+_f(\sigma^-)^\frown\sharp^\frown K_f(\sigma, i(f;\sigma))$. Otherwise, F^+_f continues the approximation of $K_f(\sigma, i(f;\sigma))$. It is not hard to see that $F^+_f(x) \in \triangledown^{(n)} \circ E(x)$ for any $x \in \mathrm{dom}(E)$ and any realizer f of F. Then, F^+_f is Weihrauch reducible to $\langle K_f, H_f\rangle$, and $\langle K_f, H_f\rangle$ is Weihrauch reducible to f. Moreover, these reduction are not depend on f. Hence, $\triangledown^{(n)} \circ E \leq_W F$.

(2,3) By the same argument as in the proof of the item (1).

(4) Assume that $E :\subseteq \mathbb{N}^\mathbb{N} \rightrightarrows \mathbb{N}^\mathbb{N}$ is Popperian, and there are partial computable functions $H :\subseteq \mathbb{N}^\mathbb{N} \to \mathbb{N}^\mathbb{N}$ and $K :\subseteq \mathbb{N}^\mathbb{N} \times \mathbb{N}^\mathbb{N} \to \mathbb{N}^\mathbb{N}$ such that $K(x, f \circ H(x)) \in \triangledown^{(n)}_\infty \circ E(x)$ for any $x \in \mathrm{dom}(\triangledown^{(n)}_\infty \circ E)$ and any realizer f of F. Then, for any realizer f of F, we have the following condition for any $x \in \mathrm{dom}(E)$.

$$K \circ (\mathrm{id} \times f) \circ \langle \mathrm{id}, H\rangle(x) = K(x, f \circ H(x)) \in \triangledown^{(n)}_\infty \circ E(x) = [\nabla_\infty]^1_n E(x).$$

As in the proof of Theorem 156, we can construct an $(n, 1)$-computable function $\gamma : [\nabla]^1_n E(x) \to E(x)$, uniformly in $x \in \mathrm{dom}(E)$. Here, note that $E(x)$ is a $\Pi^0_1(x)$ subset of Cantor space, uniformly in x. Therefore, by relativizing Theorem 180, we have a function $\gamma \leq_W \Sigma^0_2\text{-}\mathsf{LLPO}^n$ satisfying $\gamma \circ (\mathrm{id} \times f) \circ \langle \mathrm{id}, H\rangle(x) \in E(x)$ for any $x \in \mathrm{dom}(E)$ and any realizer f of F. Consequently, $E \leq_W \Sigma^0_1\text{-}\mathsf{LLPO}^n \star F$.

(5,6) By the same argument as in the proof of the item (4). □

Definition 189. The Σ^0_1 *lessor limited principle of omniscience with* (m/k) *wrong answers*, $\Sigma^0_1\text{-}\mathsf{LLPO}_{m/k}$, is the following multi-valued function.

$$\Sigma^0_1\text{-}\mathsf{LLPO}_{m/k} :\subseteq \mathbb{N}^\mathbb{N} \rightrightarrows k, \qquad x \mapsto \{l < k : (\forall n \in \mathbb{N})\ x(kn + l) = 0\}.$$

Here, $\mathrm{dom}(\Sigma^0_1\text{-}\mathsf{LLPO}_{m/k}) = \{x \in \mathbb{N}^\mathbb{N} : x(n) \neq 0, \text{ for at most } m \text{ many } n \in \mathbb{N}\}$.

Remark. It is well-known that the parallelization of $\Sigma^0_1\text{-}\mathsf{LLPO}_{1/2}$ is equivalent to Weak König's Lemma, WKL (hence, is Weihrauch equivalent to the closed choice for Cantor space, $\mathsf{C}_{2^\mathbb{N}}$).

Definition 190.

(1) (Cenzer-Hinman [21]) A set $P \subseteq k^\mathbb{N}$ is (m, k)*-separating* if $P = \prod_{n\in\mathbb{N}} F_n$ for some uniform sequence $\{F_n\}_{n\in\mathbb{N}}$ of Π^0_1 sets $F_n \subseteq k$, where $\#(k \setminus F_n) \leq m$ for any $n \in \mathbb{N}$.

(2) A function $f : \mathbb{N}^m \to k$ is k*-valued* m*-diagonally noncomputable in* $\alpha \in \mathbb{N}^\mathbb{N}$ if the value $f(\langle e_0, \dots, e_{m-1}\rangle)$ does not belong to $\{\Phi_{e_i}(\alpha; \langle e_0, \dots, e_{m-1}\rangle) : i < m\}$ for each $\langle e_0, \dots, e_{m-1}\rangle \in \mathbb{N}^m$. By $\mathsf{DNR}_{m/k}(\alpha)$, we denote the set of all k-valued functions which are m-diagonally noncomputable in α.

(3) The (m/k) *diagonally noncomputable operation* $\mathsf{DNR}_{m/k} : \mathbb{N}^\mathbb{N} \rightrightarrows k^\mathbb{N}$ is the multi-valued function mapping $\alpha \in \mathbb{N}^\mathbb{N}$ to $\mathsf{DNR}_{m/k}(\alpha)$.

Remark. Clearly $\mathsf{DNR}_{m/k}(\emptyset)$ is (m, k)-separating. The Medvedev degree structure of (m, k)-separating sets have been studied by Cenzer-Hinman [21]. Diagonally noncomputable functions are extensively studied in connection with *effective randomness*, for example, see Greenberg-Miller [34].

Proposition 191. $\mathsf{DNR}_{m/k}$ *is Weihrauch equivalent to* $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}$.

Proof. To see $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k} \leq_W \mathsf{DNR}_{m/k}$, for given $(x_i : i \in \mathbb{N})$, let e^i_t be an $\bigoplus_{i\in\mathbb{N}} x_i$-computable index of an algorithm, for any argument, which returns l at stage s if $l \in L_{s+1} \setminus L_s$ and $\#L_s = t$, where $L_s = \{l^* < k : (\exists n)\ kn + l^* < s\ \&\ x_i(kn + l^*) \neq 0\}$. Clearly, $\{e^i_t : i \in \mathbb{N}\ \&\ t < m\}$ is computable uniformly in $\bigoplus_{i\in\mathbb{N}} x_i$. For any $f \in \mathsf{DNR}_{m/k}(\bigoplus_{i\in\mathbb{N}} x_i)$, the function $i \mapsto f(\langle e^i_0, \dots, e^i_{m-1}\rangle)$ belongs to $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}(\langle x_i : i \in \mathbb{N}\rangle)$. Conversely, for given $x \in \mathbb{N}^\mathbb{N}$, for the i-th m-tuple $\langle e_0, \dots, e_{m-1}\rangle \in \mathbb{N}^m$, we set $x_i(ks + l) = 1$ if $\Phi_{e_t}(\langle e_0, \dots, e_{m-1}\rangle)$ converges to $l < k$ at stage $s \in \mathbb{N}$ for some $t < m$, and otherwise we set $x_i(ks + l) = 0$. Clearly $\{x_i : i \in \mathbb{N}\}$ is uniformly computable in x. Then, for any $\langle l_i : i \in \mathbb{N}\rangle \in \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}(\langle x_i : i \in \mathbb{N}\rangle) \subseteq k^\mathbb{N}$, we have $l_i \notin \{\Phi_{e_t}(\langle e_0, \dots, e_{m-1}\rangle) : t < m\}$ by our construction. Hence, the k-valued function $i \mapsto l_i$ is m-diagonally noncomputable in x. □

Corollary 192. *Let* $k \geq 2$ *be any natural number.*

(1) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k} \not\leq_W \Sigma^0_2\text{-}\mathsf{DNE} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k+1}$.
(2) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k} \not\leq_W \Sigma^0_2\text{-}\mathsf{LLPO} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k+1}$.
(3) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k} \leq_W \Sigma^0_2\text{-}\mathsf{LEM} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{1/k+1}$.

Proof. By Corollary 166, Theorem 178, Theorem 180, and Proposition 191, the item (1) and (2) are satisfied. It is not hard to show the item (3) by analyzing Theorem 167 and Theorem 181. □

Remark. By combining the results from Cenzer-Hinman [21] and our previous results, we can actually show the following.

(1) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{n/l} \not\leq_W \Sigma^0_2\text{-}\mathsf{DNE} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}$, whenever $0 < n < l < \lceil k/m \rceil$.
(2) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{n/l} \not\leq_W \Sigma^0_2\text{-}\mathsf{LLPO} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}$, whenever $0 < n < l < \lceil k/m \rceil$.
(3) $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{n/l} \leq_W \Sigma^0_2\text{-}\mathsf{LEM} \star \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k}$, whenever $0 < n < l$ and $0 < m < k$.

These results suggest, within some constructive setting, that the Σ^0_2 law of excluded middle is sufficient to show the formula $\Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{m/k} \to \Sigma^0_1\text{-}\widehat{\mathsf{LLPO}}_{n/l}$, whereas neither the Σ^0_2 double negation elimination nor the Σ^0_2 lessor limited principle of omniscience is sufficient.

Corollary 193. $\mathsf{DNR}_2 \leq_{\Sigma^0_2\text{-}\mathsf{LEM}} \mathsf{DNR}_3$*;* $\mathsf{DNR}_2 \not\leq_{\Sigma^0_2\text{-}\mathsf{LLPO}} \mathsf{DNR}_3$*;* $\mathsf{DNR}_2 \not\leq_{\Sigma^0_2\text{-}\mathsf{DNE}} \mathsf{DNR}_3$*;* $\mathsf{MLR} \leq_{\Sigma^0_2\text{-}\mathsf{LEM}} \mathsf{DNR}_3$*; and* $\mathsf{MLR} \not\leq_{\Sigma^0_2\text{-}\mathsf{DNE}} \mathsf{DNR}_3$. *Here, recall that* MLR *denotes the set of all Martin-Löf random reals.*

Proof. For the first three statements, see Corollary 166, and Theorem 167, 178, 180. It is easy to see that $\mathsf{MLR} \leq^1_1 \mathsf{DNR}_2 \leq_{\Sigma^0_2\text{-}\mathsf{LEM}} \mathsf{DNR}_3$. It is shown by Downey-Greenberg-Jockusch-Millans [28] that $\mathsf{MLR} \not\leq^1_1 \mathsf{DNR}_3$. By homogeneity of DNR_3 and Proposition 165, we have $\mathsf{MLR} \not\leq_{\Sigma^0_2\text{-}\mathsf{DNE}} \mathsf{DNR}_3$. □

5.5. **Some Intermediate Lattices are Not Brouwerian.** Recall from Medvedev's Theorem 3, Muchnik's Theorem 5, and Proposition 41, the degree structures $\mathcal{D}^1_1$, $\mathcal{D}^1_\omega$, and $\mathcal{D}^\omega_1$ are Browerian. Indeed, we have already observed that one can generate $\mathcal{D}^1_\omega$ from a logical principle so called the Σ^0_2*-double negation elimination*. Though $\mathcal{D}^1_{<\omega}$, $\mathcal{D}^1_{\omega|<\omega}$ and $\mathcal{D}^{<\omega}_1$ are also generated from certain logical principles over $\mathcal{D}^1_1$ as seen before, surprisingly, these degree structures are *not* Brouwerian.

Theorem 194. *The degree structures* $\mathcal{D}^1_{<\omega}$, $\mathcal{D}^1_{\omega|<\omega}$, $\mathcal{D}^{<\omega}_1$, $\mathcal{P}^1_{<\omega}$, $\mathcal{P}^1_{\omega|<\omega}$, *and* $\mathcal{P}^{<\omega}_1$ *are not Brouwerian.*

Put $\mathcal{A}(P,Q) = \{R \subseteq \mathbb{N}^\mathbb{N} : Q \leq^1_{<\omega} P \otimes R\}$, and $\mathcal{B}(P,Q) = \{R \subseteq \mathbb{N}^\mathbb{N} : Q \leq^{<\omega}_1 P \otimes R\}$. Note that $\mathcal{A}(P,Q) \subseteq \mathcal{B}(P,Q)$. Then we show the following lemma.

Lemma 195. *There are* Π^0_1 *sets* $P, Q \subseteq 2^\mathbb{N}$*, and a collection* $\{Z_e\}_{e\in\mathbb{N}}$ *of* Π^0_1 *subsets of* $2^\mathbb{N}$ *such that* $Z_e \in \mathcal{A}(P,Q)$*, and that, for every* $R \in \mathcal{B}(P,Q)$*, we have* $R \not\leq^\omega_1 Z_e$ *for some* $e \in \mathbb{N}$.

Proof. By Jockusch-Soare's theorem 69, we have a collection $\{S_i\}_{i\in\mathbb{N}}$ of nonempty Π^0_1 subsets of $2^\mathbb{N}$ such that $x_k \not\leq_T \bigoplus_{j\neq k} x_j$ for any choice $x_i \in S_i$, $i \in \mathbb{N}$. Consider the following sets.

$$P = \mathsf{CPA}^\frown\{S_{\langle e,0\rangle}{}^\frown S_{\langle e,1\rangle}{}^\frown \dots {}^\frown S_{\langle e,e\rangle}\}_{e\in\mathbb{N}}, \quad Z_e = S_{\langle e,e+1\rangle},$$

$$Q = \mathsf{CPA}^\frown\{Q_n\}_{n\in\mathbb{N}}, \quad \text{where } Q_{\langle e,i\rangle} = \begin{cases} S_{\langle e,i\rangle} \otimes Z_e, & \text{if } i \leq e, \\ (P \setminus [\rho_e]) \otimes Z_e, & \text{if } i = e+1, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Here, ρ_e is the e-th leaf of the corresponding computable tree T_{CPA} for CPA. To see $Z_e \in \mathcal{A}(P,Q)$, choose an element $f \oplus g \in P \otimes Z_e$. If $f \upharpoonright n \in T_{\mathsf{CPA}}$ or $f \upharpoonright n$ extends a leaf except ρ_e, our learner $\Psi((f \upharpoonright n) \oplus g)$ guesses an index of the identity function. If $f \upharpoonright n$ extends ρ_e, then Ψ first guesses $\Phi_{\Psi((f\upharpoonright n)\oplus g)}(f \oplus g) = (f^{\leftarrow|\rho_e|}) \oplus g$. By continuing this guessing procedure, if $f \upharpoonright n$ is of the form $\rho_e{}^\frown\tau^0{}^\frown\tau^1{}^\frown \dots {}^\frown\tau^i{}^\frown\tau$ such that τ^j is a leaf of $S_{\langle e,j\rangle}$ for each $j \leq i$, and τ does not extend a leaf of $S_{\langle e,j+1\rangle}$, then Ψ guesses $\Phi_{\Psi((f\upharpoonright n)\oplus g)}(f \oplus g) =$

$(f^{\leftharpoonup(|\rho_e|+|\tau^0|+\dots+|\tau^i|)}) \oplus g$. Note that $i < e$, since $f \in P$. It is easy to see that $Q \leq^1_{<\omega} P \otimes Z_e$ via the learner Ψ, where $\#\{n \in \mathbb{N} : \Psi((f \oplus g) \upharpoonright n+1) \neq \Psi((f \oplus g) \upharpoonright n)\} \leq e+1$. Therefore, $Z_e \in \mathcal{A}(P, Q)$.

Fix $R \in \mathcal{B}(P, Q)$. As $Q \leq^{<\omega}_1 P \otimes R$, there is $b \in \mathbb{N}$ such that, for every $f \oplus g \in P \otimes R$, we must have $\Phi_e(f \oplus g) \in Q$ for some $e < b$. Suppose for the sake of contradiction that $R \leq^{\omega}_1 Z_{b+1}$. Then, for any $h \in Z_{b+1}$, we have $g \in R$ with $g \leq_T h$. Pick $f_0 \in \rho_{b+1}{}^\frown S_{\langle b+1,0\rangle} \subset P \cap [\rho_{b+1}]$. Since $R \in \mathcal{B}(P, Q)$, there is $e_0 < b$ such that $\Phi_{e_0}(f_0 \oplus g) \in Q$. By our choice of $\{S_n\}_{n \in \mathbb{N}}$ and the property $g \leq_T h \in Z_{b+1} = S_{\langle b+1,b+2\rangle}$, if $e \neq b+1$ or $i \neq 0$, then $Q_{\langle e,i\rangle}$ has no $(f_0 \oplus g)$-computable element. Therefore, $\Phi_{e_0}(f_0 \oplus g)$ have to extend $\rho_{\langle b+1,0\rangle}$. Take an initial segment $\sigma_0 \subset f_0$ determining $\Phi_{e_0}(\sigma_0 \oplus g) \supseteq \rho_{\langle b+1,0\rangle}$. Extend σ_0 to a leaf τ^0 of $S_{b+1,0}$, and choose $f_1 \in \rho^\frown\tau^{0\frown} S_{b+1,1} \subset P$. Again we have $e_1 < b$ such that $\Phi_{e_1}(f_1 \oplus g) \in Q$. As before, $\Phi_{e_1}(f_1 \oplus g)$ have to extend $\rho_{\langle b+1,1\rangle}$. However, $\rho_{\langle b+1,1\rangle}$ is incomparable with $\rho_{\langle b+1,0\rangle}$. Hence, we have $e_1 \neq e_0$. Again take an initial segment $\sigma_1 \subset f_1$ extending σ_0 and determining $\Phi_{e_1}(\sigma_1 \oplus g) \supseteq \rho_{\langle b+1,1\rangle}$. By iterating this procedure, we see that R requires at least $b+1$ many indices e_i. This contradicts our assumption. Therefore, $R \not\leq^{<\omega}_1 Z_{b+1}$. □

Proof of Theorem 194. Let P, Q, and $\{Z_e\}_{e \in \mathbb{N}}$ be Π^0_1 sets in 195. Fix $(\alpha, \beta) \in \{(1, <\omega), (1, \omega| <\omega), (<\omega, 1)\}$. To see $\mathcal{D}^\alpha_\beta$ is not Brouwerian, it suffices to show that there is no (α, β)-least R satisfying $Q \leq^\alpha_\beta P \otimes R$. If R satisfies $Q \leq^\alpha_\beta P \otimes R$, then clearly $R \in \mathcal{B}(P, Q)$ since $\leq^\alpha_\beta$ is stronger than or equals to $\leq^{<\omega}_1$. Then, $R \not\leq^\alpha_\beta Z_e$ for some $e \in \mathbb{N}$. Moreover, $Z_e \in \mathcal{A}(P, Q)$ implies $Q \leq^\alpha_\beta P \otimes Z_e$, since $\leq^\alpha_\beta$ is weaker than or equals to $\leq^1_{<\omega}$. Hence R is not such a smallest set. By the same argument, it is easy to see that $\mathcal{P}^\alpha_\beta$ is not Brouwerian, since Z_e is Π^0_1. □

Theorem 196. *$\mathcal{D}^{<\omega}_\omega$ and $\mathcal{P}^{<\omega}_\omega$ are not Brouwerian. Moreover, the order structures that are induced by $(\mathcal{P}(\mathbb{N}^\mathbb{N}), \leq_{\Sigma^0_2\text{-LEM}})$ and (the set of all nonempty Π^0_1 subsets of $2^\mathbb{N}$, $\leq_{\Sigma^0_2\text{-LEM}}$) are not Brouwerian.*

Lemma 197. *Let $\{S_i\}_{i \leq n}$ be a collection of Π^0_1 subsets of $2^\mathbb{N}$ with the property for each $i \leq n$ that $\bigcup_{k \neq i} S_k$ has no element computable in $x_i \in S_i$. Then, there is no (n, ω)-computable function from $\blacktriangledown_{i \leq n} S_i$ to $\bigoplus_{i \leq n} S_i$.*

Proof. Assume the existence of an (n, ω)-computable function from $\blacktriangledown_{i \leq n} S_i$ to $\bigoplus_{i \leq n} S_i$ which is identified by n many learners $\{\Psi_i\}_{i<n}$. Let F_i be a partial (n, ω)-computable function identified by Ψ, i.e., $F_i(x) = \Phi_{\lim_n \Psi(x \upharpoonright n)}(x)$. Note that $\blacktriangledown_{i \leq n} S_i \subseteq \bigcup_{i<n} \mathrm{dom}(F_i)$. For each $i < n$, put $D_i = \mathrm{dom}(F_i) \cap F_i^{-1}(\bigoplus_{i \leq n} S_i)$. Let T_{S_i} denote the corresponding tree for S_i, for each $i \leq n$. Define $S^\heartsuit_E$ for each $E \subseteq n+1$ to be the set of all infinite paths through the following tree T_E.

$$T_E = \blacktriangledown_{\sigma \in T^E_0} \left(\blacktriangledown_{\sigma \in T^E_1} \left(\dots \left(\blacktriangledown_{\sigma \in T^E_{n-1}} [T^E_n] \right) \dots \right) \right).$$

$$\text{Here, } T^E_i = \begin{cases} T^{ext}_{S_i}, & \text{if } i \in E, \\ \text{some finite subtree of } T^{ext}_{S_i}, & \text{otherwise.} \end{cases}$$

Here, the choice of "some finite subtree of $T^{ext}_{S_i}$" depends on the context, and is implicitly determined when E is defined. For each $E \subseteq n+1$, clearly $S^\heartsuit_E$ is a closed subset of $\blacktriangledown_{i \leq n} S_i$. Divide $S^\heartsuit_{n+1}$ into $n+1$ many parts $\{S^*_i\}_{i \leq n}$, where $S^\heartsuit_{n+1}$ is equal to $\bigcup_{i \leq n} S^*_i$, and each S^*_i is degree-isomorphic to S_i.

For each $i \leq n$, check whether there is a string σ extendible in $S^\heartsuit_{n+1}$ such that $S^\heartsuit_{n+1} \cap D_i \cap [\sigma]$ is contained in S^*_j for some $j \leq n$. If yes, for such a least $i \leq n$, choose a witness $\sigma_0 = \sigma$, and put $A_0 = \{i\}$, and $B_0 = \{j\}$. Then, for such $j \in B_0$, "some finite subtree of $T^{ext}_{S_j}$" is choosen as the set of all strings η used in σ_0 as a part of $T^{ext}_{S_j}$ in the sense of the definition of $\blacktriangledown_{i \leq n} S_i$, or successors of such η in $T^{ext}_{S_j}$. Note that σ_0 is also extendible in $S^\heartsuit_{(n+1) \setminus \{j\}}$. Inductively, for some $s < n$ assume that σ_s, A_s, and B_s has been already defined. For each $i \notin A_s$, check whether there is a string $\sigma \supseteq \sigma_s$ extendible in $S^\heartsuit_{(n+1) \setminus B_s}$ such that $S^\heartsuit_{(n+1) \setminus B_s} \cap D_i \cap [\sigma]$ is contained in S^*_j for some $j \notin B_s$. If yes, for such a least $i \notin A_s$, choose a witness $\sigma_0 = \sigma$, and put $A_{s+1} = A_s \cup \{i\}$, and $B_{s+1} = B_s \cup \{j\}$. As before, for such $j \in B_{s+1}$, "some finite subtree of $T^{ext}_{S_j}$" is choosen as the set of all strings η used in σ_{s+1} as a part of $T^{ext}_{S_j}$, or successors of such η in $T^{ext}_{S_j}$. Note that σ_{s+1} is also extendible in $S^{\heartsuit}_{(n+1)\setminus B_{s+1}}$. If no such $i \notin A_s$ exists, finish our construction of σ, A, and B. Then, put $A = A_s$, $B = B_s$, and define σ^* to be the last witness σ_s.

Put $A^- = n \setminus A$ and $B^- = (n+1) \setminus B$. Note that $\#A^- + 1 = \#B^-$, since $\#A = \#B$. Therefore, B contains at least one element. By our assumption, for any $x \in S^{\heartsuit}_{B^-} \cap [\sigma^*] \neq \emptyset$, we must have $F_i(x) \in \bigoplus_{i \leq n} S_i$ for some $i \in A^-$. Thus, A^- is nonempty.

Fix a sequence $\alpha \in (A^-)^{\mathbb{N}}$ such that, for each $i \in A^-$, there are infinitely many $n \in \mathbb{N}$ such that $\alpha(n) = i$. First set $\tau_0 = \sigma^*$. Inductively assume that $\tau_s \supseteq \sigma^*$ has been already defined. By our definition of σ^*, A and B, if ξ extends σ^*, then the set $S^{\heartsuit}_{B^-} \cap D_{\alpha(s)} \cap [\xi]$ intersects with S^*_j for at least two $j \in B^-$. Therefore, we can choose $x \in S^{\heartsuit}_{B^-} \cap D_{\alpha(s)} \cap [\tau_s] \cap S^*_j \neq \emptyset$ for some $j \in B^-$. Then, $F_{\alpha(s)}(x;0) = j$, by our assumption of $\{S_i\}_{i \leq n}$. Find a string τ^*_s such that $\tau_s \subseteq \tau^*_s \subset x$ and $F_{\alpha(s)}(\tau^*_s;0) = j$. Again, we can choose $x^* \in S^{\heartsuit}_{B^-} \cap D_{\alpha(s)} \cap [\tau^*_s] \cap S^*_k \neq \emptyset$ for some $k \in B^- \setminus \{j\}$. Then, we must have $F_{\alpha(s)}(x^*;0) = k \neq j$. Let τ_{s+1} be a string such that $\tau^*_s \subseteq \tau_{s+1} \subset x^*$ and $F_{\alpha(s)}(\tau_{s+1};0) = k$. Therefore, between τ_s and τ_{s+1}, the learner $\Psi_{\alpha(s)}$ changes his mind.

Define $y = \bigcup_s \tau_s$. Then y is contained in $S^{\heartsuit}_{B^-}$, since $S^{\heartsuit}_{B^-}$ is closed. However, for each $i \in A^-$, by our construction of y, the value $F_i(y)$ does not converge. Moreover, for each $i \notin A^-$, by our definition of A, B, and $\sigma^* \subset y$, even if $F_i(y)$ converges, $F_i(y) \notin \bigoplus_{i \leq n} S_i$. Consequently, there is no (n, ω)-computable function from $\blacktriangledown_{i \leq n} S_i \supset S^{\heartsuit}_{B^-}$ to $\bigoplus_{i \leq n} S_i$ as desired. □

Put $\mathcal{J}(P,Q) = \{R \subseteq \mathbb{N}^{\mathbb{N}} : Q \leq_{\Sigma^0_2\text{-}\mathsf{LEM}} P \otimes R\}$, and $\mathcal{K}(P,Q) = \{R \subseteq \mathbb{N}^{\mathbb{N}} : Q \leq^{<\omega}_{\omega} P \otimes R\}$. Note that $\mathcal{J}(P,Q) \subseteq \mathcal{K}(P,Q)$. Then we show the following lemma.

Lemma 198. *There are Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$, and a collection $\{Z_e\}_{e \in \mathbb{N}}$ of Π^0_1 subsets of $2^{\mathbb{N}}$ such that $Z_e \in \mathcal{J}(P,Q)$, and that, for every $R \in \mathcal{K}(P,Q)$, we have $R \not\leq^{\omega}_1 Z_e$ for some $e \in \mathbb{N}$.*

Proof. By Jockusch-Soare's theorem 69, we have a collection $\{S_i\}_{i \in \mathbb{N}}$ of nonempty Π^0_1 subsets of $2^{\mathbb{N}}$ such that $x_k \not\leq_T \bigoplus_{j \neq k} x_j$ for any choice $x_i \in S_i$, $i \in \mathbb{N}$. Consider the following sets.

$$P = \mathsf{CPA}^\frown\{S_{\langle e,0\rangle} \blacktriangledown S_{\langle e,1\rangle} \blacktriangledown \dots \blacktriangledown S_{\langle e,e\rangle}\}_{e \in \mathbb{N}}, \quad Z_e = S_{\langle e,e+1\rangle},$$

$$Q = \mathsf{CPA}^\frown\{Q_n\}_{n \in \mathbb{N}}, \quad \text{where} \quad Q_{\langle e,i\rangle} = \begin{cases} S_{\langle e,i\rangle} \otimes Z_e, & \text{if } i \leq e, \\ (P \setminus [\rho_e]) \otimes Z_e, & \text{if } i = e+1, \\ \emptyset, & \text{otherwise.} \end{cases}$$

Here, ρ_e is the e-th leaf of the corresponding computable tree T_{CPA} for CPA. To see $Z_e \in \mathcal{J}(P,Q)$, for $f \oplus g \in P \otimes Z_e$, by using Σ^0_1-LEM, check whether f does no extend ρ_e. If no, outputs $\rho_{e,e+1}{}^\frown(f \oplus g)$. If f extends ρ_e, it is not hard to see that an finite iteration of Σ^0_2-LEM can divide $(\rho_e{}^\frown S_{\langle e,0\rangle} \blacktriangledown S_{\langle e,1\rangle} \blacktriangledown \dots \blacktriangledown S_{\langle e,e\rangle}) \otimes Z_e$ into $\{S_{\langle e,i\rangle} \otimes Z_e\}_{i \leq e}$.

Fix $R \in \mathcal{K}(P,Q)$. As $Q \leq^{\omega}_{<\omega} P \otimes R$, some (b, ω)-computable function F maps $P \otimes R$ into Q. Suppose for the sake of contradiction that $R \leq^{\omega}_1 Z_b$. Then, for any $h \in Z_b$, we have $g \in R$ with $g \leq_T h$. Then, F maps $(P \cap [\rho_b]) \otimes \{g\}$ into $Q \cap (\bigcup_{i \leq b} \rho_{\langle b,i\rangle})$ by our choice of $\{S_n\}_{n \in \mathbb{N}}$. Note that $(P \cap [\rho_b]) \otimes \{g\} \equiv^1_1 (\blacktriangledown_{i \leq b} S_{\langle e,i\rangle}) \otimes \{g\}$, and $Q \cap (\bigcup_{i \leq b} \rho_{\langle b,i\rangle}) \equiv^1_1 (\bigoplus_{i \leq b} S_{\langle e,i\rangle}) \otimes Z_e$. Therefore, by Lemma 197, F is not (b, ω)-computable. □

Proof of Theorem 196. Let P, Q, and $\{Z_e\}_{e \in \mathbb{N}}$ be Π^0_1 sets in 198. Then, by the same argument as in the proof of Theorem 194, it is not hard to show the desired statement. □

Corollary 199. *If $(\alpha, \beta) \in \{(1,1), (1,\omega), (\omega,1)\}$, and $(\gamma, \delta) \in \{(1, <\omega), (1, \omega| <\omega), (<\omega, 1), (<\omega, \omega)\}$, then, there is an elementary difference between $\mathcal{D}^{\alpha}_{\beta}$ and $\mathcal{D}^{\gamma}_{\delta}$, in the language of partial orderings $\{\leq\}$.*

Proof. By Medvedev's Theorem 3, Muchnik's Theorem 5, and Proposition 41, the degree structures $\mathcal{D}^1_1$, $\mathcal{D}^1_{\omega}$, and $\mathcal{D}^{\omega}_1$ are Browerian, i.e., they satisfy the following elementary formula ψ in the language of partial orders.

$$\psi \equiv (\forall p, q)(\exists r)(\forall s)\ (p \leq q \vee r \ \&\ (p \leq q \vee s \ \rightarrow\ r \leq s)).$$

Here, the supremum $\vee$ is first-order definable in the language of partial orders. On the other hand, by Theorem 194 and 196, $\mathcal{D}^1_{<\omega}$, $\mathcal{D}_{\omega|<\omega}$, $\mathcal{D}^{<\omega}_1$, and $\mathcal{D}^{<\omega}_{\omega}$ are not Brouwerian, i.e., they satisfy $\neg\psi$. □

5.6. **Borel Measurability, and Backtrack Games.** Berardi-Coquand-Hayashi [5] showed that a 1-*backtrack Tarski game* provides a semantics of positive arithmetical fragment of Limit Computable Mathematics (i.e., Δ^0_2-mathematics, in the sense of Kleene realizability).

Theorem 200 (Berardi-Coquand-Hayashi [5])**.** *A positive arithmetical formula A is true in the Limit Realizability Interpretation if and only if the $\exists$-player has a computable winning strategy in the* 1-*backtracking game* $\mathsf{bck}(\mathcal{G}(A))$ *associated with the Tarski game for A (for notations, see* [5]*).*

Meanwhile, in the study of Wadge degree, Van Wesep [73] introduced *backtrack game*, and Andretta [3] used this game to characterize the $\mathbf{\Delta}^0_2$ functions (also called the first level Borel functions) on Baire space $\mathbb{N}^{\mathbb{N}}$.

Definition 201. Fix any countable ordinal $\xi < \omega_1$, and a function f on Baire space $\mathbb{N}^{\mathbb{N}}$.

1. f is $\mathbf{\Delta}^0_\xi$ if the preimage $f^{-1}(A)$ is $\mathbf{\Sigma}^0_\xi$ for every $\mathbf{\Sigma}^0_\xi$ set $A \subseteq \mathbb{N}^{\mathbb{N}}$.
2. f is $\mathbf{\Sigma}^0_\xi$-*measurable* if the preimage $f^{-1}(A)$ is $\mathbf{\Sigma}^0_\xi$ for every $\mathbf{\Sigma}^0_1$ set $A \subseteq \mathbb{N}^{\mathbb{N}}$.

Clearly, every $\mathbf{\Delta}^0_\xi$ function on Baire space $\mathbb{N}^{\mathbb{N}}$ is $\mathbf{\Sigma}^0_\xi$-measurable. The effective hierarchy of Borel measurable functions is studied by Brattka [10]. However, in general, an effective Σ^0_ξ measurable function is not nonuniformly computable. For example, an effectively Σ^0_{n+1}-measurable function can map a computable point to a point of Turing degree $\mathbf{0}^{(n)}$. Indeed, the class of (effectively) Σ^0_ξ-measurable functions does not closed under composition, in general, whereas the class of $\mathbf{\Delta}^0_\xi$ functions must be closed under composition. Our piecewise computability theorem 156 suggests that our notions of learnability is not like the effective Σ^0_ξ-measurability but more like effective versions of $\mathbf{\Delta}^0_\xi$ functions, because of the following observations.

Definition 202. Let f be a partial function on Baire space $\mathbb{N}^{\mathbb{N}}$, and $\mathbf{\Gamma}$ (resp. Γ) be a boldface (resp. lightface) pointclass.

1. f is $\mathbf{\Gamma}$-*piecewise continuous* if there is a $\mathbf{\Gamma}$ partition $\{Q_i\}_{i\in\mathbb{N}}$ of $\mathrm{dom}(f)$ such that $f \upharpoonright Q_i$ is continuous for each $i \in \mathbb{N}$.
2. f is Γ-*piecewise computable* if there is a Γ partition $\{Q_i\}_{i\in\mathbb{N}}$ of $\mathrm{dom}(f)$ such that $f \upharpoonright Q_i$ is computable uniformly in $i \in \mathbb{N}$.

Theorem 203. *Let $f : \mathbb{N}^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ be a function.*

1. *(Jayne-Rogers* [42]*) f is $\mathbf{\Delta}^0_2$ if and only if f is $\mathbf{\Pi}^0_1$-piecewise continuous.*
2. *(Semmes* [59]*) f is $\mathbf{\Delta}^0_3$ if and only if f is $\mathbf{\Pi}^0_2$-piecewise continuous.*

Proposition 204. *Let P and Q be any subsets of Baire space $\mathbb{N}^{\mathbb{N}}$.*

1. *A function $f : Q \to P$ is $(1, \omega)$-computable if and only if f is Π^0_1-piecewise computable.*
2. *Whenever P is Π^0_1, an $(\omega, 1)$-computable function $f : Q \to P$ exists if and only if a Π^0_2-piecewise computable function $f^* : Q \to P$ exists.*

Proof. By analyzing the proof of Theorem 156. □

Actually, our dynamic models directly fit into the backtrack and multitape game characterization of subclasses of Borel measurable functions. We now introduce various games based on *the Wadge game*, *the backtrack game*, and *the multitape game*,

Definition 205. Fix a partial function f on $\mathbb{N}^{\mathbb{N}}$, and a set X which has no intersection with $\mathbb{N}$. The set X may contain $\mathtt{pass}$, $\mathtt{back}$, $(\mathtt{move}, i)$ for each $i \in \mathbb{N}$. Then, we introduce various two-players games on f as follows. At every round $n \in \mathbb{N}$, Player I chooses an element $x_n \in \mathbb{N}$, and Player II chooses an element $y_n \in \mathbb{N} \cup X$.

$$\begin{array}{lccccccc} \text{I:} & x_0 & & x_1 & & x_2 & & \dots \\ \text{II:} & & y_0 & & y_1 & & y_2 & \dots \end{array}$$

A pair of infinite sequences $\langle x, y\rangle \in \mathbb{N}^{\mathbb{N}} \times (\mathbb{N} \cup X)^{\mathbb{N}}$ is called *a play*. Fix a play $\langle x, y\rangle$, where $x = \langle x_n\rangle_{n\in\mathbb{N}}$ and $y = \langle y_n\rangle_{n\in\mathbb{N}}$. We say that Player I *violates the basic rule* if $x \notin \mathrm{dom}(f)$, and that Player II *violates the basic rule* if either $y_n \in \{\mathtt{pass}, (\mathtt{move}, i) : i \in \mathbb{N}\}$ for almost all $n \in \mathbb{N}$, or $y_n = \mathtt{back}$ for infinitely many $n \in \mathbb{N}$. We also say that Player II *violates the rule m* if y contains at least m many $\mathtt{back}$'s, and *violate the rule $*$* if there is an infinite increasing sequence $\langle m(n)\rangle_{n\in\mathbb{N}}$ of natural numbers with $y_{m(n)} = (\mathtt{move}, i_n)$ and $i_{n+1} \neq i_n$ for any $n \in \mathbb{N}$. For each $i \in \mathbb{N}$, *the i-th content* of the play y of Player II is a function $\mathtt{content}_i : (\mathbb{N} \cup X)^{\mathbb{N}} \to \mathbb{N}^{\mathbb{N}}$ which is inductively defined as follows. Set $\mathtt{content}_i(\langle\rangle) = \langle\rangle$ and $\mathtt{tape}(\langle\rangle) = 0$. Assume that $\mathtt{content}_i(y \upharpoonright n)$ and $\mathtt{tape}(y \upharpoonright n)$ have been already defined for each $i \in \mathbb{N}$.

$$\mathtt{content}_i(y \upharpoonright n+1) = \begin{cases} \mathtt{content}_i(y \upharpoonright n)^\frown\langle y_n\rangle & \text{if } y_n \in \mathbb{N} \ \&\ i = \mathtt{tape}(y \upharpoonright n), \\ \langle\rangle & \text{if } y_n = \mathtt{back} \ \&\ i = \mathtt{tape}(y \upharpoonright n), \\ \mathtt{content}_i(y \upharpoonright n) & \text{otherwise.} \end{cases}$$

$$\mathtt{tape}(y \upharpoonright n+1) = \begin{cases} i & \text{if } y_n = (\mathtt{move}, i), \\ \mathtt{tape}(y \upharpoonright n) & \text{otherwise.} \end{cases}$$

Then, for each $i \in \mathbb{N}$, we define $\mathtt{content}_i(y) = \lim_{n\in\mathbb{N}} \mathtt{content}_i(y \upharpoonright n)$ for any $y \in (\mathbb{N} \cup X)^{\mathbb{N}}$. We say that Player II *wins* (resp. *is winnable*) on the play $\langle x, y\rangle \in \mathbb{N}^{\mathbb{N}} \times (\mathbb{N} \cup X)^{\mathbb{N}}$ of the game $G(f, X)$ if either Player II does not violate the basic rule, and $f(x) = \mathtt{content}_i(y)$ for the least $i \in \mathbb{N}$ with $\mathtt{content}_i(y)$ being total (resp. for some $i \in \mathbb{N}$), or Player I violates the basic rule. We also say that Player II *wins* (resp. *is winnable*) on the play $\langle x, y\rangle$ of the game $G_m(f, X)$ if Player II wins (resp. is winnable) the game $G(f, X)$ and does not violate the rule m, and that Player II *wins* (resp. *is winnable*) the game $G_*(f, X)$ if Player II wins (resp. is winnable) the game $G(f, X)$ and does not violate the rule $*$.

A *strategy* of Player II is a function $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup X)^{<\mathbb{N}}$ such that $|\psi(\sigma)| = |\sigma|$ for each $\sigma \in \omega^{<\omega}$, and $\psi(\sigma) \subseteq \psi(\tau)$ whenever $\sigma \subseteq \tau$. A strategy ψ of Player II is *winning* (resp. *winnable*) in the game G if Player II wins (resp. is winnable) the game G on the play $\langle x, \bigcup_{n\in\mathbb{N}} \psi(x \upharpoonright n)\rangle$ for any $x \in \mathbb{N}^{\mathbb{N}}$.

We write P, B, and M_α for $\{\mathtt{pass}\}$, $\{\mathtt{back}\}$, and $\{(\mathtt{move}, i) : i < \alpha\}$, respectively, for each $\alpha \leq \omega$. Then, for $\mathsf{S}, \mathsf{T}, \mathsf{U} \in \{\mathsf{P}, \mathsf{B}, \mathsf{M}_\alpha\}_{\alpha\leq\omega}$, the union $\mathsf{S} \cup \mathsf{T} \cup \mathsf{U}$ is denoted by STU.

Remark. The games $G(f, \mathsf{P})$, $G(f, \mathsf{PB})$, and $G(f, \mathsf{PM}_\omega)$ are essentially same as *the Wadge game, the backtrack game*, and *the multitape game*, respectively. For original definition of variations of Wadge games, see also Motto Ros [51] and Semmes [59].

Theorem 206. *Let f be a partial function on Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) (Wadge [74]) *f is continuous if and only if Player II has a winning strategy in the game $G(f, \mathsf{P})$.*
(2) (Andretta [3]) *f is $\mathbf{\Delta}^0_2$ if and only if Player II has a winning strategy in the game $G(f, \mathsf{PB})$.*
(3) (Andretta, Semmes [58]) *f is $\mathbf{\Pi}^0_2$-piecewise continuous if and only if Player II has a winning strategy in the game $G(f, \mathsf{PM}_\omega)$.*

Theorem 207 (Game representation)**.** *Let f be a partial function on Baire space $\mathbb{N}^{\mathbb{N}}$.*

(1) *f is $(1, 1)$-computable if and only if Player II has a computable winning strategy in the game $G(f, \mathsf{P})$.*
(2) *f is $(1, m)$-computable if and only if Player II has a computable winning strategy in the game $G_m(f, \mathsf{PB})$.*

(3) *f is $(1, \omega|m)$-computable if and only if Player II has a computable winning strategy in the game $G_*(f, \mathsf{PM}_m)$.*
(4) *f is $(1, \omega)$-computable if and only if Player II has a computable winning strategy in the game $G(f, \mathsf{PB})$.*
(5) *f is $(m, 1)$-computable if and only if Player II has a computable winnable strategy in the game $G(f, \mathsf{PM}_m)$.*
(6) *f is (m, ω)-computable if and only if Player II has a computable winnable strategy in the game $G(f, \mathsf{PBM}_m)$.*
(7) *f is $(\omega, 1)$-computable if and only if Player II has a computable winnable strategy in the game $G(f, \mathsf{PM}_\omega)$.*

Proof. (1,3) We need to construct a winning strategy $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{pass}, \mathtt{back}\})^{<\mathbb{N}}$ from a given partial $(1, \omega)$-computable function $f :\subseteq \mathbb{N}^\mathbb{N} \to \mathbb{N}^\mathbb{N}$. Assume that f is $(1, \omega)$-computable via a learner Ψ. We inductively define a strategy $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{pass}, \mathtt{back}\})^{<\mathbb{N}}$ and an auxiliary parameter $\eta : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{back}\})^{<\mathbb{N}}$. Set $\psi(\langle\rangle) = \eta(\langle\rangle) = \langle\rangle$, and assume that $\psi(\sigma^-)$ and $\eta(\sigma^-)$ have been already defined. If $\Psi(\sigma) = \Psi(\sigma^-)$, then set $\eta_*(\sigma) = \eta(\sigma^-)^\frown\tau$, where $\Phi_{\Psi(\sigma)}(\sigma) = \Phi_{\Psi(\sigma)}(\sigma^-)^\frown\tau$. If $\Psi(\sigma) \neq \Psi(\sigma^-)$, then set $\eta_*(\sigma) = \eta(\sigma^-)^\frown\mathtt{back}^\frown\Phi_{\Psi(\sigma)}(\sigma)$. From the parameter $\eta_*(\sigma)$, we define $\psi(\sigma)$ as follows. If $\eta_*(\sigma) = \langle\rangle$, then $\psi(\sigma) = \psi(\sigma^-)^\frown\mathtt{pass}$, and $\eta(\sigma) = \eta_*(\sigma)$. If $\eta_*(\sigma) \neq \langle\rangle$, then $\psi(\sigma) = \psi(\sigma^-)^\frown\eta_*(\sigma)(0)$, and $\eta(\sigma) = \mathtt{sh}^1(\eta_*(\sigma))$. Note that $\{n \in \mathbb{N} : (\bigcup_k \psi(x \upharpoonright k))(n) = \mathtt{back}\} = \{n \in \mathbb{N} : \Psi(x \upharpoonright n+1) \neq \Psi(x \upharpoonright n)\}$ for any $x \in \mathrm{dom}(f)$. It is easy to see that ψ is a computable winning strategy in the game $G(f, \mathsf{PB})$.

Assume that a computable winning strategy ψ^* in the game $G(f, \mathsf{PB})$ is given. We consider the computable function $\psi(\sigma) = \mathtt{content}_0(\psi^*(\sigma))$. Then $\{n \in \mathbb{N} : \psi(x \upharpoonright n+1) \not\supseteq \psi(x \upharpoonright n)\}$ is finite, for any $x \in \mathrm{dom}(f)$, since $\bigcup_{n\in\mathbb{N}} \psi(x \upharpoonright n)$ contains finitely many $\mathtt{back}$'s. Moreover, $f(x) = \lim_n \psi(x \upharpoonright n)$. Thus, by Proposition 42, f is $(1, \omega)$-computable.

(2) By Theorem 178 (2), if f is $(1, \omega|m)$-computable, then $f \leq_W \mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m$. Assume that we have computable functions $H : \mathbb{N}^\mathbb{N} \to (\mathbb{N}^\mathbb{N})^m$ and $K : \mathbb{N}^\mathbb{N} \times m \to \mathbb{N}^\mathbb{N}$ satisfying $f(x) = K(x, G \circ H(x))$ for any $x \in \mathrm{dom}(f)$, for the unique realizer G of $\mathsf{Unique}\Sigma^0_2\text{-}\mathsf{LLPO}_m$. For each $\sigma \in \mathbb{N}^{<\mathbb{N}}$, the i-th coordinate of $H(\sigma)$ is denoted by $H_i(\sigma)$, i.e., $H(\sigma) = \langle H_i(\sigma)\rangle_{i<m}$. We inductively define a strategy $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{pass}, (\mathtt{move}, i) : i < m\})^{<\mathbb{N}}$ and an auxiliary parameter $\eta : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{(\mathtt{move}, i) : i < m\})^{<\mathbb{N}}$. Set $\psi(\langle\rangle) = \eta(\langle\rangle) = \langle\rangle$, and assume that $\psi(\sigma^-)$ and $\eta(\sigma^-)$ have been already defined. We pick the least $i < m$ such that γ_i contains 0, where $H_i(\sigma) = H_i(\sigma^-)^\frown\gamma_i$. Then, we set $\eta_*(\sigma) = \eta(\sigma^-)^\frown(\mathtt{move}, i)^\frown\tau$, where $K(x, i) = \mathtt{content}_i(\psi(\sigma^-))^\frown\tau$. If there is no such $i < m$, then we set $\eta_*(\sigma) = \eta(\sigma^-)$. From the parameter $\eta_*(\sigma)$, we define $\psi(\sigma)$ as follows. If $\eta_*(\sigma) = \langle\rangle$, then $\psi(\sigma) = \psi(\sigma^-)^\frown\mathtt{pass}$, and $\eta(\sigma) = \eta_*(\sigma)$. If $\eta_*(\sigma) \neq \langle\rangle$, then $\psi(\sigma) = \psi(\sigma^-)^\frown\eta_*(\sigma)(0)$, and $\eta(\sigma) = \mathtt{sh}^1(\eta_*(\sigma))$. For any $x \in \mathrm{dom}(f)$, if $G \circ H(x) = i$, then $H_k(x)$ contains only finitely many 0's for any $k \neq i$. Therefore, $\psi(x) = \bigcup_n \psi(x \upharpoonright n)$ contains infinitely many $(\mathtt{move}, i)$'s, and only finitely many $(\mathtt{move}, j)$'s for any $k \neq i$. Therefore, ψ does not violate the rule $*$, and we observe that $\mathtt{content}_i \circ \psi(x) = K(x, i) = K(x, G \circ H(x)) = f(x)$. Hence, ψ is a winning strategy in the game $G_*(f, \mathsf{PM}_m)$.

Assume that a computable winning strategy ψ^* in the game $G_*(f, \mathsf{PM}_m)$ is given. Let $e(i)$ be an index of a partial computable function $x \mapsto \mathtt{content}_i \circ \psi^*(x)$ for each $i < m$. Since ψ^* does not violate the rule $*$, there is a unique $i < m$ such that $\Phi_{e(i)} = \mathtt{content}_i \circ \psi^*(x)$ is total, for any $x \in \mathrm{dom}(f)$. We inductively define a learner Ψ. The learner Ψ first guesses $\Psi(\langle\rangle) = e(0)$. Set $\Psi(\sigma) = \Psi(\sigma^-)$ when there is no $i < m$ such that $|\Phi_{e(i)}(\sigma)| > |\Phi_{e(i)}(\sigma^-)|$. Otherwise, for the least such $i < m$, the learner guesses $\Psi(\sigma) = e(i)$. Clearly, $\#\{\Psi(x \upharpoonright n) : n \in \mathbb{N}\} < m$ for any $x \in \mathbb{N}^\mathbb{N}$. It is easy to check that, for any $x \in \mathrm{dom}(f)$, $\lim_n \Psi(x \upharpoonright n)$ converges to $e(i)$ for the unique $i < m$ ensuring the totality of $\mathtt{content}_i \circ \psi^*(x)$, and, for such $i < m$, we have $\Phi_{\lim_n \Psi(x \upharpoonright n)}(x) = \mathtt{content}_i \circ \psi^*(x) = f(x)$. Consequently, f is $(1, \omega|m)$-computable.

(4,6) For a given collection $\{\Phi_i\}_{i\in I}$ of partial computable functions, we can easily construct a strategy $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{pass}, (\mathtt{move}, i) : i \in I\})$ ensuring $\mathtt{content}_i \circ \psi(x) = \Phi_i(x)$ for any $x \in \mathbb{N}^\mathbb{N}$. Therefore, f is nonuniformly computable via $\{\Phi_i\}_{i\in I}$, then ψ is winnable in $G(f, \mathsf{PM}_I)$. Conversely, if a winnable strategy $\psi : \mathbb{N}^{<\mathbb{N}} \to (\mathbb{N} \cup \{\mathtt{pass}, (\mathtt{move}, i) : i \in I\})$ of the game $G(f, \mathsf{PM}_I)$ is given. Then we consider the partial computable function Γ_i computing $\Gamma_i(x) = \mathtt{content}_i \circ \psi(x)$ for any $x \in \mathbb{N}^{\mathbb{N}}$. It is easy to see that f is nonuniformly computable via $\{\Gamma_i\}_{i \in I}$.

(5) By combining the proofs of the items (3) and (4), it is not hard to see the equivalence of the (m, ω)-computability of f and the computable winnability in the game $G(f, \mathsf{PBM}_m)$. □

Remark. We may introduce more general multitape games based on our dynamic tape models, and nested (nested nested, nested nested nested, etc.) tape models.

5.7. **Open Questions.**

Question 208 (Small Questions)**.**

(1) *Determine the intermediate logic corresponding the degree structure $\mathcal{D}^1_\omega$, where recall that $\mathcal{D}^1_1$ and $\mathcal{D}^\omega_1$ are exactly Jankov's Logic.*
(2) *For every Π^0_1 set $Q \leq^1_1 \mathsf{CPA}^\frown\mathsf{CPA}$, does there exist nonempty Π^0_1 sets $P_0, P_1 \subseteq 2^{\mathbb{N}}$ such that $Q \equiv^1_1 P_0{}^\frown P_1$?*
(3) *Does there exist a special Π^0_1 set $P \subseteq 2^{\mathbb{N}}$ and a notation $\mathtt{omega} \in \mathcal{O}$ of order type ω satisfying $P \equiv^1_{<\omega} P^{(\mathtt{omega})}$?*
(4) *Does there exist Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ with $P \leq^1_\omega Q$ such that there is no $|a|$-bounded learnable function $\Gamma : Q \to P$ for any $a \in \mathcal{O}$? For a Π^0_1 set $\widehat{P}$ in Theorem 117, does there exist a function $\Gamma : \widehat{P} \to P$ $(1, \omega)$-computable via an $|a|$-bounded learner for some notation $a \in \mathcal{O}$?*
(5) *Does there exist a pair of special Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ with a function $\Gamma : Q \blacktriangledown P \to Q \oplus P$ (or $\Gamma : Q \triangledown_\infty \bigtriangledown P \to Q \oplus P$) which is learnable by a team of confident learners (or a team of eventually-Popperian learners)?*
(6) *Let P_0, P_1, Q_0, and Q_1 be Π^0_1 subsets of $2^{\mathbb{N}}$ with $Q_0 \leq^1_\omega Q_1$ and $P_0 \leq^1_\omega P_1$. Then, does $P_0 \blacktriangledown^\circ Q_0 \leq^1_\omega P_1 \blacktriangledown^\circ Q_1$ hold? Moreover, if $Q_0 \leq^1_\omega Q_1$ is witnessed by an eventually Lipschitz learner, then does $P_0 \blacktriangledown Q_0 \leq^1_\omega P_1 \blacktriangledown Q_1$ hold?*
(7) *Does there exist Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ such that there is a computable functions $\Gamma_i : Q_i \to P$ with a finite Π^0_1 d-layer $\{Q_i\}_{i<m}$ of Q, but is no computable functions $\Delta_i : Q_i \to P$ with a finite Π^0_1 cover $\{Q_i\}_{i<m}$ of Q? In other words, does there exist Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$ such that $P \leq^1_{<\omega} Q$ and $P \not\leq^{<\omega}_{tt,1} Q$?*
(8) *Determine the strength of the existence of computable functions $\Gamma_i : Q_i \to P$ with a Π^0_1 cover $\{Q_i\}_{i \in \mathbb{N}}$ of Q, for any Π^0_1 sets $P, Q \subseteq 2^{\mathbb{N}}$. In other words, compare the reducibility $\leq^\omega_{tt,1}$ and other reducibility notions (e.g., $\leq^{<\omega}_{tt,1}$, $\leq^1_{<\omega}$, $\leq^1_{\omega|<\omega}$, $\leq^{<\omega}_1$ and $\leq^1_\omega$) for Π^0_1 subsets of Cantor space $2^{\mathbb{N}}$.*

Question 209 (Big Questions)**.**

(1) *Are there elementary differences between any two different degree structures $\mathcal{D}^\alpha_{\beta|\gamma}$ and $\mathcal{D}^{\alpha'}_{\beta'|\gamma'}$ ($\mathcal{P}^\alpha_{\beta|\gamma}$ and $\mathcal{P}^{\alpha'}_{\beta'|\gamma'}$)?*
(2) *Is the commutative concatenation $\triangledown$ first-order definable in the structure $\mathcal{D}^1_1$ or $\mathcal{P}^1_1$?*
(3) *Is each local degree structure $\mathcal{P}^\alpha_{\beta|\gamma}$ first-order definable in the global degree structure $\mathcal{D}^\alpha_{\beta|\gamma}$?*
(4) *Is the structure $\mathcal{P}^1_\omega$ dense?*
(5) *Investigate properties of $(\alpha, \beta|\gamma)$-degrees* **a** *assuring the existence of* $\mathbf{b} > \mathbf{a}$ *which has the same $(\alpha', \beta'|\gamma')$-degree as* **a**.
(6) *Investigate the nested nested model, the nested nested nested model, and so on.*
(7) *Does there exist a "natural" intermediate notion between $(<\omega, \omega)$-computability (i.e., team-learnability) and $(\omega, 1)$-computability (i.e., nonuniform computability) on Π^0_1 sets?*
(8) *(Ishihara) Define a uniform (non-adhoc) interpretation (such as the Kleene realizability interpretation) translating each intuitionistic arithmetical sentence (e.g., $(\neg\neg\exists n \forall m A(n, m)) \to (\exists n \forall m A(n, m))$) into a partial multi-valued function (e.g., Σ^0_2-$\mathsf{DNE} :\subseteq \mathbb{N}^{\mathbb{N}} \rightrightarrows \mathbb{N}^{\mathbb{N}}$).*

Acknowledgements. The authors were partially supported by Grant-in-Aid for JSPS fellows. The second author would like to thank Hajime Ishihara, Dick de Jongh, Arno Pauly, and Albert Visser, for valuable comments and helpful discussion. The second author also would like to thank Makoto Tatsuta and Yoriyuki Yamagata for introducing him to the syntactical study on Limit Computable Mathematics. Finally, the second author also grateful to Sam Sanders who helped his English writing.

REFERENCES

[1] Yohji Akama, Stefano Berardi, Susumu Hayashi, and Ulrich Kohlenbach. An arithmetical hierarchy of the law of excluded middle and related principles. In *19th IEEE Symposium on Logic in Computer Science (LICS 2004), Turku, Finland, Proceedings*, pages 192–201. IEEE Computer Society, 2004.

[2] Christopher P. Alfeld. Non-branching degrees in the Medvedev lattice of Π^0_1 classes. *J. Symb. Log.*, 72(1):81–97, 2007.

[3] Alessandro Andretta. More on Wadge determinacy. *Ann. Pure Appl. Logic*, 144(1-3):2–32, 2006.

[4] Stefano Berardi. Some intuitionistic equivalents of classical principles for degree 2 formulas. *Ann. Pure Appl. Logic*, 139(1-3):185–200, 2006.

[5] Stefano Berardi, Thierry Coquand, and Susumu Hayashi. Games with 1-backtracking. *Ann. Pure Appl. Logic*, 161(10):1254–1269, 2010.

[6] Stefano Berardi and Yoriyuki Yamagata. A sequent calculus for limit computable mathematics. *Ann. Pure Appl. Logic*, 153(1-3):111–126, 2008.

[7] Stephen Binns. A splitting theorem for the Medvedev and Muchnik lattices. *Math. Log. Q.*, 49(4):327–335, 2003.

[8] Stephen Binns and Stephen G. Simpson. Embeddings into the Medvedev and Muchnik lattices of Π^0_1 classes. *Arch. Math. Log.*, 43(3):399–414, 2004.

[9] Lenore Blum and Manuel Blum. Toward a mathematical theory of inductive inference. *Information and Control*, 28(2):125–155, 1975.

[10] Vasco Brattka. Effective Borel measurability and reducibility of functions. *Math. Log. Q.*, 51(1):19–44, 2005.

[11] Vasco Brattka, Matthew de Brecht, and Arno Pauly. Closed choice and a uniform low basis theorem. *Ann. Pure Appl. Logic*, 163(8):986–1008, 2012.

[12] Vasco Brattka and Guido Gherardi. Effective choice and boundedness principles in computable analysis. *Bulletin of Symbolic Logic*, 17(1):73–117, 2011.

[13] Vasco Brattka and Guido Gherardi. Weihrauch degrees, omniscience principles and weak computability. *J. Symb. Log.*, 76(1):143–176, 2011.

[14] Vasco Brattka, Guido Gherardi, and Alberto Marcone. The Bolzano-Weierstrass theorem is the jump of weak König's lemma. *Ann. Pure Appl. Logic*, 163(6):623–655, 2012.

[15] Vasco Brattka and Arno Pauly. Computation with advice. In Xizhong Zheng and Ning Zhong, editors, *Proceedings Seventh International Conference on Computability and Complexity in Analysis*, pages 41–55, 2010.

[16] Vasco Brattka, Arno Pauly, and Stéphane le Roux. Connected choice and Brouwer's fixed point theorem. in preparation, 2011.

[17] J. Case and S. Ngo-Manguelle. Refinements of inductive inference by Popperian machines. Technical Report 152, SUNY Buffalo, Dept. of Computer Science, 1979.

[18] John Case, Matthias Ott, Arun Sharma, and Frank Stephan. Learning to win process-control games watching game-masters. *Inf. Comput.*, 174(1):1–19, 2002.

[19] Douglas Cenzer, Takayuki Kihara, Rebecca Weber, and Guohua Wu. Immunity and non-cupping for closed sets. *Tbilisi Math. J.*, 2:77–94, 2009.

[20] Douglas A. Cenzer and Peter G. Hinman. Density of the Medvedev lattice of Π^0_1 classes. *Arch. Math. Log.*, 42(6):583–600, 2003.

[21] Douglas A. Cenzer and Peter G. Hinman. Degrees of difficulty of generalized r.e. separating classes. *Arch. Math. Log.*, 46(7-8):629–647, 2008.

[22] Douglas A. Cenzer and Jefferey B. Remmel. Π^0_1 classes in mathematics. In *Handbook of Recursive Mathematics, vol* 2, Stud. Logic Found. Math., pages 623–821. Elsevier, 1998.

[23] Thomas Chadzelek and Günter Hotz. Analytic machines. *Theor. Comput. Sci.*, 219(1-2):151–167, 1999.

[24] Joshua A. Cole and Takayuki Kihara. The $\forall\exists$-theory of the effectively closed Medvedev degrees is decidable. *Arch. Math. Log.*, 49(1):1–16, 2010.

[25] Joshua A. Cole and Stephen G. Simpson. Mass problems and hyperarithmeticity. *J. Math. Log.*, 7(2):125–143, 2007.

[26] Matthew de Brecht and Akihiro Yamamoto. Interpreting learners as realizers for Σ^0_2-measurable functions (extended abstract). *Special Interests Group on Fundamental Problems in Artificial Intelligence (SIG-FPAI)*, 74:39–44, 2009.

[27] Matthew de Brecht and Akihiro Yamamoto. Topological properties of concept spaces. *Inf. Comput.*, 208(4):327–340, 2010.

[28] Rodney G. Downey, Noam Greenberg, Carl G. Jockusch Jr., and Kevin G. Milans. Binary subtrees with few labeled paths. *Combinatorica*, 31(3):285–303, 2011.

[29] Rodney G. Downey and Denis R. Hirschfeldt. *Algorithmic Randomness and Complexity*. Theory and Applications of Computability. Springer, 2010. 883 pages.

[30] Jacques Duparc. Wadge hierarchy and Veblen hierarchy Part I: Borel sets of finite rank. *J. Symb. Log.*, 66(1):56–86, 2001.

[31] Y. L. Ershov, S. S. Goncharov, A. Nerode, J. B. Remmel, and V. W. Marek, editors. *Handbook of Recursive Mathematics, Volume 2: Recursive Algebra, Analysis and Combinatorics*. Studies in Logic and the Foundations of Mathematics. North Holland, 1998. 798 pages.

[32] Rusins Freivalds and Carl H. Smith. On the role of procrastination in machine learning. *Inf. Comput.*, 107(2):237–271, 1993.

[33] E. Mark Gold. Limiting recursion. *J. Symb. Log.*, 30(1):28–48, 1965.

[34] Noam Greenberg and Joseph S. Miller. Diagonally non-recursive functions and effective Hausdorff dimension. *Bull. London Math. Soc.*, 43(4):636–654, 2011.

[35] Susumu Hayashi. Mathematics based on learning. In Nicolò Cesa-Bianchi, Masayuki Numao, and Rüdiger Reischuk, editors, *Algorithmic Learning Theory, 13th International Conference, ALT 2002, Lübeck, Germany, Proceedings*, volume 2533 of *Lecture Notes in Computer Science*, pages 7–21. Springer, 2002.

[36] Susumu Hayashi. Mathematics based on incremental learning - excluded middle and inductive inference. *Theor. Comput. Sci.*, 350(1):125–139, 2006.

[37] Peter G. Hinman. A survey of Muchnik and Medvedev degrees. (arXiv:1007.2376), Jul 2010.

[38] Sanjay Jain, Daniel N. Osherson, James S. Royer, and Arun Sharma. *Systems That Learn: An Introduction to Learning Theory*. MIT Press, 1999. 329 pages.

[39] Giorgi Japaridze. Introduction to computability logic. *Ann. Pure Appl. Logic*, 123(1-3):1–99, 2003.

[40] Giorgi Japaridze. Sequential operators in computability logic. *Inf. Comput.*, 206(12):1443–1475, 2008.

[41] Giorgi Japaridze. Toggling operators in computability logic. *Theor. Comput. Sci.*, 412(11):971–1004, 2011.

[42] J. E. Jayne and C. A. Rogers. First level Borel functions and isomorphism. *J. Math. Pure Appl.*, 61:177–205, 1982.

[43] Carl G. Jockusch. Degrees of functions with no fixed points. In *Logic, methodology and philosophy of science, VIII (Moscow, 1987)*, Stud. Logic Found. Math., pages 191–201. North-Holland, Amsterdam, 1989.

[44] Carl G. Jockusch and Robert I. Soare. Degrees of members of Π^0_1 classes. *Pacific J. Math.*, 40(3):605–616, 1972.

[45] Carl G. Jockusch and Robert I. Soare. Π^0_1 classes and degrees of theories. *Trans. Amer. Math. Soc.*, 173:33–56, 1972.

[46] Susanne Kaufmann and Martin Kummer. On a quantitative notion of uniformity. *Fundam. Inform.*, 25(1):59–78, 1996.

[47] Martin Kummer and Matthias Ott. Learning branches and learning to win closed games. In Avrim Blum and Michael Kearns, editors, *Proceedings of the Ninth Annual Conference on Computational Learning Theory*, pages 280–291, 1996.

[48] Andrew E. M. Lewis, Richard A. Shore, and Andrea Sorbi. Topological aspects of the Medvedev lattice. *Arch. Math. Log.*, 50(3-4):319–340, 2011.

[49] Y. T. Medvedev. Degrees of difficulty of the mass problems. *Dokl. Akad. Nauk SSSR*, 104:501–504, 1955. (In Russian).

[50] Yiannis N. Moschovakis. *Descriptive Set Theory*. Mathematical Surveys and Monographs. American Mathematical Society, 2009. 502 pages.

[51] Luca Motto Ros. Game representations of classes of piecewise definable functions. *Math. Log. Q.*, 57(1):95–112, 2011.

[52] A. A. Muchnik. On strong and weak reducibility of algorithmic problems. *Sibirskii Math. Zh.*, 4:1328–1341, 1963. (In Russian).

[53] Carl Mummert. Subsystems of second-order arithmetic between RCA_0 and WKL_0. *Arch. Math. Log.*, 47(3):205–210, 2008.

[54] Masahiro Nakata and Susumu Hayashi. A limiting first order realizability interpretation. *Sci. Math. Jpn.*, 55(3):567–580, 2002.

[55] Andre Nies. *Computability and Randomness*. Oxford Logic Guides. Oxford University Press, 2009. 433 pages.

[56] Matthias Ott and Frank Stephan. Structural measures for games and process control in the branch learning model. *Theor. Comput. Sci.*, 244(1-2):135–165, 2000.

[57] Hartley Rogers. *The Theory of Recursive Functions and Effective Computability*. MIT Press, 1987. 504 pages.

[58] Brian Semmes. Multitape games. In J. Van Benthem, D. Gabbay, and B. Löwe, editors, *Interactive logic*, volume 1 of *Texts in Logic and Games*, pages 195–207. Amsterdam University press, 2007.

[59] Brian Semmes. *A Game for the Borel Functions*. PhD thesis, Universiteit van Amsterdam, 2009.

[60] Paul Shafer. Coding true arithmetic in the Medvedev and Muchnik degrees. *J. Symb. Log.*, 76(1):267–288, 2011.

[61] Stephen G. Simpson. Mass problems and randomness. *Bull. Symb. Log.*, 11(1):1–27, 2005.

[62] Stephen G. Simpson. An extension of the recursively enumerable Turing degrees. *J. London Math. Soc.*, 75:287–297, 2007.

[63] Stephen G. Simpson. Some fundamental issues concerning degrees of unsolvability. In *Computational prospects of infinity Part II. Presented talks*, volume 15 of *Lect. Notes Ser. Inst. Math. Sci. Natl. Univ. Singap.*, pages 313–332. World Sci. Publ., Hackensack, NJ, 2007.

[64] Stephen G. Simpson. *Subsystems of Second Order Arithmetic*. Perspectives in Logic. Cambridge University Press, 2009. 464 pages.
[65] Stephen G. Simpson. Mass problems associated with effectively closed sets. *Tohoku Math. J.*, 63(4):489–517, 2011.
[66] Stephen G. Simpson and Theodore Slaman. Medvedev degrees of Π^0_1 subsets of 2^ω. unpublished, 2001.
[67] Robert I. Soare. *Recursively Enumerable Sets and Degrees*. Perspectives in Mathematical Logic. Springer, Heidelberg, 1987. XVIII+437 pages.
[68] Andrea Sorbi. Some remarks on the algebraic structure of the Medvedev lattice. *J. Symb. Log.*, 55(2):831–853, 1990.
[69] Andrea Sorbi and Sebastiaan Terwijn. Intermediate logics and factors of the Medvedev lattice. *Ann. Pure Appl. Logic*, 155(2):69–85, 2008.
[70] Frank Stephan, Yue Yang, and Liang Yu. Turing degrees and the Ershov hierarchy. In T. Arai, J. Brendle, C. T. Chong, R. Downey, Q. Feng, H. Kikyo, and H.Ono, editors, *Proceedings of the 10th Asian Logic Conference*, pages 300–321, 2010.
[71] Makoto Tatsuta and Stefano Berardi. Non-commutative infinitary Peano arithmetic. In *CSL*, pages 538–552, 2011.
[72] Michael Toftdal. A calibration of ineffective theorems of analysis in a hierarchy of semi-classical logical principles: (extended abstract). In *ICALP*, pages 1188–1200, 2004.
[73] R. A. Van Wesep. *Subsystems of Second-Order Arithmetic and Descriptive Set Theory under the Axiom of Determinateness*. PhD thesis, University of California, Barkeley, 1977.
[74] W. W. Wadge. *Reducibility and determinateness on the Baire space*. PhD thesis, University of California, Barkeley, 1983.
[75] Klaus Weihrauch. *Computable Analysis: An Introduction*. Texts in Theoretical Computer Science. Springer, 2000. 285 pages.
[76] Mariko Yasugi, Takakazu Mori, and Yoshiki Tsujii. The effective sequence of uniformities and its limit as a methodology in computable analysis. *Ann. Japan Assoc. Phil. Sci.*, 15(2):99–121, 2007.
[77] Mariko Yasugi and Yoshiki Tsujii. Computability of a function with jumps –effective uniformity and limiting recursion–. *Topol. and its Appl.*, 146-147:563–582, 2005.
[78] Martin Ziegler. Real hypercomputation and continuity. *Theory Comput. Syst.*, 41(1):177–206, 2007.
[79] Martin Ziegler. Revising type-2 computation and degrees of discontinuity. *Electr. Notes Theor. Comput. Sci.*, 167:255–274, 2007.
[80] Martin Ziegler. (short) survey of real hypercomputation. In *Computation and Logic in the Real World, Third Conference on Computability in Europe, CiE 2007, Siena, Italy, June 18-23, 2007, Proceedings*, pages 809–824, 2007.
[81] Martin Ziegler. Real computation with least discrete advice: A complexity theory of nonuniform computability with applications to effective linear algebra. *Ann. Pure Appl. Logic*, 163(8):1108–1139, 2012.

(K. Higuchi) Mathematical Institute, Tohoku University, Sendai 980-8578, Japan
E-mail address: `sa7m24@math.tohoku.ac.jp`

(T. Kihara) School of Information Science, Japan Advanced Institute of Science and Technology, Nomi 923-1292, Japan
E-mail address: `kihara@jaist.ac.jp`